

Theoretical Studies and Computer Modeling of Supramolecular Chemical Systems: Structure, Properties and Reactivity [†]

 Alexander S. Novikov ^{1,2,3} 
¹ Institute of Chemistry, Saint Petersburg State University, Universitetskaya Nab., 7/9, 199034 Saint Petersburg, Russia; a.s.novikov@spbu.ru

² Infochemistry Scientific Center, ITMO University, Kronverksky Pr., 49, Bldg. A, 197101 Saint Petersburg, Russia

³ Research Institute of Chemistry, Peoples' Friendship University of Russia (RUDN University), Miklukho-Maklaya St., 6, 117198 Moscow, Russia

[†] Presented at the 26th International Electronic Conference on Synthetic Organic Chemistry, 1–15 November 2022; Available online: <https://sciforum.net/event/ecsoc-26>.

Abstract: The results of my research in the fields of theoretical studies and computer modeling of supramolecular chemical systems were presented. The main attention was focused on theoretical studies in the following topics: cycloaddition and nucleophilic addition reactions involving substrates with multiple CC and CN bonds, their mechanisms, driving forces, kinetics and thermodynamics; the consideration of the catalysis of hydrocarbons oxidation processes and their conversion to alcohols, ethers, aldehydes, ketones and carboxylic acids; investigations of various unusual types of non-covalent interactions (from quite trivial hydrogen bonds to more exotic σ -hole, π -hole and metal-philic interactions) in organic, organometallic and coordination compounds. Some fundamental issues of supramolecular chemical systems were also discussed (e.g., structure and properties of chemical compounds and their supramolecular associates; conformational transitions and rotation barriers of functional groups; nature of chemical bonds; orbital and charge factors; photophysical properties).

Keywords: coordination chemistry; organometallic chemistry; inorganic chemistry; organic chemistry; quantum chemical calculations; computer modeling; supramolecular systems; catalysis; reactivity



Citation: Novikov, A.S. Theoretical Studies and Computer Modeling of Supramolecular Chemical Systems: Structure, Properties and Reactivity. *Chem. Proc.* **2022**, *12*, 88. <https://doi.org/10.3390/ecsoc-26-13717>

Academic Editor: Julio A. Seijas

Published: 18 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

In this Keynote presentation, I have presented the results of my research in fields of theoretical studies and computer modeling of supramolecular chemical systems. The main attention was focused on theoretical studies in the following topics: cycloaddition and nucleophilic-addition reactions involving substrates with multiple CC and CN bonds (viz. metal-assisted (by Au(I), Au(III), Pt(II), Pt(IV), Re(V) and Pd(II)) 1,3-dipolar cycloaddition of aldo- and ketonitrone to isocyanides [1]; Cu(I)-catalyzed 1,3-dipolar cycloaddition of ketonitrone to dialkylcyanamides [2]; nucleophilic addition of hydroxyguanidines, amidoximes and ketoximes to nitrilium *closo*-decaborate clusters [3]; nucleophilic addition of amidoximes to aromatic nitrile complexes *trans*-[PtCl₂(RC₆H₄CN)₂] (R = *p*-CF₃, H, *o*-Cl) [4]), their mechanisms, driving forces, kinetics and thermodynamics and correlations between activation and reaction energies and the calculated unscaled stretching vibration frequencies of CN group in metal-free and coordinated isocyanides were also discussed [5]; the consideration of the catalysis of hydrocarbons oxidation processes and their conversion to alcohols, ethers, aldehydes, ketones and carboxylic acids (mechanisms of the rate-limiting stages of alkanes oxidation and olefins epoxidation with hydrogen peroxide assisted by the formation of hydroxyl radicals, catalyzed by aqua complexes [M(H₂O)_n]³⁺ of the group III metals exhibiting a unique stable non-zero oxidation state (M = Ga, In, Sc, Y or La) [6]); investigations of various unusual types of non-covalent interactions (from

quite trivial hydrogen bonds to more exotic σ -hole, π -hole and metallophilic interactions) in organic, organometallic and coordination compounds (viz. intermolecular hydrogen bonds and anagostic interactions in Pd(II) acyclic diaminocarbene complexes [7]; supramolecular polymers derived from Pd(II) and Pt(II) N,N-diaryldiazabutadiene Schiff base complexes via C(sp²)-H...Hal hydrogen bonding [8]; halogen bonds with metal centers [9] and isocyano carbon atom [10]; non-covalent B-H... π (Ph) interactions determines stabilization of the configuration around the amidrazone C=N bond in *closo*-decaborato amidrazones [11]; the effect of halogen bonding on solubility of Pd/Pt isocyanide complexes [12]; chalcogen bonds in Pd(II) carbene complexes [13]; stacking interactions and π -hole interactions in transition metal complexes and various supramolecular building blocks [14]; ligation-enhanced π -hole... π , π -hole...metal center, and π -hole...lone pair interactions involving isocyanides [15]; reverse-arene sandwich structures based upon π -hole...[M(II)] (d⁸ M = Pt, Pd) interactions intramolecular [16]; xenon derivatives as aerogen bond-donating catalysts for organic transformations [17]; sulfonium and selenonium salts as noncovalent organocatalysts for the multicomponent Groebke–Blackburn–Bienaymé reaction [18]). Some fundamental issues of supramolecular chemical systems were also discussed (e.g., structure and properties of chemical compounds and their supramolecular associates (viz. comprehensive theoretical study of the aminonitrone-imino hydroxamic acid tautomerism [19]; melamine–barbiturate supramolecular assembly as a pH-dependent organic radical trap material [20]); conformational transitions and rotation barriers of functional groups; nature of chemical bonds; orbital and charge factors; photophysical properties of phosphorescent Pt(II) complexes featuring chelated acetoxime pyrazoles [21]). This research is interdisciplinary in nature and lies at the intersection of computer modeling with such natural science disciplines as chemistry, physics, crystallography, biology and medicine, as well as directly related to materials science and nanotechnology. The results obtained in the course of this research can be promising for finding new original ways to create high-tech functional materials with valuable mechanical, magnetic and optical properties; porous structures with a developed surface, sensors, battery cells and liquid crystals; synthesis of antibacterial, antiviral and antineoplastic drugs; labels for neutron capture therapy of oncological diseases; catalysts for cross-coupling processes (for example, the Suzuki, Heck and Sonogashira reactions) and the conversion of hydrocarbons (for example, to alcohols, epoxides, ethers, aldehydes, ketones and carboxylic acids), as well as multicomponent reactions.

Funding: The support of my research focused on theoretical studies and computer modeling of supramolecular chemical systems within the RUDN University Strategic Academic Leadership Program is gratefully acknowledged.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Novikov, A.S.; Kuznetsov, M.L.; Pombeiro, A.J.L. Theory of the formation and decomposition of N-heterocyclic amino oxycarbenes through metal-assisted [2+3]-dipolar cycloaddition/retro-cycloaddition. *Chem. Eur. J.* **2013**, *19*, 2874–2888. [[CrossRef](#)] [[PubMed](#)]
2. Melekhova, A.A.; Smirnov, A.S.; Novikov, A.S.; Panikorovskii, T.L.; Bokach, N.A.; Kukushkin, V.Y. Copper(I)-catalyzed 1,3-dipolar cycloaddition of ketonitrone to dialkylcyanamides. A step toward sustainable generation of 2,3-dihydro-1,2,4-oxadiazoles. *ACS Omega* **2017**, *2*, 1380–1391. [[CrossRef](#)] [[PubMed](#)]
3. Bolotin, D.S.; Burianova, V.K.; Novikov, A.S.; Demakova, M.Y.; Pretorius, C.; Mokolokolo, P.P.; Roodt, A.; Bokach, N.A.; Suslonov, V.V.; Zhdanov, A.P.; et al. Nucleophilicity of oximes based upon addition to a nitrilium *closo*-decaborate cluster. *Organometallics* **2016**, *35*, 3612–3623. [[CrossRef](#)]
4. Bolotin, D.S.; Demakova, M.Y.a.; Novikov, A.S.; Avdontceva, M.S.; Kuznetsov, M.L.; Bokach, N.A.; Kukushkin, V.Y. Bifunctional reactivity of amidoximes observed upon nucleophilic addition to metal-activated nitriles. *Inorg. Chem.* **2015**, *54*, 4039–4046. [[CrossRef](#)] [[PubMed](#)]

5. Novikov, A.S. 1,3-Dipolar cycloaddition of nitrones to transition metal-bound isocyanides: DFT and HSAB principle theoretical model together with analysis of vibrational spectra. *J. Organomet. Chem.* **2015**, *797*, 8–12. [\[CrossRef\]](#)
6. Novikov, A.S.; Kuznetsov, M.L.; Pombeiro, A.J.L.; Bokach, N.A.; Shul'pin, G.B. Generation of HO• radical from hydrogen peroxide catalyzed by aqua-complexes of the group III metals $[M(H_2O)_n]^{3+}$ (M = Ga, In, Sc, Y, La): A theoretical study. *ACS Catal.* **2013**, *3*, 1195–1208. [\[CrossRef\]](#)
7. Mikhaylov, V.N.; Sorokoumov, V.N.; Korvinson, K.A.; Novikov, A.S.; Balova, I.A. Synthesis and simple immobilization of palladium(II) acyclic diaminocarbene complexes on polystyrene-support as efficient catalysts for Sonogashira and Suzuki-Miyaura cross-coupling. *Organometallics* **2016**, *35*, 1684–1697. [\[CrossRef\]](#)
8. Tskhovrebov, A.G.; Novikov, A.S.; Odintsova, O.V.; Mikhaylov, V.N.; Sorokoumov, V.N.; Serebryanskaya, T.V.; Starova, G.L. Supramolecular polymers derived from the Pt^{II} and Pd^{II} Schiff base complexes via C(sp²)-H...Hal hydrogen bonding: Combined experimental and theoretical study. *J. Organomet. Chem.* **2019**, *886*, 71–75. [\[CrossRef\]](#)
9. Ivanov, D.M.; Novikov, A.S.; Ananyev, I.V.; Kirina, Y.V.; Kukushkin, V.Y. Halogen bonding between metal centers and halocarbons. *Chem. Commun.* **2016**, *52*, 5565–5568. [\[CrossRef\]](#) [\[PubMed\]](#)
10. Mikherdov, A.S.; Katkova, S.A.; Novikov, A.S.; Efremova, M.M.; Reutskaya, E.Y.; Kinzhalov, M.A. (Isocyano group)...lone pair interactions involving coordinated isocyanides: Experimental, theoretical and CSD study. *CrystEngComm* **2020**, *22*, 1154–1159. [\[CrossRef\]](#)
11. Burianova, V.K.; Bolotin, D.S.; Mikherdov, A.S.; Novikov, A.S.; Mokolokolo, P.P.; Roodt, A.; Boyarskiy, V.P.; Dar'in, D.; Krasavin, M.; Suslonov, V.V.; et al. Mechanism of generation of closo-decaborato amidrazones. Intramolecular non-covalent B-H... π (Ph) interaction determines stabilization of the configuration around the amidrazone C=N bond. *New J. Chem.* **2018**, *42*, 8693–8703. [\[CrossRef\]](#)
12. Kinzhalov, M.A.; Kashina, M.V.; Mikherdov, A.S.; Mozheeva, E.A.; Novikov, A.S.; Smirnov, A.S.; Ivanov, D.M.; Kryukova, M.A.; Ivanov, A.Y.; Smirnov, S.N.; et al. Dramatically enhanced solubility of halide-containing organometallic species in diiodomethane: The role of solvent complex halogen bonding. *Angew. Chem. Int. Ed.* **2018**, *57*, 12785–12789. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Mikherdov, A.S.; Kinzhalov, M.A.; Novikov, A.S.; Boyarskiy, V.P.; Boyarskaya, I.A.; Dar'in, D.V.; Starova, G.L.; Kukushkin, V.Y. Difference in energy between two distinct types of chalcogen bonds drives regioisomerization of binuclear (diaminocarbene)Pd^{II} complexes. *J. Am. Chem. Soc.* **2016**, *138*, 14129–14137. [\[CrossRef\]](#)
14. Ivanov, D.M.; Kirina, Y.V.; Novikov, A.S.; Starova, G.L.; Kukushkin, V.Y. Efficient π -stacking with benzene provides 2D assembly of *trans*-[PtCl₂(*p*-CF₃C₆H₄CN)₂]. *J. Mol. Struct.* **2016**, *1104*, 19–23. [\[CrossRef\]](#)
15. Mikherdov, A.S.; Novikov, A.S.; Kinzhalov, M.A.; Boyarskiy, V.P.; Starova, G.L.; Ivanov, A.Y.; Kukushkin, V.Y. Halides held by bifurcated chalcogen-hydrogen bonds. Effect of μ (S,N-H)Cl contacts on dimerization of Cl(Carbene)Pd^{II} species. *Inorg. Chem.* **2018**, *57*, 3420–3433. [\[CrossRef\]](#)
16. Rozhkov, A.V.; Krykova, M.A.; Ivanov, D.M.; Novikov, A.S.; Sinelshchikova, A.A.; Volostnykh, M.V.; Konovalov, M.A.; Grigoriev, M.S.; Gorbunova, Y.G.; Kukushkin, V.Y. Reverse arene sandwich structures based upon π -hole...[M^{II}](d⁸M = Pt, Pd) interactions, where positively charged metal centers play the role of a nucleophile. *Angew. Chem. Int. Ed.* **2019**, *58*, 4164–4168. [\[CrossRef\]](#)
17. Novikov, A.S.; Bolotin, D.S. Xenon derivatives as aerogen bond-donating catalysts for organic transformations: A theoretical study on the metaphorical “spherical cow in a vacuum” provides insights into noncovalent organocatalysis. *J. Org. Chem.* **2023**, *88*, 1936–1944. [\[CrossRef\]](#)
18. Il'in, M.V.; Novikov, A.S.; Bolotin, D.S. Sulfonium and selenonium salts as noncovalent organocatalysts for the multicomponent Groebke-Blackburn-Bienaymé reaction. *J. Org. Chem.* **2022**, *87*, 10199–10207. [\[CrossRef\]](#)
19. Il'in, M.V.; Novikov, A.S.; Bolotin, D.S. Aminonitrone-iminohydroxamic acid tautomerism: Theoretical and spectroscopic study. *J. Mol. Struct.* **2019**, *1176*, 759–765. [\[CrossRef\]](#)
20. Shilovskikh, V.V.; Timralieva, A.A.; Nesterov, P.V.; Novikov, A.S.; Sitnikov, P.A.; Konstantinova, E.A.; Kokorin, A.I.; Skorb, E.V. Melamine-barbiturate supramolecular assembly as a pH-dependent organic radical trap material. *Chem. A Eur. J.* **2020**, *26*, 16603–16610. [\[CrossRef\]](#)
21. Bolotin, D.S.; Novikov, A.S.; Kolesnikov, I.E.; Suslonov, V.V.; Novozhilov, Y.; Ronzhina, O.; Dorogov, M.; Krasavin, M.; Kukushkin, V.Y. Phosphorescent platinum(II) complexes featuring chelated acetoxime pyrazoles: Synthetic, structural, and photophysical study. *ChemistrySelect* **2016**, *3*, 456–461. [\[CrossRef\]](#)