

An In Silico Approach to Enzymatic Synthesis of Fucooligosaccharides Using α -L-Fucosidase from *Thermotoga maritima*[†]

Emmanuel Pérez-Escalante¹, Luis Guillermo González-Olivares¹, Araceli Castañeda-Ovando¹, Alma Elizabeth Cruz-Guerrero², John F. Trant³, Wendolyne López-Orozco¹, Luis Humberto Mendoza-Huizar^{1,*} and Sergio Alatorre-Santamaría^{2,*}

¹ Área Académica de Química, Universidad Autónoma del Estado de Hidalgo, Carretera Pachuca-Tulancingo km 4.5 Mineral de la Reforma, Hidalgo 42184, Mexico; emmanuel_perez@uaeh.edu.mx (E.P.-E.); lgonzales@uaeh.edu.mx (L.G.G.-O.); ovandoa@uaeh.edu.mx (A.C.-O.); wen.lopez.orocho@gmail.com (W.L.-O.)

² Departamento de Biotecnología, División de Ciencias Biológicas y de la Salud, Universidad Autónoma Metropolitana, Unidad Iztapalapa, Av. San Rafael Atlixco 186, Ciudad de Mexico 09340, Mexico; aec@xanum.uam.mx

³ Department of Chemistry and Biochemistry, University of Windsor, 401 Sunset Ave., Windsor, ON N9B 3P4, Canada; j.trant@uwindsor.ca

* Correspondence: hhuizar@uaeh.edu.mx (L.H.M.-H.); salatorre@xanum.uam.mx (S.A.-S.)

[†] Presented at the 24th International Electronic Conference on Synthetic Organic Chemistry, 15 November–15 December 2020; Available online: <https://ecsoc-24.sciforum.net/> (accessed on 14 November 2020).

Citation: Pérez-Escalante, E.; González-Olivares, L.G.; Castañeda-Ovando, A.; Cruz-Guerrero, A.E.; Trant, J.F.; López-Orozco, W.; Mendoza-Huizar, L.H.; Alatorre-Santamaría, S. An In Silico Approach to Enzymatic Synthesis of Fucooligosaccharides Using α -L-Fucosidase from *Thermotoga maritima*. **2021**, *3*, 10. <https://doi.org/10.3390/ecsoc-24-08303>

Published: 14 November 2020

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Abstract: Fucooligosaccharides comprise the primary group of human milk oligosaccharides. Due to their beneficial properties, a series of synthetic methods have been proposed to obtain them. Enzymatic methods show great promise, and α -L-fucosidase from *Thermotoga maritima* has emerged as a powerful catalyst for their production. Nonetheless, the enzyme's limited substrate scope has delayed its wider application. The present work aims to compare the relative reactivity of fucose, pNP-fucose, and ethyl-fucose, while also exploring the molecular interactions of these fucosyl-donors with the enzyme through a combination DFT and docking analysis. The HOMO-LUMO band gaps range from -7.14571 to -4.24429 eV, with α/β -pNP-fucose and α -fucose being the three most reactive compounds. Moderate association energies between -6.4 to -5.5 kcal·mol⁻¹ were found in the docking analysis, with α -pNP-fucose and both anomers of ethyl-fucose demonstrating the poorest affinity. In the case of α/β -lactose affinity to the β -fucose/enzyme complex, no significant differences were shown. We conclude that the best fucosyl-donors for transfucosylation are those that maintain an enzyme affinity and reactivity similar to pNP-fucose.

Keywords: Fucooligosaccharides; α -L-fucosidase; DFT study; molecular docking

1. Introduction

Fucooligosaccharides (FucOS) are the main oligosaccharides in human milk comprising 65–77% of the total oligosaccharide content [1]. Due to their antimicrobial, immunomodulatory, and prebiotic activities, as well as their promise to function as developmental cognitive enhancers, their incorporation into commercial formulations has become highly desirable [1,2]. As their isolation is complex due to their low abundance in animal milk, attention has turned to synthesis [1,2]. Fermentation is the most efficient, with generally recognize as safe (GRAS) certification from the U.S. Food and Drug Administration (FDA) permitting the addition of 2'-fucosyllactose (2'FL) to infant formula [2]. Another synthetic alternative with recent promising results is the use of isolated en-

zyme, which requires the yield optimization of FucOS by fucosyl-transferases. Unfortunately, this approach presents the inconvenience of requiring nucleotide sugars as fucosyl-donors, which are more expensive than those used for fucosidases [1,3]. Consequently, FucOS synthesis by fucosyl-hydrolases like the α -L-fucosidase from *Thermotoga maritima* has gained importance, as this pathway allows the use either of less expensive fucosyl-donors or even agro-industrial waste [3–8]. However, this enzymatic route provides lower yields than the transferase, or involves the release of toxic compounds such as *p*-nitrophenol [5–8]. Alternatives are highly desirable. Thus, the present work aimed to determine the relative reactivity of three non-classical fucosyl-donors through an in silico study to propose substrate alternatives for the enzymatic synthesis of FucOS.

2. Methods

2.1. Geometry Optimization and HOMO-LUMO Parameters

All compounds were totally geometry optimized through the Density Functional Theory (DFT) with the B3LYP/6-311++G(2d,2p) basis set using water as solvent. The highest occupied molecular orbital (HOMO) and lowest unoccupied molecular orbital (LUMO) density surfaces were visualized with Gabedit 2.5.0. [9].

2.2. Molecular Docking for Hydrolysis and Transfucosylation Process

The A chain of the α -L-fucosidase crystal from *Thermotoga maritima* (PDB: 1ODU) was prepared with the DockPrep tool implemented in Chimera 1.13.1 [10], prior to its use as the receptor for molecular docking. All molecular dockings were performed by Autodock VINA [11] through the PyRx software [12], taking as reference the amino acids from the active site cited by Sulzenbacher et al. [13], with coordinates for the search space centered on x: -20.63, y: 19.03, and z: 63.32, with a grid cube with dimensions of 25.00 Å. In the case of the hydrolysis, a single docking step was performed for each fucosyl-donor and the receptor. Meanwhile for the transfucosylation process, a sequential docking sequence was employed. First, β -fucose was docked to the enzyme in order to form the pre-complex, then docking was performed again with lactose. The best binding mode for each interaction was obtained and its interactions were processed with the BIOVIA Discovery Studio Visualizer© v19.1.0.18287 [14].

3. Results and Discussion

3.1. HOMO-LUMO Parameters

The HOMO-LUMO frontier orbitals and the bandgaps were calculated for both anomers of fucose, ethyl-fucose, *p*-nitrophenyl (pNP)-fucose, and lactose (Figure 1). In general, β -anomers showed a lower bandgap compared with the α -anomers, except for the pNP-fucose anomers, which showed similar bandgaps. As the magnitude of the HOMO-LUMO gap directly relates to chemical reactivity, where larger bandgaps predict lower reactivity [15,16], we predict that the β -anomers of fucose, ethyl-fucose, and lactose should prove less reactive than their α -anomers; in contrast, both anomers of pNP-fucose should demonstrate similar reactivity. Ranking the series, α/β -pNP-fucose with its electron withdrawing nature, unsurprisingly, is predicted to be the most reactive followed by α -fucose. In general, the α -anomers promise greater reactivity than the β anomers (again, with the exception of the hot pNP electrophile) based on the distribution of density in the frontier molecular orbitals.




Figure 1. HOMO and LUMO surfaces (blue: positive and red: negative) of (a) α -fucose, (b) α -ethyl-fucose, (c) α -pNP-fucose, (d) α -lactose, (e) β -fucose, (f) β -ethyl-fucose, (g) β -pNP-fucose, and (h) β -lactose, as well as their HOMO-LUMO band gap.


3.2. Molecular Docking for Hydrolysis Processes

Single molecular docking simulated the hydrolysis catalyzed by *T. maritima*'s α -L-fucosidase. Docking scores varied between -6.4 and -5.5 kcal·mol $^{-1}$, with β -pNP-fucose showing the best affinity to the receptor, while α -ethyl-fucose presented the worst (Table 1). A lower substrate–enzyme affinity could be correlated with lower complex stability and a tendency to destroy it [17]. Thus, among the six different tested molecules, α -ethyl-fucose, β -ethyl-fucose, and α -pNP-fucose appear to be the fucosyl donors most readily hydrolyzed once bound to the enzyme.

Table 1. Coupling energies obtained for each fucosyl-donor and the α -L-fucosidase from *Thermotoga maritima*.

Fucosyl-Donor	Coupling Energy (kcal·mol ⁻¹)
α -fucose	−6.0
β -fucose	−6.3
α -ethyl-fucose	−5.5
β -ethyl-fucose	−5.8
α -pNP-fucose	−5.9
β -pNP-fucose	−6.4

The key non-covalent interactions were identified (Figure 2). Those identified for β -fucose/enzyme docking (Figure 2d), agree with those reported by Sulzenbacher et al. [13]. All showed essential π -interactions between the C-5 methyl group and aromatic residues on the enzyme. The last interactions propitiate that the sugar ring takes a perpendicular position to this hydrophobic region, favoring van der Waals interactions observed with the rest of the molecule sites [13]. An important change was found for the interaction with the Asp224, the amino acid responsible for the nucleophilic attack that forms the covalent glycosyl-intermediate [13]. This pre-reaction interaction is a hydrogen-bond for β -fucose; but is van der Waals type for α -fucose and α/β -pNP-fucose; is an unfavorable repulsive interaction for α -ethyl-fucose; and a hydrogen bond with the C-4 OH for β -ethyl-fucose, indicating the different binding modes of these two substrates. These insights could explain the differential docking energies and suggests that hydrolysis would prove more challenging.

**Figure 2.** Mappings in 2D of the binding interactions among fucosyl-donors and the α -L-fucosidase from *Thermotoga maritima*: (a) α -fucose, (b) α -ethyl-fucose, (c) α -pNP-fucose, (d) β -fucose, (e) β -ethyl-fucose, and (f) β -pNP-fucose.

3.3. Molecular Docking for Studying Transfucosylation

In order to simulate a transfucosylation process, a sequential docking was performed. First, the β -fucose/enzyme complex was established by a single docking. The

results were consistent with the reports of Sulzenbacher et al. [13], with α - or β -lactose, providing docking scores of -5.7 and -5.8 kcal·mol $^{-1}$, respectively. However, the conformation was different for each (Figure 3). While α -lactose adopts a position near the β -fucose binding site, β -lactose adheres preferentially to a more distal site. β -lactose has only a series of weak van der Waals interactions holding it in place, including with the key Glu266 residue, while α -lactose forms strong hydrogen bonds with both Glu66 and Glu266, the amino acids responsible for the activation of acceptor groups for transufucosylation [13]. This suggests that the alpha anomer should be far more reactive, and indicates that mutarotation likely forms this anomer prior to transformation.




Figure 3. Binding position of α -lactose (green colored) and β -lactose (blue colored) in the complex β -fucose (orange colored)/enzyme, as well as the molecular interactions of (a) α -lactose and (b) β -lactose to the receptor.

On the other hand, the effect of the fucosyl-donors on the transufucosylation process can be related to the reactivity showed in the HOMO-LUMO gap, because previous *in vitro* studies have shown the effective transference of pNP-fucose to lactose to synthesize FucOS [5,6], while other studies have found low yields or long process when fucose itself is used as the donor [7,8]. Thus, ethyl-fucose could show similar results to those obtained with fucose mainly due to both molecules showing similar reactivity. Finally, according to *in vitro* results obtained earlier and the *in silico* insights found here, it is possible to hypothesize that fucosyl-donors with similar structure and/or reactivity to that of pNP-fucose could act as good substrates for transufucosylation with the α -L-fucosidase from *Thermotoga maritima*.

4. Conclusions

In silico insights of the reactivity and molecular interactions with the α -L-fucosidase from *Thermotoga maritima* obtained for each fucosyl-donor and *in vitro* results from literature allowed us to conclude that the best fucosyl-donors for transufucosylation reaction will be those with similar reactivity to pNP-fucose, so that the synthesis of compounds

with similar structures, but lower toxicity, should be prioritized to find next generation fucosyl transfer agents.

Author Contributions: Conceptualization, E.P.-E., L.G.G.-O. and S.A.-S.; methodology, E.P.-E. and L.H.M.-H.; software, L.H.M.-H. and W.L.-O.; validation, A.E.C.-G., J.F.T. and A.C.-O.; formal analysis, A.E.C.-G. and L.G.G.-O.; investigation, E.P.-E. and L.H.M.-H.; resources, L.H.M.-H., L.G.G.-O. and S.A.-S.; data curation, J.F.T. and A.C.-O.; writing—original draft preparation, E.P.-E.; writing—review and editing, S.A.-S. and J.F.T.; visualization, L.G.G.-O. and S.A.-S.; supervision, L.H.M.-H. and S.A.-S.; project administration, L.G.G.-O. and S.A.-S.; funding acquisition, L.H.M.-H., L.G.G.-O. and S.A.-S. All authors have read and agreed to the published version of the manuscript.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available within the article.

Acknowledgments: E.P.-E. would like to acknowledge the Consejo Nacional de Ciencia y Tecnología (CONACYT) for granting the scholarship number 473399.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Pérez-Escalante, E.; Alatorre-Santamaría, S.; Castañeda-Ovando, A.; Salazar-Pereda, V.; Bautista-Ávila, M.; Cruz-Guerrero, A.E.; Flores-Aguilar, J.F.; González-Olivares, L.G. Human milk oligosaccharides as bioactive compounds in infant formula: Recent advances and trends in synthetic methods. *Crit. Rev. Food Sci. Nutr.* **2020**, *1*, 1–34.
- Bych, K.; Mikš, M.H.; Johanson, T.; Hederos, M.J.; Vignæs, L.K.; Becker, P. Production of HMOs using microbial hosts—From cell engineering to large scale production. *Curr. Opin. Biotechnol.* **2019**, *56*, 130–137.
- Zeuner, B.; Meyer, A.S. Enzymatic transufucosylation for synthesis of human milk oligosaccharides. *Carbohydr. Res.* **2020**, *493*, 108029.
- Wan, L.; Zhu, Y.; Zhang, W.; Mu, W. α -L-Fucosidases and their applications for the production of fucosylated human milk oligosaccharides. *Appl. Microbiol. Biotechnol.* **2020**, *104*, 5619–5631.
- Guzmán-Rodríguez, F.; Alatorre-Santamaría, S.; Gómez-Ruiz, L.; Rodríguez-Serrano, G.; García-Garibay, M.; Cruz-Guerrero, A. Synthesis of a fucosylated trisaccharide via transglycosylation by α -L-fucosidase from *Thermotoga maritima*. *Appl. Biochem. Biotechnol.* **2018**, *186*, 681–691.
- Shi, R.; Ma, J.; Yan, Q.; Yang, S.; Fan, Z.; Jiang, Z. Biochemical characterization of a novel α -L-fucosidase from *Pedobacter* sp. and its application in synthesis of 3'-fucosyllactose and 2'-fucosyllactose. *Appl. Microbiol. Biotechnol.* **2020**, *104*, 5813–5826.
- Uvalampi, A.; Medrano, M.R.; Maaheimo, H.; Salminen, H.; Tossavainen, O.; Frey, A.D. Production and characterization of *Aspergillus niger* GH29 family α -fucosidase and production of a novel non-reducing 1-fucosyllactose. *Glycoconj. J.* **2020**, *37*, 221–229.
- Zeuner, B.; Muschiol, J.; Holck, J.; Lezyk, M.; Gedde, M.R.; Jers, C.; Mikkelsen, J.D.; Meyer, A.S. Substrate specificity and transufucosylation activity of GH29 α -L-fucosidases for enzymatic production of human milk oligosaccharides. *New Biotechnol.* **2018**, *41*, 34–45.
- Allouche, A.R. Gabedit—A graphical user interface for computational chemistry softwares. *J. Comput. Chem.* **2011**, *32*, 174–182.
- Pettersen, E.F.; Goddard, T.D.; Huang, C.C.; Couch, G.S.; Greenblatt, D.M.; Meng, E.C.; Ferrin, T.E. UCSF Chimera—A visualization system for exploratory research and analysis. *J. Comput. Chem.* **2004**, *13*, 1605–1612.
- Trott, O.; Olson, A.J. AutoDock Vina: Improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *J. Comput. Chem.* **2010**, *31*, 455–461.
- Dallakyan, S.; Olson, A.J. Small-molecule library screening by docking with PyRx. *Methods Mol. Biol.* **2015**, *1263*, 243–250.
- Sulzenbacher, G.; Bignon, C.; Nishimura, T.; Tarling, C.A.; Withers, S.G.; Henrissat, B.; Bourne, Y. Crystal Structure of *Thermotoga maritima* α -L-Fucosidase insights into the catalytic mechanism and the molecular basis for fucosidosis. *J. Biol. Chem.* **2004**, *279*, 13119–13128.
- Dassault Systèmes BIOVIA. *Discovery Studio Visualizer*; v19.1.0.18287; Dassault Systèmes: San Diego, CA, USA, 2019.
- Talmaciu, M.M.; Bodoki, E.; Oprean, R. Global chemical reactivity parameters for several chiral beta-blockers from the Density Functional Theory viewpoint. *Clujul Med.* **2016**, *89*, 513–518.
- Boukli-Hacene, F.; Merad, M.; Ghalem, S.; Soufi, W. DFT Study of the Interaction of Cu (II), Zn (II), Sn (II) with Carbohydrates in Aqueous Solution. *J. Chem.* **2014**, *8*, 1009–1017.
- Bettelheim, F.A.; Brown, W.H.; Campbell, M.K.; Farrell, S.O. *Introduction to Organic and Biochemistry*, 7th ed.; Cengage Learning: Belmont, CA, USA, 2010; pp. 353–355.