

Proceeding Paper

# Synthesis and Characterization of a Water-Soluble Pentamethine Indocyanine Dye for Peptide Labeling †

Cátia D. F. Martins , Maria Manuela M. Raposo  and Susana P. G. Costa \* 

Centre of Chemistry, University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal; catiadf\_martins@hotmail.com (C.D.F.M.); mfox@quimica.uminho.pt (M.M.M.R.)

\* Correspondence: spc@quimica.uminho.pt

† Presented at the 25th International Electronic Conference on Synthetic Organic Chemistry, 15–30 November 2021; Available online: <https://ecsoc-25.sciforum.net/>.

**Abstract:** A water-soluble pentamethine cyanine dye was synthesized for further conjugation with peptides. The dye preparation was accomplished by solution-phase chemistry and the synthesized compounds were characterized by the usual techniques (nuclear magnetic resonance, UV-vis absorption and fluorescence spectroscopies and mass spectrometry). The photophysical properties of cyanine **4** were studied in aqueous media, namely in phosphate-buffered saline (PBS) at pH 7.4. Sulfo-Cy5 carboxylic acid **4** exhibited a narrow absorption band and a high molar extinction coefficient ( $\log \epsilon = 5.11$ ) at 646 nm. The wavelength of maximum fluorescence was found in the near-infrared (NIR) region at 661 nm. Therefore, due to its excellent spectral properties, good water solubility and emission in the NIR region, dye **4** can be employed in the synthesis of fluorescence resonance energy transfer (FRET) probes for studying enzymatic hydrolysis in vitro.

**Keywords:** Cy5; cyanine dye; FRET probes; peptide labeling



**Citation:** Martins, C.D.F.; Raposo, M.M.M.; Costa, S.P.G. Synthesis and Characterization of a Water-Soluble Pentamethine Indocyanine Dye for Peptide Labeling. *Chem. Proc.* **2022**, *8*, 91. <https://doi.org/10.3390/ecsoc-25-11768>

Academic Editor: Julio A. Seijas

Published: 18 July 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Cyanine dyes are widely used as fluorescent labels in many biomedical applications such as bioimaging and diseases diagnosis [1], nucleic acid detection [2] and biomolecular labeling [3]. This is due to their excellent spectral properties, including narrow absorption bands, large molar absorptivities, good stability and high sensitivity [4–6].

The basic structure of cyanine dyes includes two heterocyclic rings containing nitrogen centers, which are linked by a conjugated chain of methine groups [6]. Depending on the length of the cyanine methine chain, the fluorescence range extends from the visible to the near-infrared (NIR) region [7,8]. For example, trimethine cyanine (Cy3) dyes with 3 methine groups emit visible light, while pentamethine cyanine (Cy5) dyes with 5 methine groups emit in the NIR region (>650 nm) [1,9,10].

Fluorescence detection in the NIR has many advantages, such as low tissue autofluorescence, small light scattering, background interference and deep tissue penetration, among others [11–13]. Therefore, due to the NIR fluorescence characteristics, pentamethine dyes have been employed as donor/acceptor groups in fluorescence resonance energy transfer (FRET) based enzyme probes to study biological processes in vitro and in vivo [14,15].

Among cyanine dyes, water-soluble sulfoindocyanine dye developed by Waggoner's group is one of the most used type of dyes for biological applications [16]. Sulfoindocyanine dyes have two or more  $\text{SO}_3^-$  groups on the ring systems, which enhance water solubility, prevent aggregation, and reduce non-specific binding to biomolecules and cellular constituents [4,17].

Bearing these facts in mind and considering our interest in the synthesis of FRET-labelled peptide substrates for in vitro validation of a nanoconstructed drug-delivery system, we report herein the synthesis of a water-soluble pentamethine cyanine dye.

The dye preparation was performed by solution-phase chemistry and its photophysical characterization was investigated in phosphate-buffered saline (PBS) at pH 7.4.

## 2. Experimental Section


### 2.1. Instruments and Materials

Nuclear Magnetic Resonance (NMR) spectra were obtained on a Bruker Avance III 400 at an operating frequency of 400 MHz for  $^1\text{H}$  and 100.6 MHz for  $^{13}\text{C}$  using the solvent peak as an internal reference. The solvents are indicated in parentheses before the chemical shift values ( $\delta$  relative to tetramethylsilane (TMS) and given in ppm). Assignments were supported by two-dimensional heteronuclear correlation techniques. Mass spectrometry analyses were performed at the “C.A.C.T.I. Unidad de Espectrometría de Masas” at the University of Vigo, Spain. Thin-layer chromatography (TLC) analyses were carried out on 0.25 mm-thick silica plates coated with fluorescent indicator F<sub>254</sub> (Merck KGaA, Darmstadt, Germany) and spots were visualized in a CN15 viewing cabinet under UV lamp at 365 nm (Vilber Lourmat, Marne-la-Vallée, France). Normal-phase column chromatography was performed on silica gel 0.035–0.070 mm, 60 Å (Acros Organics, Geel, Belgium). UV/Vis absorption spectra were obtained using a Shimadzu UV/2501PC spectrophotometer (Shimadzu Europa GmbH, Duisburg, Germany) and fluorescence spectra were collected using a FluoroMax-4 spectrofluorometer (HORIBA Europe GmbH, Darmstadt, Germany) in standard quartz cuvettes. All reagents were purchased from Acros Organics and Sigma-Aldrich (St. Louis, MO, USA), and used as received.

### 2.2. Synthesis

#### 2.2.1. Preparation of Sulfonated Indolium Salts

Potassium 2,3,3-trimethyl-3*H*-indole-5-sulfonate (**1**): 4-Hydrazinobenzenesulfonic acid hemihydrate (25 g, 127 mmol) and 3-methyl-2-butanone (40 mL, 371 mmol) were dissolved in acetic acid (75 mL) and the mixture was heated to reflux for 3 h. The solution was cooled to 4 °C, and the precipitate was collected by filtration. The residue was dissolved in methanol and then precipitated by addition of a saturated solution of potassium hydroxide in 2-propanol. The product was collected by filtration, washed with 2-propanol and diethyl ether, and dried in vacuo. Compound **1** (Figure 1) was obtained as purple solid (8.41 g, 24%). The product was pure enough to be used for the next reaction.




**Figure 1.** Structure of indolenine 1.

$^1\text{H}$  NMR (400 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  = 1.34 (6H, s, 2x $\text{C}3\text{-CH}_3$ ), 7.57 (1H, d,  $J$  = 8.0 Hz, H-7), 7.81 (1H, dd,  $J$  = 2.0 and 8.0 Hz, H-6), 7.86 (1H, d,  $J$  = 1.6 Hz, H-4) ppm. The  $\text{CH}_3$  group at position 2 was not observed in  $\text{D}_2\text{O}$ , in agreement with the literature [16].

$^{13}\text{C}$  NMR (100.6 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  = 21.64 (2x $\text{C}3\text{-CH}_3$ ), 54.17 (C3), 118.75 (C7), 119.30 (C4), 125.50 (C6), 139.67 (C5), 146.28 (C3a), 153.84 (C7a), 194.98 (C2) ppm.

1,2,3,3-Tetramethyl-3*H*-indolium-5-sulfonate iodide (**2**): The potassium salt of 2,3,3-trimethyl-3*H*-indole-5-sulfonate **1** (4 g, 14.4 mmol) was suspended in iodomethane (15 mL) and the reaction mixture was stirred at 40 °C for 6 h. The precipitate was collected by filtration, washed with acetone and dried in vacuo. Compound **2** (Figure 2) was obtained as purple solid (3.02 g, 55%). The product was used in the next reaction without additional purification.




**Figure 2.** Structure of indolium salt 2.

$^1\text{H}$  NMR (400 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  = 1.58 (6H, s,  $2\times\text{C}3\text{-CH}_3$ ), 4.05 (3H, s,  $\text{NCH}_3$ ), 7.56 (1H, d,  $J$  = 8.0 Hz, H-7), 7.81 (1H, dd,  $J$  = 1.6 and 8.0 Hz, H-6), 7.86 (1H, d,  $J$  = 1.6 Hz, H-4) ppm. The C2- $\text{CH}_3$  group was not observed in  $\text{D}_2\text{O}$ , in agreement with the literature [16].

$^{13}\text{C}$  NMR (100.6 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  = 21.73 ( $2\times\text{C}3\text{-CH}_3$ ), 34.63 ( $\text{NCH}_3$ ), 54.27 (C3), 118.73 (C7), 119.46 (C4), 125.64 (C6), 140.02 (C5), 146.21 (C3a), 153.27 (C7a), 198.85 (C2) ppm.

MS  $m/z$  (ESI, %): 254 ( $[\text{M}+\text{H}]^+$ , 100). HRMS:  $m/z$  (ESI) calc. for  $\text{C}_{12}\text{H}_{16}\text{NO}_3\text{S}$  254.0845; found 254.0848.

1-(5-Carboxypentyl)-2,3,3-trimethyl-3H-indolium-5-sulfonate bromide (**3**): The potassium salt of 2,3,3-trimethyl-3H-indole-5-sulfonate **1** (4 g, 14.4 mmol) and 6-bromohexanoic acid (3.25 g, 16.7 mmol) were mixed in chlorobenzene (35 mL) and heated at 110 °C for 12 h. The mixture was cooled, chlorobenzene was decanted, and the residue was triturated with 2-propanol, collected by filtration, washed with 2-propanol and dried in vacuo. Compound **3** (Figure 3) was obtained as purple solid (3.86 g, 62%). The product was used in the next reaction without additional purification.



**Figure 3.** Structure of indolium salt 3.

$^1\text{H}$  NMR (400 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  = 1.47 (2H, qui,  $\text{CH}_2\text{-c}$ ), 1.62 (6H, s,  $2\times\text{C}3\text{-CH}_3$ ), 1.68 (2H, qui,  $\text{CH}_2\text{-d}$ ), 2.00 (2H, qui,  $\text{CH}_2\text{-b}$ ), 2.39 (2H, t,  $J$  = 7.6 Hz,  $\text{CH}_2\text{-e}$ ), 4.54 (2H, t,  $J$  = 7.6 Hz,  $\text{CH}_2\text{-a}$ ), 7.92 (1H, d,  $J$  = 8.4 Hz, H-7), 8.04 (1H, dd,  $J$  = 1.6 and 8.4 Hz, H-6), 8.14 (1H, d,  $J$  = 1.6 Hz, H-4) ppm. The C2- $\text{CH}_3$  group was not observed in  $\text{D}_2\text{O}$ , in agreement with the literature [16].


$^{13}\text{C}$  NMR (100.6 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  = 21.54 ( $2\times\text{C}3\text{-CH}_3$ ), 23.74 ( $\text{CH}_2\text{-d}$ ), 25.22 ( $\text{CH}_2\text{-c}$ ), 26.76 ( $\text{CH}_2\text{-b}$ ); 33.37 ( $\text{CH}_2\text{-e}$ ); 48.16 ( $\text{CH}_2\text{-a}$ ), 54.88 (C3), 115.91 (C7), 120.88 (C4), 126.88 (C6), 142.57 (C3a), 142.74 (C7a), 144.18 (C5), 178.52 (C=O), 199.01 (C2) ppm.

MS  $m/z$  (ESI, %): 354 ( $[\text{M}+\text{H}]^+$ , 100). HRMS:  $m/z$  (ESI) calc. for  $\text{C}_{17}\text{H}_{24}\text{NO}_5\text{S}$  354.1370; found 354.1367.

#### 2.2.2. Synthesis of Sulfo-Cy5 Carboxylic Acid 4

1-(5-Carboxypentyl)-3,3-dimethyl-2-[5-(1',3',3'-trimethyl-5'-sulfonatoindol-1-ium-2'-yl)penta-2,4-dienylidene]indole-5-sulfonate chloride, sulfo-Cy5 carboxylic acid (**4**): A solution of indolium salt **3** (200 mg, 0.56 mmol) and malonaldehyde dianilide hydrochloride (176 mg, 0.68 mmol) in acetic anhydride (1.5 mL) was stirred at 120 °C for 30 min. The reaction mixture was cooled to room temperature and a solution of the indolium salt **2** (199 mg, 0.79 mmol) in pyridine (1.5 mL) was added. The reaction mixture was stirred at room temperature in the dark for 24 h. The product was precipitated by addition of ethyl acetate and collected by filtration. The crude residue was purified by a silica gel chromatography column, using as eluent mixtures of ethyl acetate and methanol with

increasing order of polarity (ethyl acetate: methanol, 7:3 > 6.5:3.5 > 6:4). The sulfo-Cy5 carboxylic acid **4** (Figure 4) was obtained as dark blue solid (72.8 mg, 19%).



**Figure 4.** Structure of dye **4**.

$^1\text{H}$  NMR (400 MHz,  $\text{MeOH-}d_4$ ):  $\delta$  = 1.49–1.55 (2H, m,  $\text{CH}_2$ -c), 1.67–1.77 (14H, m,  $\text{CH}_2$ -d +  $4\times\text{CH}_3$ ), 1.81–1.87 (2H, q,  $\text{CH}_2$ -b), 2.30 (2H, t,  $J$  = 7.2 Hz,  $\text{CH}_2$ -e), 3.67 (1H, d,  $J$  = 3.6 Hz,  $\text{NCH}_3$ ), 4.15 (2H, t,  $J$  = 6.8 Hz,  $\text{CH}_2$ -a); 6.36 (2H, dd,  $J$  = 5.6 and 13.6 Hz,  $\text{H-}\alpha$  +  $\text{H-}\alpha'$ ), 6.71 (1H, t,  $J$  = 12.4 Hz,  $\text{H-}\gamma$ ), 7.34–7.37 (2H, m,  $\text{H-}7$  +  $\text{H-}7'$ ), 7.89–7.91 (4H, m,  $\text{H-}4$  +  $\text{H-}6$  +  $\text{H-}4'$  +  $\text{H-}6'$ ), 8.32 (2H, t,  $J$  = 13.2 Hz,  $\text{H-}\beta$  +  $\text{H-}\beta'$ ) ppm.

$^{13}\text{C}$  NMR (100.6 MHz,  $\text{MeOH-}d_4$ ):  $\delta$  = 26.11 ( $\text{CH}_2$ -c), 27.42 ( $\text{CH}_2$ -d), 27.71 ( $2\times\text{CH}_3$ ), 27.84 ( $2\times\text{CH}_3$ ), 28.12 ( $\text{CH}_2$ -b), 31.77 ( $\text{NCH}_3$ ), 35.83 ( $\text{CH}_2$ -e), 45.06 ( $\text{CH}_2$ -a), 50.50 (C-3 or C-3'), 50.55 (C-3 or C-3'), 105.22 (C- $\alpha$  or C- $\alpha'$ ), 105.34 (C- $\alpha$  or C- $\alpha'$ ), 111.43 (C-7 or C-7'), 111.64 (C-7 or C-7'), 121.22 (C-6 or C-6'), 121.33 (C-6 or C-6'), 127.75 (C-4 or C-4' or C- $\gamma$ ), 128.01 (C-4 or C-4' or C- $\gamma$ ), 128.06 (C-4 or C-4' or C- $\gamma$ ), 142.48 (C-3a or C-3a'), 142.59 (C-3a or C-3a'), 143.31 (C-5 or C-5'), 143.35 (C-5 or C-5'), 144.93 (C-7a), 145.60 (C-7a'), 156.21 (C- $\beta$  + C- $\beta'$ ), 175.26 (C-2), 176.00 (C-2'), 178.88 (C=O) ppm.

### 2.3. Photophysical Characterization


The photophysical characterization of sulfo-Cy5 carboxylic acid **4** was carried out by UV-vis absorption and fluorescence spectroscopy of a  $1 \times 10^{-6}$  M solution in phosphate-buffered saline (PBS) at pH 7.4. The fluorescence spectrum of dye **4** was obtained by excitation at the wavelength of maximum absorption.

The relative fluorescence quantum yield was calculated relative to a standard solution of Nile Blue in methanol ( $\Phi_F = 0.27$ ) [18].

## 3. Results and Discussion

### 3.1. Synthesis of Sulfo-Cy5 Carboxylic Acid **4**

The preparation of sulfo-Cy5 carboxylic acid **4** was accomplished by solution-phase chemistry described by Waggoner and co-workers [16], according to Scheme 1. The key intermediate, indoleninium-5-sulfonate, was prepared from 4-hydrazinobenzenesulfonic acid hemihydrate and 3-methyl-2-butanone by conventional Fisher indole synthesis [16]. The indolenine thus obtained was converted into a potassium salt before quaternization in 24% yield. This compound was alkylated using iodomethane or 6-bromohexanoic acid to give compound **2** in 55% and compound **3** in 62%, respectively, which were used in the next reaction steps without additional purification.



**Scheme 1.** Synthesis of sulfo-Cy5 carboxylic acid **4**: (a) 3-methyl-2-butanone, acetic acid, 3 h, reflux; (b) iodomethane, 6 h, 40 °C; (c) 6-bromohexanoic acid, chlorobenzene, 12 h, 110 °C; (d) (i) acetic anhydride, 30 min, 120 °C; (ii) **2**, pyridine, 24, r.t.

Stepwise condensation reaction of the indolenines **2** and **3** with a polyene-chain precursor (malonaldehyde dianilide hydrochloride) gave good conversion to the desired sulfoindocyanine **4**. This was precipitated from the reaction mixture by addition of ethyl acetate and the precipitate was further purified by column chromatography on silica gel. Isolation of this compound proved to be very difficult, as side products and cleavage product had nearly the same polarity. Nevertheless, sulfo-Cy5 carboxylic acid **4** was obtained in the form of a dark blue solid in 19% yield.

The synthesized compounds were characterized by  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopy and mass spectroscopy, and the obtained data were in agreement with the expected structure.

### 3.2. Photophysical Characterization


A solution of sulfo-Cy5 carboxylic acid **4** ( $1 \times 10^{-6}$  M) at pH 7.4 in phosphate-buffered saline (PBS) was analyzed and the wavelengths of maximum absorption and fluorescence,  $\lambda_{\text{abs}}$  and  $\lambda_{\text{flu}}$ , molar absorptivity at the absorption maximum,  $\epsilon$ , relative fluorescence quantum yield,  $\Phi_F$ , and Stokes' shifts are compiled in Table 1.

**Table 1.** UV-vis absorption and fluorescence data of sulfo-Cy5 carboxylic acid **4** ( $1 \times 10^{-6}$  M) at pH 7.4 in PBS.

Compound	UV-Vis Absorption		Fluorescence		
	$\lambda_{\text{abs}}$ (nm)	$\log \epsilon$	$\lambda_{\text{flu}}$ (nm)	$\Phi_F$	Stokes' Shift
Dye <b>4</b>	646	5.11	661	0.27	15

The synthesized dye **4** exhibited a narrow absorption band (Figure 5), a high molar absorptivity ( $\log \epsilon = 5.11$ ) at 646 nm, and a sharp emission band with a wavelength

of maximum fluorescence in the NIR region ( $\lambda_{\text{flu}} = 661 \text{ nm}$ ). The relative fluorescence quantum yield, determined by using Nile Blue in methanol as standard ( $\Phi_F = 0.27$ ), was found to be low ( $\Phi_F = 0.27$ ), which is in agreement with the literature [16]. Although the extension of the polymethine bridge presents red shifted emission, lower quantum yields were found for NIR cyanine dyes in water. However, the presence of sulfo groups could significantly enhance water solubility and prevent aggregation and fluorescence self-quenching in aqueous media [4,17]. Taking this into account, sulfo-Cy5 carboxylic acid **4** can be activated to form succinimidyl ester for conjugation with a specific peptide.



**Figure 5.** Normalized absorption and emission spectra of sulfo-Cy5 carboxylic acid **4** ( $1 \times 10^{-6} \text{ M}$ ) at pH 7.4 in PBS.

#### 4. Conclusions

In summary, a water-soluble pentamethine dye bearing two sulfo groups was synthesized by solution solid phase synthesis in 19% yield. However, future work will be developed to optimize the isolation method to increase the overall synthesis yield.

Photophysical characterization of sulfo-Cy5 carboxylic acid **4** showed an intense absorption band with a high molar absorptivity at 646 nm and the wavelength of maximum fluorescence was found in the NIR region at 661 nm. Therefore, due to its excellent spectral properties, good solubility in water and emission in the NIR region, dye **4** can be used for our intentions for it as a donor group in FRET probes to study enzymatic hydrolysis in vitro.

**Author Contributions:** Conceptualization, S.P.G.C.; methodology, C.D.F.M. and S.P.G.C.; validation, S.P.G.C. and M.M.M.R.; formal analysis, C.D.F.M., M.M.M.R. and S.P.G.C.; investigation, C.D.F.M.; resources, S.P.G.C. and M.M.M.R.; writing—original draft preparation, C.D.F.M.; writing—review and editing, C.D.F.M., M.M.M.R. and S.P.G.C.; supervision, S.P.G.C. and M.M.M.R.; project administration, S.P.G.C.; funding acquisition, M.M.M.R. and S.P.G.C. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the Foundation for Science and Technology (FCT) for financial support to CQ/UM (UID/QUI/00686/2020) and project PTDC/QUI-COL/28052/2017 and a PhD grant to C. D. F. Martins (SFRH/BD/05277/2020). The NMR spectrometer Bruker Avance III 400 is part of the National NMR Network and was purchased within the framework of the National Program for Scientific Re-equipment, contract REDE/1517/RMN/2005 with funds from POCI 2010 (FEDER) and FCT.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. An, H.W.; Wang, M.D. The self-assembly of cyanine dyes for biomedical application in vivo. In *In Vivo Self-Assembly Nanotechnology for Biomedical Applications*; Wang, H., Li, L.L., Eds.; Springer: Singapore, 2018; pp. 31–55.
2. Zhang, X.-H.; Wang, L.-Y.; Nan, Z.-X.; Tan, S.-H.; Zhang, Z.-X. Microwave-assisted solvent-free synthesis and spectral properties of some dimethine cyanine dyes as fluorescent dyes for DNA detection. *Dyes Pigm.* **2008**, *79*, 205–209. [[CrossRef](#)]
3. Patonay, G.; Salon, J.; Sowell, J.; Streckowski, L. Noncovalent labeling of biomolecules with red and near-infrared dyes. *Molecules* **2004**, *9*, 40–49. [[CrossRef](#)]
4. Wang, L.; Fan, J.; Qiao, X.; Peng, X.; Dai, B.; Wang, B.; Sun, S.; Zhang, L.; Zhang, Y. Novel asymmetric Cy5 dyes: Synthesis, photostabilities and high sensitivity in protein fluorescence labeling. *J. Photochem. Photobiol. A Chem.* **2010**, *210*, 168–172. [[CrossRef](#)]
5. Owens, E.A.; Bruschi, N.; Tawney, J.G.; Henary, M. A microwave-assisted and environmentally benign approach to the synthesis of near-infrared fluorescent pentamethine cyanine dyes. *Dyes Pigm.* **2015**, *113*, 27–37. [[CrossRef](#)]
6. Pham, W.; Medarova, Z.; Moore, A. Synthesis and application of a water-soluble near-infrared dye for cancer detection using optical imaging. *Bioconjugate Chem.* **2005**, *16*, 735–740. [[CrossRef](#)] [[PubMed](#)]
7. Wolf, N.; Kersting, L.; Herok, C.; Mihm, C.; Seibel, J. High-yielding water-soluble asymmetric cyanine dyes for labeling applications. *J. Org. Chem.* **2020**, *85*, 9751–9760. [[CrossRef](#)] [[PubMed](#)]
8. Pisoni, D.S.; Todeschini, L.; Borges, A.C.A.; Petzhold, C.L.; Rodembusch, F.S.; Campo, L.F. Symmetrical and asymmetrical cyanine dyes. Synthesis, spectral properties, and BSA association study. *J. Org. Chem.* **2014**, *79*, 5511–5520. [[CrossRef](#)] [[PubMed](#)]
9. Shindy, H.A. Fundamentals in the chemistry of cyanine dyes: A review. *Dyes Pigm.* **2017**, *145*, 505–513. [[CrossRef](#)]
10. Bříza, T.; Rimpelová, S.; Králová, J.; Záruba, K.; Kejík, Z.; Ruml, T.; Martásek, P.; Král, V. Pentamethinium fluorescent probes: The impact of molecular structure on photophysical properties and subcellular localization. *Dyes Pigm.* **2014**, *107*, 51–59. [[CrossRef](#)]
11. Yi, X.; Wang, F.; Qin, W.; Yang, X.; Yuan, J. Near-infrared fluorescent probes in cancer imaging and therapy: An emerging field. *Int. J. Nanomed.* **2014**, *9*, 1347–1365. [[CrossRef](#)] [[PubMed](#)]
12. Luo, S.; Zhang, E.; Su, Y.; Cheng, T.; Shi, C. A review of NIR dyes in cancer targeting and imaging. *Biomaterials* **2011**, *32*, 7127–7138. [[CrossRef](#)] [[PubMed](#)]
13. Feng, L.; Chen, W.; Ma, X.; Liu, S.H.; Yin, J. Near-infrared heptamethine cyanines (Cy7): From structure, property to application. *Org. Biomol. Chem.* **2020**, *18*, 9385–9397. [[CrossRef](#)] [[PubMed](#)]
14. Lee, S.; Park, K.; Lee, S.-Y.; Ryu, J.H.; Park, J.W.; Ahn, H.J.; Kwon, I.C.; Youn, I.-C.; Kim, K.; Choi, K. Dark quenched matrix metalloproteinase fluorogenic probe for imaging osteoarthritis development in vivo. *Bioconjugate Chem.* **2008**, *19*, 1743–1747. [[CrossRef](#)] [[PubMed](#)]
15. Jia, X.; Chen, Q.; Yang, Y.; Tang, Y.; Wang, R.; Xu, Y.; Zhu, W.; Qian, X. FRET-based mito-specific fluorescent probe for ratiometric detection and imaging of endogenous peroxynitrite: Dyad of Cy3 and Cy5. *J. Am. Chem. Soc.* **2016**, *138*, 10778–10781. [[CrossRef](#)] [[PubMed](#)]
16. Mujumdar, R.B.; Ernst, L.A.; Mujumdar, S.R.; Lewis, C.J.; Waggoner, A.S. Cyanine dye labeling reagents: Sulfoindocyanine succinimidyl esters. *Bioconjugate Chem.* **1993**, *4*, 105–111. [[CrossRef](#)] [[PubMed](#)]
17. Lopalco, M.; Koini, E.N.; Cho, J.K.; Bradley, M. Catch and release microwave mediated synthesis of cyanine dyes. *Org. Biomol. Chem.* **2009**, *7*, 856–859. [[CrossRef](#)] [[PubMed](#)]
18. Jose, J.; Ueno, Y.; Burgess, K. Water-soluble Nile Blue derivatives: Syntheses and photophysical properties. *Chem. Eur. J.* **2008**, *15*, 418–423. [[CrossRef](#)]