

Home Composting: A Review of Scientific Advances [†]

Raquel Barrena and Antoni Sánchez * 

Department of Chemical, Biological and Environmental Engineering, Universitat Autònoma de Barcelona (UAB), 08193 Bellaterra, Barcelona, Spain; raquel.barrena@uab.cat

* Correspondence: antoni.sanchez@uab.cat

[†] Presented at the 1st International Electronic Conference on Processes: Processes System Innovation, 17–31 May 2022; Available online: <https://sciforum.net/event/ECP2022>.

Abstract: Composting has been demonstrated to be a sustainable technology for treating organic wastes. The process is based on the microbial decomposition of organic matter under aerobic conditions to obtain compost: An organic amendment that can be safely used in agriculture and other applications. Among the composted wastes, the organic fraction of municipal waste is commonly used. In this sense, the interest in composting at home or community scale is exponentially growing in recent years, as it permits the self-management of waste and obtaining a product that can be used by the own producer. However, some questions about the quality of the obtained compost or the environmental impact of home composting are in an early stage of this development. In this review, the main points related to home composting are analyzed in detail according to the current scientific knowledge. Among them: (i) The performance of the process, especially in the temperature reached and the fact that if home compost is sanitized, (ii) the quality of home composting, especially in terms of maturity and stability, (iii) the main environmental impacts of home composting, that is, gaseous emissions and (iv) the main trends related to community composting, a step forward from home composting, to make this alternative attractive for municipal organic waste management. The main advantages and possible drawbacks of home composting are also highlighted.

Keywords: organic fraction of municipal waste; home composting; process performance; compost quality; environmental impact

check for
updates

Citation: Barrena, R.; Sánchez, A. Home Composting: A Review of Scientific Advances. *Eng. Proc.* **2022**, *19*, 35. <https://doi.org/10.3390/ECP2022-12625>

Academic Editor: Dariusz Dziki

Published: 20 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Among municipal waste, food waste (FW) comprises the main fraction (45%) of total municipal solid waste in Europe [1]. This percentage can increase up to 55% in developing countries [2]. Some years ago, the final destination of FW was either the disposal in controlled/uncontrolled landfills or incineration with/without energy recovery.

FW treatment is usually performed by biological processes, such as composting and anaerobic digestion, although new strategies are being developed to obtain added value bioproducts from organic waste [3]. Composting is based on the biological degradation of organic matter under aerobic conditions, and compost is the product of composting. The process is considered a sustainable and environmentally friendly alternative for treating FW and is used worldwide at a commercial scale in industrial composting plants [4].

In contrast to these big facilities, an increasing number of initiatives using home or community composting have emerged in different parts of the world. Initially, these experiences were explained as hobbies and low scientific information could be obtained from them [5]. Today, we have scientifically based information about home and community composting in different aspects, such as the performance of the process, the quality of home compost, or the environmental impact and Life Cycle Assessment.

The objective of this communication is to present a general perspective of home composting, especially in those studies with scientific data, to provide updated information to the readers about the state-of-the-art of this technology.

2. Home Composting

2.1. Process Performance

Home composting performance is different from that of industrial full-scale composting. On the one hand, it is difficult to maintain the thermophilic conditions that allow the compost to be sanitized [6] or the presence of considerable stratification of temperature found in home composters [7]. The results obtained in the composters indicate that small-scale composting was viable since thermophilic sanitization temperatures (55 °C) were maintained for three consecutive days. However, stability indicators showed a different organic matter biodegradation process along the compost bin's height, with the bottom layer requiring a longer period to be stable than the upper layers. The authors conclude that these phenomena can have importance when designing commercial home composting reactors. On the other hand, other works studied other important aspects of home composting performance, such as mass balances [8], which are very important for further studies related to Life Cycle Inventories and Life Cycle Assessment (LCA) [9].

Regarding the performance of the process, other aspects are considered in recent literature. Among them, the need for a bulking agent has special relevance to provide porosity to the mixture to have a strict aerobic process and to avoid unpleasant odors and unwanted gaseous emissions in the form of greenhouse gases, especially methane and nitrous oxide (Figure 1).



Figure 1. Detail of a home composter with a proper porosity adjustment.

In this sense, Castiglioni Guidoni et al. (2018) investigated how different ratios of bulking agent and food waste can affect the progress of the composting process [10]. Results showed that the ratio of bulking agents has an important effect on the biodegradation of organic matter, nitrogen dynamics, and the toxicity of the product. It is evident that this ratio must be carefully studied in home composting to reach an equilibrium between biodegradation and environmental impact [11]. Regarding including FW in a home composting process, there is some discussion on including meat and, in general, food waste of animal origin. Storino et al. (2016) studied the comparison of home composting with and without the presence of meat in the initial mixture [11]. The authors concluded that meat has several positive effects on the process activity and an acceleration of the biodegradation of organic matter without altering the main physicochemical characteristics (pH, salinity, or phytotoxicity) and a low pathogen level with proper handling of the home composters. However, no information on gaseous emissions expected from meat waste, such as ammonia, is provided. Other authors showed low gaseous emissions when composting leftovers of raw fruits and vegetables without animal wastes [12]. In the case of inoculation, home composting is a perfect system to use the obtained compost as a source of active microorganisms to maintain a semi-continuous process, although some authors point out that the

presence of the so-called “Effective Microorganisms” is positive in only particular aspects, such as odor control and humification [13].

Finally, some authors comment that collection is a critical step to having a good home composting process. Thus, Puyuelo et al. (2013) performed a comparison of several methods to collect food waste before composting, showing that the use of perforated passively aerated bins jointly with compostable bags was superior to other conventional systems [14]. For instance, this system did not imply more gaseous emissions, and it was suitable for preparing the organic waste for further composting. Besides, in terms of weight loss, temperature, respiration index, and organic matter reduction, the best results were also achieved with the aerated system (Figure 2).



Figure 2. Collection of food waste using compostable bags and aerated bins.

2.2. Home Compost Quality

In reference to the quality of compost obtained from home systems, there is a consensus in the sense that this quality is similar, if not better, than that of industrial facilities. The absence of impurities in the initial mixture is the main reason for this high quality of home composting. Thus, Vázquez and Soto (2017) present a study including 880 experiences of home composting in rural areas, using household biowaste, including meat and fish leftovers [15]. Ninety home compost samples were analyzed, showing excellent properties: A low C/N ratio (10–15), no physical contaminant materials (less than 0.3% in dry matter), low heavy metal content, and high nutrient content (2.1% N, 0.6% P, 2.5% K, 0.7% Mg, and 3.7% Ca, respectively). Other studies show similar results of home compost quality in terms of physicochemical characterization (pH, moisture, carbon, nitrogen, and C/N ratio) [16,17]. Regarding the presence of pathogenic microorganisms, they were not detected, even when thermophilic temperatures were not fully reached [11].

Apart from these studies, the literature is scarce on the comparison of home and industrial composting. An especially interesting study is that of Barrena et al. (2014), where a large number of home and industrial composts were compared in terms of stability, using respiration techniques [18]. The main conclusion is that home composting, when properly managed, can reach high levels of stability, similar to those of full-scale composting. If only physicochemical properties are compared, industrial and home composts are not significantly different.

2.3. Environmental Impact

Regarding environmental impacts of home composting, they have been quite extensively studied in the literature. On the one hand, the main advantage of home composting is the absence of waste transport and lower energy requirements, whereas a possible disadvantage is the lack of gaseous emissions control as in full-scale facilities, which is practically unavoidable [19]. On the other hand, there is a need to compare home composting with industrial composting from the environmental point of view to have an experimentally-

based comparison. The preliminary studies on this topic were presented by Colón et al. (2010) and Andersen et al. (2012) [9,20]. Both papers performed a very complete study on the environmental impacts, which were compiled in an LCA. The conclusions of both works are similar: Home composting performed better than other waste management technologies in most of the impact categories. Both studies also agree with the fact that gaseous emissions are the main contributors to negative environmental impacts in different environmental categories, especially in the global warming potential.

Accordingly, further studies on the environmental impact focused on gaseous emissions during the home composting process [21]. For instance, Ermolaev et al. (2014) studied the greenhouse gases (GHG) emissions of several home composters treating FW and compared them with literature data from full-scale composting [22]. In this case, home composting emitted less methane than large-scale composts but similar amounts of nitrous oxide. Other works have focused on this comparison in rigorous studies with the same initial mixture, season, and location, with more environmental categories studied. This is the case of Colón et al. (2012) [23]. In this study, four different full-scale facilities treating a source-selected organic fraction of Municipal Solid Waste (OFMSW) were environmentally evaluated with an LCA, including composting technologies (in-vessel, turned piles, and home composting) and anaerobic digestion plus composting. In this case, home composting was better in terms of environmental impact than the other composting technologies. In a previous work, Martínez-Blanco et al. (2010) directly compared a complete LCA home and industrial composting [24]. In summary, the results were as expected: Ammonia and GHG emissions (methane and nitrous oxide) released from home composting were considerably higher than those of industrial composting. However, this latter option involved 2 and 53 times more needs for transport, energy, water, and infrastructure.

3. Community Composting

Although used extensively in many countries, especially in central Europe [25], and having an extraordinary development in recent years, community composting has received low attention in the world of research. Thus, there are starting experiences in universities, hospitals, municipal markets, or just groups of households. However, the information found in the scientific literature is very scarce.

Most of the studies on community composting are related to economic feasibility. From these studies, it is evident that many entities can use community composting. In this case, universities can play an important role as the first stakeholders to impulse this strategy that can be easily extrapolated to other organic waste producers, such as hotels, hospitals, or schools. Community composting can have an important influence on two items: On the one hand, it can treat a significant amount of organic waste and, on the other hand, it can be a stimulus to promote home composting among citizens [26]. In Figure 3, a recently developed program of community composting in Pontevedra (Spain) is presented.



Figure 3. Community composting in Pontevedra (Spain).

4. Conclusions

It can be concluded that home and community composting has a great potential to be a massively implemented strategy in organic waste management when developing national and regional programs. One important reason for this expansion is the fact that home composting has passed from being a hobby to a scientifically-based technology. From the environmental and economic points of view, home and community composting appears to be superior to industrial composting in most of the environmental categories, which again makes this strategy attractive to be included in waste management programs.

Author Contributions: Conceptualization, A.S.; methodology, R.B.; formal analysis, A.S.; investigation, R.B. and A.S.; writing—original draft preparation, A.S.; writing—review and editing, A.S.; funding acquisition, A.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by *Agència de Gestió d'Ajuts Universitaris i de Recerca of Generalitat de Catalunya*, grant number Pandèmies Project Ref. 2020PANDE00021.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. IPCC—Intergovernmental Panel on Climate Change. Waste Generation, Compositions and Management Data Guidelines for National Greenhouse Gas Inventories. Intergovernmental Panel on Climate Change, WMO/UNEP. 2006. Available online: <https://www.ipcc-nggip.iges.or.jp/public/2006gl/vol5.html> (accessed on 31 March 2022).
2. Troschinetz, A.M.; Mihelcic, J.R. Sustainable recycling of municipal solid waste in developing countries. *Waste Manag.* **2009**, *29*, 915–923. [[CrossRef](#)] [[PubMed](#)]
3. Cerda, C.; Artola, A.; Barrera, R.; Font, X.; Gea, T.; Sánchez, A. Innovative Production of Bioproducts From Organic Waste Through Solid-State Fermentation. *Front. Sustain. Food Syst.* **2019**, *3*, 63. [[CrossRef](#)]
4. Cerda, A.; Artola, A.; Font, X.; Barrera, R.; Gea, T.; Sánchez, A. Composting of food wastes: Status and challenges. *Bioresour. Technol.* **2019**, *248*, 57–67. [[CrossRef](#)]
5. Mayoral, J.; Sánchez, A. Backyard composting in Catalonia (Spain). Results from an extensive pilot program. *Biocycle* **2005**, *46*, 75–78.

6. Tatàno, F.; Pagliaro, G.; Giovanni, P.; Floriani, E.; Mangani, F. Biowaste home composting: Experimental process monitoring and quality control. *Waste Manag.* **2015**, *38*, 72–85. [[CrossRef](#)] [[PubMed](#)]
7. Arrigoni, J.P.; Paladino, G.; Garibaldi, L.A.; Laos, F. Inside the small-scale composting of kitchen and garden wastes: Thermal performance and stratification effect in vertical compost bins. *Waste Manag.* **2018**, *76*, 284–293. [[CrossRef](#)]
8. Andersen, J.K.; Boldrin, A.; Christensen, T.H.; Scheutz, C. Mass balances and life cycle inventory of home composting of organic waste. *Waste Manag.* **2011**, *31*, 1934–1942. [[CrossRef](#)]
9. Colón, J.; Martínez-Blanco, J.; Gabarrell, X.; Artola, A.; Sánchez, A.; Rieradevall, J.; Font, X. Environmental assessment of home composting. *Resour. Conserv. Recycl.* **2010**, *54*, 893–904. [[CrossRef](#)]
10. Castiglioni Guidonia, L.L.; Vasques Marquesa, R.; Bilhalva Moncks, R.; Torma Botelho, F.; da Paza Luciara, F.; Corrêa, B.; Corrêa, E. Home composting using different ratios of bulking agent to food waste. *J. Environ. Manag.* **2018**, *207*, 141–150. [[CrossRef](#)]
11. Storino, F.; Arizmendiarieta, J.S.; Irigoyen, I.; Muro, J.; Aparicio-Tejo, P.M. Meat waste as feedstock for home composting: Effects on the process and quality of compost. *Waste Manag.* **2016**, *56*, 53–62. [[CrossRef](#)]
12. Ballardó, C.; Vargas-García, M.C.; Sánchez, A.; Barrena, R.; Artola, A. Adding value to home compost: Biopesticide properties through *Bacillus thuringiensis* inoculation. *Waste Manag.* **2020**, *106*, 32–43. [[CrossRef](#)] [[PubMed](#)]
13. Fan, Y.V.; Lee, C.T.; Klemeš, J.J.; Chua, L.S.; Sarmidi, M.R.; Leow, C.W. Evaluation of Effective Microorganisms on home scale organic waste composting. *J. Environ. Manag.* **2018**, *216*, 41–48. [[CrossRef](#)] [[PubMed](#)]
14. Puyuelo, B.; Colón, J.; Martín, P.; Sánchez, A. Comparison of compostable bags and aerated bins with conventional storage systems to collect the organic fraction of municipal solid waste from homes. A Catalonia case study. *Waste Manag.* **2013**, *33*, 1381–1389. [[CrossRef](#)] [[PubMed](#)]
15. Vázquez, M.A.; Soto, M. The efficiency of home composting programmes and compost quality. *Waste Manag.* **2017**, *64*, 39–50. [[CrossRef](#)] [[PubMed](#)]
16. Papadopoulos, A.E.; Stylianou, M.A.; Michalopoulos, C.P.; Moustakas, K.G.; Hapeshis, K.M.; Vogiatzidaki, E.E.I.; Loizidou, M.D. Performance of a new household composter during in-home testing. *Waste Manag.* **2019**, *29*, 204–213. [[CrossRef](#)]
17. Kuchel, M.; Raclavská, H.; Růžicková, J.; Švédová, B.; Sassmanová, V.; Drozdová, J.; Raclavský, K.; Juchelková, D. Properties of composts from household food waste produced in automatic composters. *J. Environ. Manag.* **2019**, *236*, 657–666. [[CrossRef](#)]
18. Barrena, R.; Font, X.; Gabarrell, X.; Sánchez, A. Home composting versus industrial composting: Influence of composting system on compost quality with focus on compost stability. *Waste Manag.* **2014**, *34*, 1109–1116. [[CrossRef](#)]
19. Colón, J.; Martínez-Blanco, J.; Gabarrell, X.; Rieradevall, J.; Font, X.; Artola, A.; Sánchez, A. Performance of an industrial biofilter from a composting plant in the removal of ammonia and VOCs after material replacement. *J. Chem. Technol. Biotechnol.* **2009**, *84*, 1111–1117. [[CrossRef](#)]
20. Andersen, J.K.; Boldrin, A.; Christensen, T.H.; Scheutz, C. Home composting as an alternative treatment option for organic household waste in Denmark: An environmental assessment using life cycle assessment-modelling. *Waste Manag.* **2012**, *32*, 31–40. [[CrossRef](#)]
21. Quirós, R.; Villalba, G.; Muñoz, P.; Colón, J.; Font, X.; Gabarrell, X. Environmental assessment of two home composts with high and low gaseous emissions of the composting process. *Resour. Conserv. Recycl.* **2014**, *90*, 9–20. [[CrossRef](#)]
22. Ermolaev, E.; Sundberg, C.; Pell, M.; Jönsson, H. Greenhouse gas emissions from home composting in practice. *Bioresour. Technol.* **2014**, *151*, 174–182. [[CrossRef](#)] [[PubMed](#)]
23. Colón, J.; Cadena, E.; Pognani, M.; Barrena, R.; Sánchez, A.; Font, X.; Artola, A. Determination of the energy and environmental burdens associated to the biological treatment of source-separated Municipal Solid Wastes. *Energy Environ. Sci.* **2012**, *5*, 5731–5741. [[CrossRef](#)]
24. Martínez-Blanco, J.; Colón, J.; Gabarrell, X.; Font, X.; Sánchez, A.; Artola, A.; Rieradevall, J. The use of life cycle assessment for the comparison of biowaste composting at home and full scale. *Waste Manag.* **2010**, *30*, 983–994. [[CrossRef](#)] [[PubMed](#)]
25. Regions for Recycling, 2014. Good Practice Styria: Biowaste Collection. Available online: https://www.acrplus.org/images/project/R4R/Good_Practices/GP_Styria_biowaste-collection.pdf (accessed on 31 March 2022).
26. Catalan Waste Agency. Government of Catalonia. Guide for the Implementation of Community Composting. Available online: <https://residus.gencat.cat/ca/detalls/Publicacions/Guia-compost-comunitari> (accessed on 31 March 2022).