



## Abstract Photophysical Properties of Some Naphthalimide Derivatives <sup>+</sup>

Radu Ionut Tigoianu<sup>1,\*</sup>, Anton Airinei<sup>1</sup>, Florentina Georgescu<sup>2</sup>, Alina Nicolescu<sup>1</sup> and Calin Deleanu<sup>1,3</sup>

- <sup>1</sup> "Petru Poni" Institute of Macromolecular Chemistry, Gr. Ghica Voda 41A, 700487 Iasi, Romania
- <sup>2</sup> Enpro Soctech Com srl, Str. Elefterie 51, 050524 Bucharest, Romania
  <sup>3</sup> C. D. Nanitsony Contra of Organic Chamistry, Spl. Indexendente: 20
- C. D. Nenitescu Centre of Organic Chemistry, Spl. Independentei 202B, 060023 Bucharest, Romania
- \* Correspondence: tigoianu.radu@icmpp.ro
- + Presented at the 9th International Electronic Conference on Sensors and Applications, 1–15 November 2022; Available online: https://ecsa-9.sciforum.net/.

**Abstract:** Naphthalimide derivatives possess many interesting properties such as strong emission, high quantum efficiency, good photostability, thermal stability, etc. The electronic absorption and fluorescence spectra of naphthalimides are sensitive to the polarity of surrounding environment, and these derivatives can be excellent candidates for fluorescent sensors for water detection in solution because the emission is strongly depended on the solvent polarity and it is quenched even at low water levels. In order to find out more information about the excited state dynamics of the naphthalimide derivatives, time-resolved fluorescence experiments were conducted in solvents of different polarities, and lifetimes from 0.5 ns to 9 ns were obtained. The transient absorption map in dioxane, dimethylformamide and methanol in the presence or absence of water revealed ground state bleaching bands (GSB) in the range of 230–290 nm, whereas an absorption band in excited state (ESA) occurred at shorter wavelengths from 210 to 295 nm. At longer wavelength, negative bands appeared, which can be assigned to the stimulated emissions (SE). In addition, the quantum yields with absolute values from 0.01 to 0.87 were found depending on the solvent nature.

Keywords: naphthalimides; transient absorption; water sensing; lifetimes



Citation: Tigoianu, R.I.; Airinei, A.; Georgescu, F.; Nicolescu, A.; Deleanu, C. Photophysical Properties of Some Naphthalimide Derivatives. *Eng. Proc.* 2022, *27*, 4. https://doi.org/ 10.3390/ecsa-9-13356

Academic Editor: Francisco Falcone

Published: 1 November 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). **Supplementary Materials:** The following material is available at https://www.mdpi.com/article/ 10.3390/ecsa-9-13356/s1. Conference poster: Photophysical Properties of some Naphthalimide Derivatives.

Funding: This research received no external funding.

Informed Consent Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.