

Effect of Thermal Processing on Carotenoids in Fortified Bread [†]

Edward Muntean

Department of Food Sciences, Faculty of Food Sciences and Technology, University of Agricultural Sciences and Veterinary Medicine, 400372 Cluj-Napoca, Romania; edimuntean@yahoo.com

[†] Presented at the 2nd International Electronic Conference on Processes: Process Engineering—Current State and Future Trends (ECP 2023), 17–31 May 2023; Available online: <https://ecp2023.sciforum.net/>.

Abstract: Bread is a staple food that provides essential nutrients to millions of people worldwide, hence its fortification with provitamin A carotenoids can help combat vitamin A deficiencies effectively. This study investigates fortified bread made with *Cucurbita maxima* Duch. fruits using a straight-dough procedure, focusing on changes caused by thermal processing to carotenoids. High performance liquid chromatography (HPLC) with photodiode-array detection was used for the sensitive and selective analysis of carotenoids. HPLC showed that lutein, cucurbitaxanthin A, and β -carotene are the major carotenoids in fortified bread, highlighting lutein and cucurbitaxanthin A as the most stable carotenoids during thermal processing. Fortified bread is a functional food that can not only combat vitamin A deficiency and prevent age-related macular degeneration, but also has antioxidant properties with related health effects, offering new market opportunities for producers.

Keywords: bread; fortification; carotenoids; chromatography; HPLC; analysis; functional foods

1. Introduction

Vitamin A deficiency is a major public health issue in many parts of the world, particularly in low-income countries; because it can lead to serious health problems and can have a profound impact on child development, it has been addressed by numerous studies in recent decades, some of which involve food fortification approaches [1,2]. Bread is an ideal food vector for fortification because it is a staple food in many parts of the world, which provides an essential source of carbohydrates, proteins and minerals to millions of people [3,4]. In such a context, bread fortification is necessary because bread alone is not sufficient to meet the nutritional requirements of a balanced diet as it lacks important micronutrients; bread fortification with synthetic or natural provitamin A carotenoids can provide a solution for vitamin A deficiency and *Cucurbita maxima* fruits can be used for this purpose [5–7].

Carotenoids are biologically active compounds widely recognized for their advantageous impact on human health; numerous studies have shown that consuming carotenoid-rich foods is linked to a decreased risk of lung, breast, colon, and prostate cancers, as well as UV-induced skin damage, coronary heart disease, cataracts, and macular degeneration, and certain carotenoids with β -ring end groups can act as provitamin A within the human body [8,9]. Unfortunately, these compounds are also sensitive to environmental conditions such as heat, light, extreme pH values, and contact with oxidants [10,11]. Therefore, their use in technological applications must be properly handled in order to obtain the desired effects.

The fortification of bread with carotenoids of natural origin is an important factor in ensuring consumer acceptance and compliance, hence this study targets fortified bread obtained through a straight-dough procedure, in which a puree made from *Cucurbita maxima* Duch. fruits was added. Because carotenoids are heat-sensitive biomolecules, high performance liquid chromatography (HPLC) with photodiode-array detection was used as the method of analysis, this being the most appropriate analytical technique to date for their analysis [12,13].



Citation: Muntean, E. Effect of Thermal Processing on Carotenoids in Fortified Bread. *Eng. Proc.* **2023**, *37*, 70. <https://doi.org/10.3390/ECP2023-14694>

Academic Editor: Chi-Fai Chau

Published: 17 May 2023



Copyright: © 2023 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

The major objective of this study was to investigate the effect of thermal processing on carotenoids in fortified bread, using HPLC as analytical tool, to provide relevant information for food technology, nutritional studies and to support the creation of a functional food that offers health benefits beyond combating vitamin A deficiency.

2. Materials and Methods

2.1. Materials

All solvents used for chromatography were of HPLC grade purity (Merck). They were filtered through 0.45 µm Whatman filters and then degassed in an ultrasonic bath, under vacuum, before use. Solvents used for extraction were of p.a. quality; water was bidistilled, then degassed. Potassium hydroxide was from Merck. The reference carotenoids neoxanthin, violaxanthin, antheraxanthin, lutein zeaxanthin and all-E-β,β-carotene were from CaroteNature GmbH, Lupsingen, Switzerland. All the analytical operations were carried out in reduced light, avoiding sample heating at more than 40 °C; prior to injection into the HPLC system, all samples were filtered (0.45 µm, Whatman).

2.2. Biological Material & Bread Fortification

Mature *Cucurbita maxima* Duch. fruits were harvested from the experimental field of the University of Agricultural Sciences and Veterinary Medicine, Cluj Napoca, Romania. The fruits were sliced, the seeds and the placental tissues were removed, and then they were baked at 200 °C for 60 min. After cooling, the crust was removed from each slice, and the slices were homogenized in a blender.

Laboratory-scale bread preparation was accomplished using a straight-dough procedure by mixing 300 g of commercial wheat flour (Baneasa—000), 5 g of dry yeast, 4 g of sodium chloride, 150 g of water and 100 g + of puree. The dough was mixed to optimum development, then was left for fermentation at 25 °C for two hours, and was then divided into portions of 100 g which were manually molded. The final dough fermentation was accomplished at room temperature for one hour, then baking was performed at 215 °C for 25 min; the baked breads were allowed to cool on wooden racks at room temperature.

2.3. Sampling and Sample Preparation

Breads were sliced and samples were collected from each bread core; core bread samples of ~1 g were thoroughly homogenized. Then, average samples of ~20 g were weighed and subjected to extraction with 100 mL acetone in a blender, filtered under vacuum using a sintered glass funnel, and then the solid material was re-extracted twice with 50 mL acetone.

The combined filtrates were subjected to liquid–liquid extraction in a separation funnel in which 100 mL of diethyl ether was added and washed five times with 100 mL of distilled water. The resulting extract was evaporated to dryness under reduced pressure in a Buchi rotary evaporator at 40°C, re-dissolved in 25 mL diethyl ether, and then saponified with a 25 mL solution of 30% KOH in methanol on a magnetic stirrer (350 rpm, room temperature, 16 h). Carotenoids were next extracted with 50 mL diethyl ether and washed repeatedly with distilled water until free of alkali; the extract was evaporated to dryness under reduced pressure, and then the residue was dissolved in 10 mL of ethyl acetate, from which an aliquot was filtrated through a 0.47 µm membrane filter and subjected to HPLC analysis.

2.4. HPLC Analysis

HPLC analysis was performed according to a published procedure [14]; carotenoid identification was completed based on retention times, HPLC co-chromatography with standards, and visible spectral characteristics compared to reference carotenoids and literature data [15]. Quantifications were achieved using the external standard method [14].

2.5. Data Processing

Chromatographic data analysis was accomplished using Waters 990 software (Waters Corporation, Milford, MA, USA), then chromatographic data were further processed using Excel (Microsoft Corporation, Redmond, WA, USA). The provitamin A concentrations were expressed in retinol equivalents (R.E.) according to FAO/WHO requirements [16].

3. Results and Discussion

Reversed-phase HPLC enabled the quantification of three carotenes (all-E- β , β -carotene, 9Z- β -carotene and 15Z- β -carotene) and five xanthophylls (neoxanthin, violaxanthin, cucurbitaxanthin A, lutein and zeaxanthin), highlighting that the major carotenoids in fortified bread were lutein, cucurbitaxanthin A and all-E- β , β -carotene (Figures 1–3).




Figure 1. Concentration profile of carotenoids in baked *Cucurbita* fruit puree used for fortification.




Figure 2. Concentration profile of carotenoids in the initial dough (a) and in the fortified bread (b).




Figure 3. Comparative percentage of carotenoid degradation as a result of thermal processing.

Because the baked *Cucurbita maxima* fruit puree contains considerable amounts of carotenoids (Figure 1), its addition to the dough increased the carotenoid content (Figure 2), the major ones being lutein, β -carotene and cucurbitaxanthin A—the last one being a carotenoid specific to the *Cucurbita* genus, with no provitamin A activity.

Thermal processing caused massive carotenoid degradation, which affected both xanthophylls and carotenes. The effects of thermal processing on carotenoids are revealed in Figures 2 and 3, which prove that zeaxanthin and violaxanthin were the most affected carotenoids, having been degraded to the greatest extent, whereas cucurbitaxanthin A and lutein proved to be the most stable carotenoids under these circumstances. A special case is that of 9Z- β , β -carotene, a product of all-E- β , β -carotene isomerization, because its overall concentration increased after thermal processing; this is the reason why it was not included in Figure 3.

Because thermal processing had an important effect on the most important provitamin A carotenoid in the fortified bread (β -carotene), the retinol equivalent was affected accordingly (Figure 4).




Figure 4. Comparative values for retinol equivalents in the investigated food matrices.

4. Conclusions

- HPLC analysis demonstrated that the major carotenoids in fortified bread were lutein, cucurbitaxanthin A and β -carotene. It also highlighted the stability of the targeted compounds and the change in provitamin A activity as a result of thermal processing during baking; the most stable carotenoids proved to be lutein and cucurbitaxanthin A.
- The behavior of the involved carotenoids can provide relevant information for nutritional studies.
- Bread fortified with baked *Cucurbita* is not only a possible solution for combating vitamin A deficiency, but it is also a true functional food.
- Bread products fortified with naturally occurring carotenoids may be of interest to consumers because of the health benefit of the antioxidant effect, as compared with non-fortified products. The use of natural sources of carotenoids, such as *Cucurbita maxima* fruits, can also increase consumer acceptance and compliance.
- Fortified bread can also open new market opportunities for producers; being a solution to a serious public health issue, it can be used as a selling point for health-conscious consumers.

Supplementary Materials: The presentation materials can be downloaded at: <https://www.mdpi.com/article/10.3390/ECP2023-14694/s1>.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Akhtar, S.; Ahmed, A.; Randhawa, M.A.; Atukorala, S.; Arlappa, N.; Ismail, T.; Ali, Z. Prevalence of vitamin A deficiency in South Asia: Causes, outcomes, and possible remedies. *J. Health Popul. Nutr. Nutr.* **2013**, *31*, 413–423. [CrossRef] [PubMed]
2. Gernand, A.D.; Schulze, K.J.; Stewart, C.P.; West, K.P., Jr.; Christian, P. Micronutrient deficiencies in pregnancy worldwide: Health effects and prevention. *Nat. Rev. Endocrinol.* **2016**, *12*, 274–289. [CrossRef] [PubMed]
3. Aghalari, Z.; Dahms, H.U.; Sillanpää, M. Evaluation of nutrients in bread: A systematic review. *J. Health Popul. Nutr.* **2022**, *41*, 50. [CrossRef] [PubMed]
4. Ibrahim, U.K.; Salleh, R.M.; Maqsood-ul-Haque, S.N.S. Bread towards functional food: An overview. *Int. J. Food Eng.* **2015**, *1*, 39–43. [CrossRef]
5. Muntean, E.; Rotar, I. HPLC assessment of provitamin A carotenoids from Cucurbita maxima Duch. ex. lam.(Marița cultivar) fruits. *Res. J. Agric. Sci.* **2010**, *42*, 517–520.
6. Muntean, E.; Muntean, N.; Duda, M.M. Cucurbita maxima Duch. as a medicinal plant. *Hop Med. Plants* **2013**, *1*, 75–80.
7. Muntean, E. Carotenoids in Cucurbita fruits. In Proceedings of the 2nd International Electronic Conference on Nutrients, On-line, 15–30 March 2022; Available online: <https://sciforum.net/manuscripts/12381/slides.pdf> (accessed on 15 April 2023).
8. Perera, C.O.; Yen, G.M. Functional properties of carotenoids in human health. *Int. J. Food Prop.* **2007**, *10*, 201–230. [CrossRef]
9. Rao, A.V.; Rao, L.G. Carotenoids and human health. *Pharmacol. Res.* **2007**, *55*, 207–216. [CrossRef] [PubMed]
10. Boon, C.S.; McClements, D.J.; Weiss, J.; Decker, E.A. Factors influencing the chemical stability of carotenoids in foods. *Crit. Rev. Food Sci. Nutr.* **2010**, *50*, 515–532. [CrossRef] [PubMed]
11. Christaki, E.; Bonos, E.; Giannenas, I.; Florou-Paneri, P. Functional properties of carotenoids originating from algae. *J. Sci. Food Agr.* **2013**, *93*, 5–11. [CrossRef] [PubMed]
12. Felzl, L.; Pacakova, V.; Stulik, K.; Volka, K. Reliability of carotenoid analyses: A review. *Curr. Anal. Chem.* **2005**, *1*, 93–102. [CrossRef]
13. Zhao, X.; Liang, K.; Zhu, H. Carotenoids in Cereals and Related Foodstuffs: A Review of Extraction and Analysis Methods. *Food Rev. Int.* **2022**, *14*, 1–16. [CrossRef]
14. Muntean, E.; Muntean, N.; Modoran, C. HPLC analysis of carotenoids in bread. *Sci. Res. Agroaliment. Process. Technol.* **2004**, *10*, 186–190.
15. Britton, G. UV/Visible spectroscopy. In *Carotenoids—Vol. 1: Spectroscopy*; Britton, G., Liaaen-Jensen, S., Pfander, H., Eds.; Birkhauser: Basel, Switzerland, 1996; pp. 13–62.
16. World Health Organization. *Vitamin and Mineral Requirements in Human Nutrition*, 2nd ed.; World Health Organization: Geneva, Switzerland, 2004; pp. 24–26. Available online: <http://apps.who.int/iris/bitstream/handle/10665/42716/9241546123.pdf?sequence=1> (accessed on 15 April 2023).

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.