

Abstract

Structural Transformation in Liquid Tellurium from Stillinger–Weber Potential [†]

Hafiz Ghulam Abbas ¹, Kamal Prasad Sapkota ², Md. Akherul Islam ³, Md. Abu Hanif ³, Jeasmin Akter ² and Jae Ryang Hahn ^{1,4,*}

¹ Department of Nanoscience and Technology, Research Institute of Physics and Chemistry, Jeonbuk National University, Jeonju 54896, Korea; hgabbas71@gmail.com

² Department of Chemistry, Research Institute of Physics and Chemistry, Jeonbuk National University, Jeonju 54896, Korea; mychemistry2037@gmail.com (K.P.S.); tina44445@gmail.com (J.A.)

³ Department of Bioactive Material Sciences, Jeonbuk National University, Jeonju 54896, Korea; akherulraju@gmail.com (M.A.I.); hanif4572@gmail.com (M.A.H.)

⁴ Textile Engineering, Chemistry and Science, North Carolina State University, 2401 Research Dr., Raleigh, NC 27695-8301, USA

* Correspondence: jrhahn@bnu.ac.kr

[†] Presented at the 2nd International Online-Conference on Nanomaterials, 15–30 November 2020; Available online: <https://iocn2020.sciforum.net/>.

Abstract: Structural evolutions in liquid tellurium (Te) are observed by employing molecular dynamics simulations at various temperatures ranging from 1500 K to 300 K. Local structural variations are noticed in radial correlation functions, structure factors, bond angle distribution functions, Honeycutt–Anderson index, Voronoi tessellation, and coordination numbers. Upon quenching, we found that icosahedral short-range motifs dominated in a stable and supercooled liquid state. The first peak of the radial distribution function at 970 K and 722 K is in excellent agreement with the findings of neutron diffraction. The transformation to a supercooled liquid state with distorted icosahedral patterns is observed at 600 K and to a body-centered cubic cluster after 600 K. Finally, we also show that near the melting point diffusion coefficient of liquid tellurium is fairly consistent with the tight-binding and experimental models. We assume that our findings not only replicate all the remarkable characteristics but also predict useful transition mechanisms through the use of the well-known Stillinger–Weber potential.

Citation: Abbas, H.G.; Sapkota, K.P.; Islam, M.A.; Hanif, M.A.; Akter, J.; Hahn, J.R. Structural Transformation in Liquid Tellurium from Stillinger–Weber Potential. *Mater. Proc.* **2021**, *4*, 83. <https://doi.org/10.3390/IOCN2020-07971>

Published: 12 November 2020

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).