

Article

Field Evaluation of Preventive Efficacy of Monthly Multimodal Prophylactic Treatment (Milbactor[®] and Vectra[®] 3D) against *Dirofilaria* spp. in Dogs

Younes Laidoudi ^{1,2,*} , Hacène Medkour ^{1,2} , Djamel Tahir ^{1,2} , Handi Dahmana ^{1,2}, Jean-Lou Marié ³, Marie Varloud ⁴ , Oleg Mediannikov ^{1,2}  and Bernard Davoust ^{1,2,3,*} 

¹ IRD, AP-HM, MEPHI, Aix Marseille University, IHU Méditerranée Infection, 19-21, Bd Jean Moulin, CEDEX 05, 13385 Marseille, France; hacenevet1990@yahoo.fr (H.M.); djamel.tahir@yahoo.fr (D.T.); handy92@hotmail.fr (H.D.); olegusss1@gmail.com (O.M.)

² IHU Méditerranée Infection, Assistance Publique-Hôpitaux de Marseille, 19-21, Bd Jean Moulin, CEDEX 05, 13385 Marseille, France

³ Animal Epidemiology Expert Group, French Military Health Service, 37100 Tours, France; jean-lou.marie@intradef.gouv.fr

⁴ Ceva Santé Animale, 10, Av de la Ballastière, 33500 Libourne, France; marie.varloud@ceva.com

* Correspondence: younes.laidoudi@yahoo.com (Y.L.); bernard.davoust@gmail.com (B.D.); Tel.: +33-0413732401 (B.D.)



Citation: Laidoudi, Y.; Medkour, H.; Tahir, D.; Dahmana, H.; Marié, J.-L.; Varloud, M.; Mediannikov, O.; Davoust, B. Field Evaluation of Preventive Efficacy of Monthly Multimodal Prophylactic Treatment (Milbactor[®] and Vectra[®] 3D) against *Dirofilaria* spp. in Dogs. *Parasitologia* **2021**, *1*, 130–136. <https://doi.org/10.3390/parasitologia1030014>

Academic Editors: Maurizio Francesco Brivio and Geoff Hide

Received: 30 June 2021

Accepted: 22 July 2021

Published: 30 July 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: *Dirofilaria immitis* and *Dirofilaria repens* are mosquito-borne pathogens responsible for dirofilariasis in humans and animals. Their transmission and spread depend on the activity of vectors and the frequency of hosts in a given area. Here, we investigated the efficacy of a monthly multimodal prophylactic (MMP) strategy against canine dirofilariasis on Corsica Island (France). The study was conducted as evidence of an efficacy trial in which eighty dogs were divided into two groups: (i) one test group consisted of 25 dogs under the MMP [per-os administration of 1.5 tablets of milbemycin-oxime-praziquantel (Milbactor[®]) and a topical line-on application of a 3.6 mL solution of dinotefuran-permethrin-pyriproxyfen (Vectra[®] 3D)] and (ii) a control group under different real-life prophylactic treatments (RLP) based on the use of ectoparasiticides (different formulations: deltamethrin, fluralaner, fipronil) and/or macrocyclic lactone-based products (milbemycin-oxime/praziquantel, milbemycin-oxime, moxidectin) during the period from June to October 2017. All animals were followed up for one year, with blood collected at day 0, with follow-up at 6 and 12 months. Samples were tested for *Dirofilaria* spp. by species-specific qPCR. At the end of the study, no new case of *Dirofilaria* spp. infection was detected in the test group. However, the cumulative incidence of *Dirofilaria* spp. infection was 16.4% (n = 9; p = 0.027) in the control group. The data illustrate that, in contrast to RLP treatment, which failed to protect at least 16.4% of dogs, the MMP based on the simultaneous administration of milbemycin oxime-praziquantel and dinotefuran-permethrin-pyriproxyfen efficiently protects dogs in a high-risk area from *Dirofilaria* spp. infection.

Keywords: dog; *Dirofilaria* spp.; multimodal prophylaxis; prevention

1. Introduction

Dirofilaria immitis and *Dirofilaria repens* are nematodes of tissues and cavities responsible for cardiopulmonary and subcutaneous and/or subcutaneous/ocular dirofilariasis in canids and humans, respectively [1]. These heteroxenous parasites produce blood-dwelling microfilariae, which are assumed by blood-feeding vectors (i.e., Culicidae of the genera *Culex*, *Aedes*, *Ochlerotatus*, *Anopheles*, *Coquillettidia*, *Armigeres*, *Mansonia* and *Psorophora*) [2] and go through two stages to reach the infective stage (L3), which can then be transmitted to a new receptive host. These parasites appear to be associated with an episytem complex involving multiple factors such as temperature, movement and abundance of

vectors and hosts, as well as the pathogen itself [3], which complicates disease control [4]. Epidemiological studies indicate the continuous spread of *Dirofilaria* spp. in human and canine populations, especially in Europe, where endemicity and sympatric occurrence have already been reported.

Two different preventive strategies are used by practitioner veterinarians against *Dirofilaria* spp. infection, mainly based on seasonal application of repellent/insecticidal products against vector bites and thus parasite transmission, or on macrocyclic lactones (ML) targeting *Dirofilaria* larvae. Recent recommendations from the European Society for Dirofilariosis and Angiostrongylosis (ESDA), the European Scientific Counsel Companion Animal Parasites (ESCCAP) and the American Heartworm Society (AHS) indicate the need to strengthen preventive measures by combining both methods as a multimodal prophylaxis in endemic areas [5,6]. The multimodal prophylactic (MMP) strategy was initially developed as an urgent solution to prevent the spread of ML-resistant strains of *D. immitis*, which has been proven to be a promising chemoprophylaxis under experimental conditions [7]. However, no data are available on the efficacy of the MMP strategy against *Dirofilaria* spp. infection under natural conditions.

Corsica is a French island known to be endemic for *Dirofilaria* spp. for four decades. Moreover, the island is surrounded by highly endemic areas and represents a touristic destination for more than 750,000 visitors and their pets each year, which places it in first rank in the dissemination of *Dirofilaria* spp. and other vector-borne pathogens to non-endemic areas [8–11]. Such an epidemiological context encompasses all risk factors for *Dirofilaria* spp. transmission and spread and thus represents best challenge for the evaluation of the newly developed preventive strategies under natural conditions. In the current study, the efficacy of the MMP strategy based on simultaneous treatment with Vectra® 3D (dinotefuran-permethrin-pyriproxyfen) (Ceva Santé Animale, Libourne, France) and Milbactor® (milbemycin oxime-praziquantel) (Ceva Santé Animale, Libourne, France) against mosquito vectors and *Dirofilaria* spp. larvae, respectively, was investigated.

2. Results

At day 180 of follow-up, no new cases of *Dirofilaria* spp. infection were detected in the studied dogs from both the test (n = 25) and control (n = 55) group. However, unlike the test group, 9 (16.4%) new infections consisting of 8 (14.6%) and 1 (1.8%) *D. immitis* and *D. repens*, respectively, were recorded in the control group at Day 365, showing a significant difference between these two groups (Table 1) and a protection level of 100% against *Dirofilaria* spp. transmission in test group.

Table 1. Incidence density rates (IDR) of *Dirofilaria* spp. infection in the studied dogs. P-values compare the IDR rates between the test and control group.

IDRs (%)			Pairwise Comparison (Gray's Test)	
Pathogens/Time	Test Group (n = 25)	Control Group (n = 55)	Statistic	p-Value
<i>Dirofilaria</i> spp.				
Day 180	0	0	4.55	0.033
Day 365	0	16.4		
<i>D. immitis</i>				
Day 180	0	0	4.0	0.046
Day 365	0	14.6		
<i>D. repens</i>				
Day 180	0	0	0.46	0.500
Day 365	0	1.8		

3. Discussion

The present study demonstrates the field efficacy of consistent administration of larvicidal (macrocyclic lactones) plus vector-preventive products (repellents) as MMP against *Dirofilaria* spp. infection in endemic areas and the first European field study confirming the year-round efficacy of MMP on a monthly schedule against these vector-borne parasites. *Dirofilaria immitis* is responsible for canine cardiopulmonary dirofilariasis (known as heartworm disease) with varied symptoms including exercise intolerance, fatigue to right heart and lung failure [12]. Treatment of canine heartworm can be life-threatening, expensive and complex [6,12]. In contrast, *D. repens* infections are often asymptomatic [3]. However, fatal outcomes were observed in *D. repens* infestation due to the massive release of the filarial endosymbiotic bacterium *Wolbachia* after microfilaricidal treatment [13]. Therefore, to avoid these complications, preventive chemoprophylaxis remains the best approach to control these parasitoses [5,6].

The efficacy evaluation of the MMP protocol was conducted here as evidence of an efficacy trial using a positive control group. It is expected that the protective success in the test group compared to the control group is due to several elements, such as compliance, which is known to be low in France (6%) [14], leading to prevention failures [15]. Another key factor could be the treatment period, where year-round chemoprophylaxis was challenged compared to seasonal chemoprophylaxis in the control group. In southern Europe, such as Corsica, the presence of year-round active vectors of *Dirofilaria* spp. such as *Stegomyia albopicta* has been reported [16,17]. Consequently, year-round control of such vectors is recommended [4]. Another key factor for the effective protection in the test group could be the combined use of two different products against *Dirofilaria* vectors and parasite larvae. This approach has been registered to provide broad-spectrum activity against nematodes and ectoparasites in dogs [7,18,19]. In addition, the separate use of Vectra® 3D (dinotefuran-permethrin-pyriproxyfen) and Milbactor® (milbemycin oxime/praziquantel) have been proven to be effective against a wide range of vectors and *Dirofilaria* larvae [20–24], leading to a reinforcement of their synergetic effect against *Dirofilaria* transmission when administrated concomitantly.

In addition to the direct effect on mosquito-borne helminths, the intended multimodal prophylactic strategy based on year-round administration of Vectra® 3D and Milbactor® on a monthly schedule provides reliable control against *Dirofilaria* spp. infections in dogs in highly endemic areas, and it should be promoted more widely in and around these areas to prevent the spread of *Dirofilaria* spp. to dogs and humans.

4. Materials and Methods

4.1. Study Design

From March 2017 to April 2018, a cohort study was conducted in three neighboring areas on Corsica Island (i.e., Ventiseri 41°55'52.5" N; 9°23'58.3" E, Marine de Solaro 41°50.1114' N; 9°22.482' E and Aleria 42°6'15.293" N; 9°30'44.744" E) as evidence of efficacy trial [25], according to the guidelines on data requirements for veterinary medicinal products for the prevention of transmission of vector-borne diseases in dogs and cats [26].

Dogs were recruited after obtaining owner consent. Inclusion criteria were qPCR-negative for at least one *Dirofilaria* spp., with the absence of clinical signs consistent with heartworm, which could be a threat to a dog's life during the follow-up, an age of at least one year at the time of inclusion and the application of prophylactic treatment against dirofilariosis. Finally, dogs lost during follow-up were excluded from the study (Figure 1).




Figure 1. Flow diagram showing the semi-annual evolution of the cohort.

4.2. Animals and Treatment Protocol

A total of 80 dogs were included and allocated into two groups:

1. The test group (test treatment group, n = 25) consisted of military working dogs (MWD) housed individually in wire cages in a kennel in Ventiseri (Figure 2a). The dogs received monthly multimodal prophylactic treatment (MMP) (per os administration of 1.5 tablets of milbemycin oxime-praziquantel (Milbactor®, Ceva Santé Animale, Libourne, France) and topical line-on application of a 3.6 mL solution of dinotefuran-permethrin-pyriproxyfen (Vectra® 3D, Ceva Santé Animale)).
2. The control group (active control, n = 55) consisted of civilian dogs from Marine de Solaro and those from the Aleria kennel. All dogs lived outdoor in wire cages or free in gardens (Figure 2b). These dogs received various real-life prophylactic treatments (RLP) destined to prevent against vector biting or to block the development of *Dirofilaria* larvae.



Figure 2. Photo shoot showing housing conditions of dogs from (A) test and (B) control groups.

Citrated blood samples were collected at the time of enrollment (Day 0), after six (Day 180) and after 12 (Day 360) months, in accordance with the requirements of the Animal Ethics Procedures of French veterinarians and with the consent of dog owners. Genomic DNA was extracted individually from each whole blood using the Biorobot EZ1 System with the EZ1 DNA tissue kit (Qiagen, Courtaboeuf, France) according to the manufacturer's instructions, and then screened for the presence of *Dirofilaria* spp. at each time point using species-specific multiplex qPCR, as previously described [27].

4.3. Estimation of Incidence and Data Analysis

Animals were considered positive if the infection with at least one *Dirofilaria* species was detected. The incidence density rate (IDR) represented by the rate of newly infected dogs (i.e., new infections) was calculated. Gray's test [28] was applied to compare between the test and control group for *Dirofilaria* spp. incidence. Significance was defined at $\alpha \leq 0.05$. Statistical analysis were performed using Addinsoft XLSTAT 2018 (Data analysis and statistical solution for Microsoft Excel, Paris, France). The protection level of MMP was expressed as the percentage of success in blocking the transmission of *Dirofilaria* spp. infections and calculated at the end of the follow-up period using the following formula: $\text{Protection (\%)} = 100 \times ((\text{IDRc} - \text{IDRt}) / \text{IDRc})$, where IDRc and IDRt represent the incidence density rate of *Dirofilaria* spp. infection in the control and test group, respectively [26].

Author Contributions: Y.L., B.D., M.V., J.-L.M. and O.M. designed the study. Y.L., B.D., D.T., H.D. and J.-L.M. intervened in the field study. Y.L. and H.M. performed the laboratory analyses. Y.L., B.D. and O.M. designed, and Y.L. carried out, the data analysis. Y.L. wrote the original draft. Y.L., B.D., M.V. and O.M. reviewed and edited the final draft. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by the Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, the National Research Agency under the program "Investissements d'avenir", reference ANR-10-IAHU-03, the Région Provence-Alpes-Côte d'Azur and European funding FEDER PRIM1. This work was also funded by Ceva Santé Animale (supply of pharmaceutical products).

Institutional Review Board Statement: Ethical review and approval were waived according to French and European legislation because the study does not involve laboratory animals and protocols.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: We thank Sandrine Ferrandi, Marie-Hélène Filippi, Mickaël Boni, Laetitia Guidicelli and all the dog owners for their help.

Conflicts of Interest: All the authors declare no conflicts of interest related to this article. One author (Marie Varloud) is currently an employee of Ceva Santé Animale. However, there were no conflicts of interest that could have biased the work presented in this paper.

References

1. Simón, F.; Siles-Lucas, M.; Morchón, R.; González-Miguel, J.; Mellado, I.; Carretón, E.; Montoya-Alonso, J.A. Human and animal dirofilariasis: The emergence of a zoonotic mosaic. *Clin. Microbiol. Rev.* **2012**, *25*, 507–544. [CrossRef] [PubMed]
2. Aonuma, H.; Yoshimura, A.; Perera, N.; Shinzawa, N.; Bando, H.; Oshiro, S.; Nelson, B.; Fukumoto, S.; Kanuka, H. Loop-mediated isothermal amplification applied to filarial parasites detection in the mosquito vectors: *Dirofilaria immitis* as a study model. *Parasit. Vectors* **2009**, *2*, 15. [CrossRef]
3. Simón, F.; González-Miguel, J.; Diosdado, A.; Gómez, P.J.; Morchón, R.; Kartashev, V. The complexity of zoonotic filariasis episystem and its consequences: A multidisciplinary view. *Biomed. Res. Int.* **2017**, *2017*, 6436130. [CrossRef]
4. Schorderet-Weber, S.; Noack, S.; Selzer, P.M.; Kaminsky, R. Blocking transmission of vector-borne diseases. *Int. J. Parasitol. Drugs Drug Resist.* **2017**, *7*, 90–109. [CrossRef]
5. European Scientific Counsel Companion Animal Parasites (ESCCAP). Control of Vector-Borne 5 Diseases in Dogs and Cats. Guideline 05, 3rd ed 2019. Available online: http://www.t2kkcbgl_0775_ESCCAP_Guideline_GL5_v9_1p.pdf (accessed on 15 June 2021).
6. Nelson, C.T.; McCall, J.W.; Jones, S.; Moorhead, A. Current canine guidelines for the prevention, diagnosis, and management of heartworm (*Dirofilaria immitis*) infection in dogs. *Am. Hear. Soc.* **2018**, 1–35. Available online: <https://heartwormsociety.org/images/pdf/2014-AHS-Canine-Guidelines.pdf> (accessed on 15 June 2021).
7. McCall, J.W.; Varloud, M.; Hodgkins, E.; Mansour, A.; Dicosty, U.; McCall, S.; Carmichael, J.; Carson, B.; Carter, J. Shifting the paradigm in *Dirofilaria immitis* prevention: Blocking transmission from mosquitoes to dogs using repellents/insecticides and macrocyclic lactone prevention as part of a multimodal approach. *Parasit. Vectors* **2017**, *10* (Suppl. 2), 525. [CrossRef]
8. Moraes-Da-Silva, M.D.F.C.V.; Mendes-De-Almeida, F.; Abdalla, L.; Merlo, A.; Paiva, J.P.; Labarthe, N.V. Selamectin for the prevention of canine *Dirofilaria immitis* infection: Field efficacy in client-owned dogs in a high risk area. *Parasit. Vectors* **2016**, *9*, 1–5. [CrossRef]
9. Pantchev, N.; Etzold, M.; Dauschies, A.; Dyachenko, V. Diagnosis of imported canine filarial infections in Germany 2008–2010. *Parasitol. Res.* **2011**, *109* (Suppl. 1), S61–S76. [CrossRef]
10. Zahler, M.; Glaser, B.; Gothe, R. Imported parasites in dogs: *Dirofilaria repens* and *Dipetalonema reconditum*. *Tierarztl. Prax.* **1997**, *25*, 388–392. (In German)
11. Berger, S.M. *Infectious Diseases of France*, 2019th ed.; Gideon Informatics, Inc.: Los Angeles, CA, USA, 2019; ISBN 978-1-4988-2193-3.
12. McCall, J.W.; Genchi, C.; Kramer, L.H.; Guerrero, J.; Venco, L. Heartworm disease in animals and humans. *Adv. Parasitol.* **2008**, *66*, 193–285. [CrossRef]
13. Wyszczek, M.E.; Klockiewicz, M.; Sobczak-Filipiak, M. Case studies of severe microfilaremia in four dogs naturally infected with *Dirofilaria repens* as the primary disease or a disease complicating factor. *Front. Vet. Sci.* **2020**, *7*, 577466. [CrossRef]
14. Roussel, C.; Drake, J.; Ariza, J.M. French national survey of dog and cat owners on the deworming behaviour and lifestyle of pets associated with the risk of endoparasites. *Parasit. Vectors* **2019**, *12*, 480. [CrossRef] [PubMed]
15. McTier, T.L.; Kryda, K.; Wachowski, M.; Mahabir, S.; Ramsey, D.; Rugg, D.; Mazaleski, M.; Terrien, C.; Adams, E.; Wolff, T.; et al. ProHeart® 12, a moxidectin extended-release injectable formulation for prevention of heartworm (*Dirofilaria immitis*) disease in dogs in the USA for 12 months. *Parasit. Vectors* **2019**, *12*, 369. [CrossRef] [PubMed]
16. Tahir, D.; Bittar, F.; Barré-Cardi, H.; Sow, D.; Dahmani, M.; Mediannikov, O.; Raoult, D.; Davoust, B.; Parola, P. Molecular survey of *Dirofilaria immitis* and *Dirofilaria repens* by new real-time TaqMan® PCR assay in dogs and mosquitoes (Diptera: Culicidae) in Corsica (France). *Vet. Parasitol.* **2017**, *235*, 1–7. [CrossRef] [PubMed]
17. Otranto, D.; Dantas-Torres, F.; Brianti, E.; Traversa, D.; Petrić, D.; Genchi, C.; Capelli, G. Vector-borne helminths of dogs and humans in Europe. *Parasit. Vectors* **2013**, *6*, 1–14. [CrossRef] [PubMed]
18. Prichard, R.K.; Geary, T.G. Perspectives on the utility of moxidectin for the control of parasitic nematodes in the face of developing anthelmintic resistance. *Int. J. Parasitol. Drugs Drug Resist.* **2019**, *10*, 69–83. [CrossRef]
19. Labarthe, N.V.; Willi, L.M.V.; Paiva, J.P.; De Miranda, M.G.N.; Zoreck, K.; Mendes De Almeida, F. Chemoprophylaxis of *Dirofilaria immitis* (Leidy 1856) infection at a high challenge environment. *Parasit. Vectors* **2015**, *8*, 6–9. [CrossRef] [PubMed]
20. European Medicines Agency. Vectra3D Product Information EPAR 2015. 2013. Available online: http://www.ema.europa.eu/docs/en_GB/document_library (accessed on 15 June 2021).

21. Tahir, D.; Davoust, B.; Almeras, L.; Berenger, J.M.; Varloud, M.; Parola, P. Anti-feeding and insecticidal efficacy of a topical administration of dinotefuran–pyriproxyfen–permethrin spot-on (Vectra® 3D) on mice against *Stegomyia albopicta* (= *Aedes albopictus*). *Med. Vet. Entomol.* **2017**, *31*, 351–357. [[CrossRef](#)]
22. Tahir, D.; Davoust, B.; Varloud, M.; Berenger, J.M.; Raoult, D.; Almeras, L.; Parola, P. Assessment of the anti-feeding and insecticidal effects of the combination of dinotefuran, permethrin and pyriproxyfen (Vectra® 3D) against *Triatoma infestans* on rats. *Med. Vet. Entomol.* **2017**, *31*, 132–139. [[CrossRef](#)] [[PubMed](#)]
23. Laidoudi, Y.; Tahir, D.; Medkour, H.; Varloud, M.; Mediannikov, O.; Davoust, B. Effect of dinotefuran, permethrin, and pyriproxyfen (Vectra® 3D) on the foraging and blood-feeding behaviors of *aedes albopictus* using laboratory rodent model. *Insects* **2020**, *11*, 507. [[CrossRef](#)]
24. di Cesare, A.; Braun, G.; Di Giulio, E.; Paoletti, B.; Aquilino, V.; Bartolini, R.; La Torre, F.; Meloni, S.; Drake, J.; Pandolfi, F.; et al. Field clinical study evaluating the efficacy and safety of an oral formulation containing milbemycin oxime/praziquantel (Milbemax®, Novartis Animal Health) in the chemoprevention of the zoonotic canine infection by *Dirofilaria repens*. *Parasit. Vectors* **2014**, *7*, 347. [[CrossRef](#)]
25. U.S. Department of Health and Human Services; Food and Drug Administration. E10 Choice of Control Group and Related Issues in Clinical Trials.. Available online: <https://www.fda.gov/media/71349/download> (accessed on 15 June 2021).
26. European Medicines Agency. Guideline on Data Requirements for Veterinary Medicinal Products for the Prevention of Transmission of Vector-Borne Diseases in Dogs and Cats. 2018. Available online: <https://www.ema.europa.eu/en> (accessed on 15 June 2021).
27. Laidoudi, Y.; Davoust, B.; Varloud, M.; Niang, E.H.A.; Fenollar, F.; Mediannikov, O. Development of a multiplex qPCR-based approach for the diagnosis of *Dirofilaria immitis*, *D. repens* and *Acanthocheilonema reconditum*. *Parasit. Vectors* **2020**, *13*, 319. [[CrossRef](#)] [[PubMed](#)]
28. Dignam, J.J.; Kocherginsky, M.N. Choice and interpretation of statistical tests used when competing risks are present. *J. Clin. Oncol.* **2008**, *26*, 4027–4034. [[CrossRef](#)] [[PubMed](#)]