

Proceeding Paper

Phytochemical and Antioxidant Properties of *Athamanta turbith* (L.) Brot Collected from Serbia †

Sofija N. Kilibarda ^{1,*} , Sandra Z. Vuković ¹, Danijel D. Milinčić ¹ , Marina P. Mačukanović-Jocić ¹, Snežana Jarić ² and Aleksandar Ž. Kostić ¹ 

¹ Faculty of Agriculture, University of Belgrade, Nemanjina 6, 11080 Belgrade, Serbia; sandra.vukovic@agrif.bg.ac.rs (S.Z.V.); danijel.milincic@agrif.bg.ac.rs (D.D.M.); marmajo@agrif.bg.ac.rs (M.P.M.-J.); akostic@agrif.bg.ac.rs (A.Ž.K.)

² Institute for Biological Research “Siniša Stanković”-National Institute of Republic of Serbia, University of Belgrade, Bulevar Despota Stefana 142, 11060 Belgrade, Serbia; nena2000@ibiss.bg.ac.rs

* Correspondence: sofija@ymail.com

† Presented at the 2nd International Electronic Conference on Plant Sciences—10th Anniversary of Journal Plants, 1–15 December 2021; Available online: <https://iecps2021.sciforum.net/>.

Abstract: *Athamanta turbith* (L.) Brot. is the endemic flowering plant from the Apiaceae family originated from Đetinja Canyon (Serbia). The aim of this study was to determine the content of selected plant bioactive compounds present in rhizome, vegetative shoot, and inflorescence. Extraction was performed in 80% methanol as a solvent with two different approaches: powdered plant material was extracted with solvent for 3 h in ratio 1:10 without (classical solvent extraction, SE,) and with application of ultrasound (ultrasound-assisted extraction, UAE). Analysis of total phenolic content (TPC), total flavonoid content (TFC) and total hydroxycinnamic acid derivative content (HCA) was performed via spectrophotometric methods. The inflorescence had the highest TPC in UAE obtained-extract (2.73 ± 0.13 mg GAE/g), as well as the highest TFC (1.56 ± 0.02 mg/g QE) and HCA (1.45 ± 0.11 mg/g CGAE) in SE-prepared extract where GAE, QE and CGAE are gallic acid equivalents, quercetin equivalents and chlorogenic acid equivalents respectively. The lowest amount of TPC, TFC and HCA was detected in the rhizome regardless of extraction methods. Additionally, antioxidant properties of extracts were determined with five assays: ABTS⁺, DPPH[·], ferric reducing power (FRP), in vitro phosphomolybdenum total antioxidant capacity (TAC), and cupric reducing antioxidant capacity (CUPRAC). The inflorescence had the highest antioxidant activity in both quencher assays (at $\gamma = 0.1$ g/mL) with 92.1% of inhibition for ABTS⁺ (UAE extract,) and 77.7% inhibition of DPPH[·] (for both extracts). In addition, it exhibited the highest FRP (18.4 mg/g AAE, SE extract and CUPRAC (~40 mg/g AAE for both extracts) values where AAE stands for ascorbic acid equivalents, whereas the rhizome had the lowest values for all antioxidant assays concerning both SE and UEA. In conclusion, exhibited antioxidant properties are mostly in line with the determined content of selected bioactive compounds. Further statistical analysis is applied to confirm/oppose this.

Keywords: *Athamanta turbith*; bioactive compounds; antioxidant activity; Đetinja Canyon



Citation: Kilibarda, S.N.; Vuković, S.Z.; Milinčić, D.D.; Mačukanović-Jocić, M.P.; Jarić, S.; Kostić, A.Ž. Phytochemical and Antioxidant Properties of *Athamanta turbith* (L.) Brot Collected from Serbia. *Biol. Life Sci. Forum* **2022**, *11*, 30. <https://doi.org/10.3390/IECPS2021-11947>

Academic Editor: Giedrė Samuoliene

Published: 30 November 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Plants are a remarkable source of different bioactive compounds with proved antioxidant properties. In order to obtain valuable natural compounds from plants it is important to apply adequate extraction procedures including appropriate solvents as well as extraction techniques. Among several procedures classical solvent extraction (SE) and ultrasound-assisted extraction (UAE) stand out as the most present. It is proven that UAE can improve extraction yield of different plant bioactive compounds such as carotenoids [1] and/or phenolics [2]. Recently there has been a trend of examining endemic plants as novel valuable sources of different bioactives. Among them is *Athamanta turbith* (L.) Brot., flowering plant from the Apiaceae family that grows in Đetinja Canyon. River Đetinja

springs on the slopes of Tara Mountain, near village Kremna in Western Serbia (Figure 1). This distinguished area has great historical, cultural, geological, ecological and biological importance. *A. turbith* L. prefers chalky, dry soils and gravel, which are exposed to sun. The average height of the plant is 30 cm. Stems are branched, bright green leaves are triangular, two to four times pinnated. In summer months, this lithophyte forms inflorescences that belong to compound umbel type with tiny, white, star-shaped flowers as shown in Figure 1.




Figure 1. Location of Djetinja Canyon (left) and *Athamanta turbith* L. Brot (right).

In the literature there is still a lack of data about phytochemical analysis for this plant. Accordingly, the main objective of this study was to determine the content of selected bioactive compounds present in three distinct plant parts—rhizome, vegetative shoot, and inflorescence by application of usual spectrophotometric methods. For this purpose, two extraction techniques were performed: solvent extraction (SE) and ultrasound-assisted extraction (UAE) in order to obtain maximal yield of bioactive compounds. After that, antioxidant properties of prepared extracts were analyzed and correlated with determined bioactive compounds of *A. turbith* L.

2. Material and Methods

Plant material (rhizome, vegetative shoot, and inflorescence) of *A. turbith* was collected from Djetinja Canyon (western Serbia) and further used to prepare extracts.

2.1. Preparation of Extracts

Extraction was performed in 80% methanol as the solvent (plant material: solvent ratio, 1:10) with two different approaches: without (classical solvent extraction, SE) and with application of ultrasound (ultrasound-assisted extraction, UAE). In the case of SE, samples were placed in plastic cuvettes, soaked with solvent, and intensively shaken with protection from light, at room temperature. For UAE, samples were prepared in the same way and then sonicated in an ultrasonic bath (VAB SB 3 LD, maximum power 440 W, operating frequency 40 Hz) at maximum frequency (40 Hz), at room temperature. The extracts were filtered through suitable filter paper. Obtained supernatants were left in a dark place, at 4 °C until further analysis.

2.2. Determination of Bioactive Compounds

Total phenolic content (TPC). Determination of TPC was performed using standard Folin-Ciocalteu (FC) method [3]. After that, the reaction mixtures were intensively vortexed and left in the dark, at room temperature, for 90 min. The absorbance was measured at 765 nm. Quantification was done on the basis of the calibration curve of gallic acid (GA). Results were expressed as mg of gallic acid equivalents (GAE) per g of dry weight (DW).

Total flavonoid content (TFC). TFC was estimated using common AlCl_3 method [4]. After a 10 min mixture incubation at room temperature, absorbance was measured against the blank (blank = acidified methanol) at 430 nm. The flavonoid content was determined using quercetin as the standard and the obtained results were expressed as mg of quercetin equivalent (QE) per g of dry sample weight.

Total dihydroxycinnamic acid derivatives (HCAs). Total HCA content was determined by application of standard Arnow's reagent [5], with a slight modification. In a 2 mL plastic tube, there were added: 0.2 mL of plant extract, 0.4 mL of 0.5 M HCl, 0.4 mL of Arnow's reagent, 0.4 mL of 2.215 M NaOH and 0.6 mL of distilled H_2O . Blank was prepared by replacing 0.2 mL plant extract with distilled water. After that, the reaction mixtures were intensively vortexed and left in the dark, at room temperature, for 20 min. The absorbance was measured at 525 nm. Chlorogenic acid (CGA) was used as standard for the calibration curve. Total HCA content was expressed as mg of CGA equivalents (CGAE) per g of DW.

2.3. Determination of Antioxidant Activity

DPPH. and ABTS^+ assays. The free radical scavenging activity of plant extracts was tested by DPPH. scavenging assay [6]. Briefly, 105 μL of extracts were mixed with 840 μL of 150 μM DPPH. solution. The reaction mixture was well vortexed and left in the dark at room temperature. After 30 min the absorbance was measured spectrophotometrically at 515 nm. Antioxidant activity of plant extracts is reflected through the decolorization of purple DPPH. solution compared to the blank. For ABTS^+ quenching ability determination 500 μL of plant extract was mixed with 1 mL of green ABTS^+ solution and well vortexed [7]. After a 7-min incubation of reaction mixture in dark place, the absorbance was measured, against the blank at 734 nm. For both assays the percentage inhibition of radicals caused by the action of plant phytochemicals was calculated by using the following equation:

$$\% \text{ inhibition} = [A_b - A_s] / A_b * 100$$

A_b —the absorbance of blank; A_s —the absorbance of the sample extracts.

TAC assay. Total antioxidant capacity (TAC) was determined spectrophotometrically [8]. Namely, 0.3 mL of plant extract was mixed with 3 mL of phospho-molybdate reagent. The reaction mixtures were heated for 90 min at 95 °C to complete the reaction. After cooling reaction mixtures at room temperature, the absorbance was measured at 695 nm. Ascorbic acid (AA) was used as the standard and the obtained results were expressed as mg of ascorbic acid equivalents (AAE) per g of DW.

FRP assay. For ferric reducing power (FRP) assay [9] 0.5 mL of plant extract, 0.5 mL of 0.2 M sodium phosphate buffer (pH 6.6) and 0.5 mL of 1% potassium ferricyanide, were mixed in a glass tube. The reaction mixtures were incubated at 50 °C, for 20 min. Then 0.5 mL of 10% trichloroacetic acid was added and the total mixture was centrifuged ($4000 \times g$) for 5 min. Next, 1.5 mL of clear supernatant was withdrawn from glass tube and mixed with 1.5 mL of distilled water and 0.3 mL of 0.1% of ferric chloride. The absorbance of colored solutions was measured at 700 nm. The calibration curve was prepared with ascorbic acid as the standard and the results were expressed as mg of ascorbic acid equivalents (AAE) per g of DW.

CUPRAC assay. Cupric ion antioxidant capacity (CUPRAC) assay was conducted by the spectrophotometric method [10]. Briefly, 0.35 mL of plant extract, 0.35 mL of 0.01 M CuCl_2 , 0.35 mL of 0.0075 M neocuproine solution (previously dissolved in concentrated absolute ethanol), 0.350 mL 1 M ammonium acetate buffer solution were mixed in a plastic tube. The reaction mixtures were left in the dark for 30 min and then the absorbance of solutions was read at 450 nm. Ascorbic acid (AA) was used to set up the standard curve. The obtained results were expressed as mg of ascorbic acid equivalents (AAE) per g of dry sample weight.

2.4. Statistics

All assays were carried out in triplicate ($n = 3$). The results were shown as means \pm standard deviation (SD). Statistical analyses of the data were performed using the STATISTICA 12.0. Statistical significance was evaluated employing Tukey's test. Differences were considered significant at $p < 0.05$. Correlation analysis between analyzed phenolic classes and antioxidant properties, that is, between different types of extraction for specific part of investigated *A. turbith* plant was evaluated using Pearson's correlation coefficient (r), at $p < 0.01$.

3. Results and Discussion

3.1. Phytochemical Composition

The content of total phenolics (TPC), total flavonoids (TFC) and total dihydroxycinnamic acid derivatives (HCAs) observed in rhizome, vegetative shoot, and inflorescence extracts of *A. turbith* is shown in Table 1. The inflorescence had the highest TPC in UAE obtained-extract, as well as the highest TFC and HCA in SE-prepared extract. The obtained results are in line with literature where it was reported that flower of Apiaceae plants had higher TPC compared to vegetative parts and rhizome [11]. In general, the extract of inflorescence obtained by UAE had significantly higher ($p < 0.05$) content of TPC and TFC than extract obtained by SE. However, there was no significant difference in the content of HCAs, achieved in the inflorescence, for both extracts. The lowest amount of TPC and HCAs was detected in the rhizome regardless of the extraction methods ($p < 0.05$), while the flavonoids were not detected.

Table 1. Phytochemical composition of *Athamanta turbith*.

Sample	Extraction Technique	TPC * (mg/g GAE)	TFC (mg/g QE)	HCAs (mg/g CGAE)
Inflorescence	UAE	2.73 \pm 0.13 ^{a,**}	1.36 \pm 0.02 ^a	1.41 \pm 0.004 ^a
	SE	1.95 \pm 0.15 ^b	1.56 \pm 0.02 ^b	1.45 \pm 0.11 ^a
Vegetative shoot	UAE	1.06 \pm 0.02 ^c	0.70 \pm 0.002 ^c	1.07 \pm 0.009 ^b
	SE	0.87 \pm 0.01 ^c	0.53 \pm 0.05 ^d	0.85 \pm 0.008 ^c
Rhizome	UAE	0.37 \pm 0.03 ^d	n.d.	0.71 \pm 0.00 ^d
	SE	0.40 \pm 0.01 ^d	n.d.	0.66 \pm 0.00 ^d

* TPC—total phenolic content; TFC—total flavonoid content; HCA—total dihydroxycinnamic acid derivative content; GAE—gallic acid equivalents; QE—quercetin equivalents; CGAE—chlorogenic acid equivalents; n.d.—not detected. ** Different superscript letters (a–d) in a same column indicate significant differences at $p < 0.05$.

Antioxidant properties of *A. turbith* extracts determined with five assays: ABTS⁺, DPPH[·], ferric reducing power (FRP), in vitro phosphomolybdenum total antioxidant capacity (TAC) and cupric reducing antioxidant capacity (CUPRAC) are shown in Table 2.

Table 2. Antioxidant properties of *A. turbith* extracts.

Sample	Extraction Technique	ABTS ⁺ * (% inh.)	DPPH [·] (% inh.)	TAC (mg/g AAE)	CUPRAC (mg/g AAE)	FRP (mg/g AAE)
Inflorescence	UAE	51.43 \pm 0.06 ^{a,**}	77.68 \pm 0.55 ^a	3.60 \pm 0.07 ^a	39.15 \pm 3.03 ^a	11.06 \pm 0.52 ^a
	SE	92.11 \pm 0.48 ^b	77.77 \pm 0.57 ^a	3.53 \pm 0.29 ^a	41.83 \pm 1.29 ^a	18.37 \pm 1.70 ^b
Vegetative shoot	UAE	23.67 \pm 0.00 ^c	33.86 \pm 0.14 ^b	1.75 \pm 0.00 ^b	12.52 \pm 1.10 ^b	1.59 \pm 0.09 ^{c,d}
	SE	34.00 \pm 0.06 ^d	50.34 \pm 0.41 ^c	1.54 \pm 0.01 ^b	8.42 \pm 1.03 ^b	3.33 \pm 0.30 ^c
Rhizome	UAE	13.91 \pm 0.13 ^e	10.00 \pm 0.17 ^d	1.72 \pm 0.14 ^b	n.d.	0.46 \pm 0.06 ^d
	SE	9.14 \pm 0.66 ^f	5.67 \pm 0.31 ^e	1.78 \pm 0.007 ^b	n.d.	0.46 \pm 0.05 ^d

* ABTS⁺—2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) radical cation; DPPH[·]—2,2-diphenylpicrylhydrazyl cation; TAC—total antioxidant capacity determined via in vitro phosphomolybdenum assay; CUPRAC—Cupric Reducing Antioxidant Capacity; FRP—Ferric Reducing Power; AAE—ascorbic acid equivalents. ** Different superscript letters (a–f) in a same column indicate significant differences at $p < 0.05$.

The inflorescence had the highest antioxidant activity in both quencher assays (at $\gamma = 0.1$ g/mL) with 92.1% of inhibition for ABTS^{•+} (UAE extract,) and 77.7% inhibition of DPPH[•] (for both extracts). In addition, it exhibited the highest FRP (18.4 mg/g AAE, SE extract), CUPRAC (~40 mg/g AAE for both extracts) and TAC (~35 mg/g AAE for both extracts). The inflorescence extracts obtained by SE and UAE show a statistically significant difference ($p < 0.05$) between antioxidant activity, only in the ABTS^{•+} and FRP assays. The rhizome had the lowest values for all antioxidant assays concerning both SE and UEA. Differences in antioxidant activity between SE and UAE-obtained rhizome extracts were statistically significant ($p < 0.05$) for ABTS^{•+} and DPPH[•] assays.

3.2. Correlation Analysis

Correlation analysis showed a significant ($p < 0.01$) positive correlation between TPC and CUPRAC ($r = 0.95$); while TFC showed a positive correlation between DPPH[•] ($r = 0.96$), FRP ($r = 0.92$) and CUPRAC ($r = 0.98$). Further, HCAs show positive correlation with DPPH[•] ($r = 0.92$) and CUPRAC ($r = 0.98$). Based on the correlation results, it can be concluded that extracted dihydrocinnamic acid derivatives and flavonoids from different parts of the plant, together or individually show good cupric reducing capacity (CUPRAC), while, on the other hand, only flavonoids contribute to effective ferric reducing capacity (FRP). In addition, individual flavonoids and dihydrocinnamic acid derivatives of plant rhizome, vegetative shoot, and inflorescence, have good hydrogen donor properties and effective DPPH[•] scavenging capacity. Furthermore, CUPRAC significantly correlated with other antioxidant assays such as FRP ($r = 0.94$), DPPH[•] ($r = 0.94$), and TAC ($r = 0.95$). This may mean that the same phenolic compounds or phenolic compounds which possess the same mechanism of antioxidant action (extracted with SE and UAE), contribute to the strong correlation of CUPRAC with the above methods. However, based on analyzed phenolic classes and antioxidant properties, both types of extraction (solvent and ultrasound-assisted extraction) strongly correlate for each part of the plant individually (rhizome, vegetative shoot, and inflorescence), which means that both methods can potentially be used to extract phenolics from different parts of the *A. turbith* plant, without significant differences in the content of different phenolic classes and antioxidant properties.

4. Conclusions

Phytochemical analysis of different parts of *A. turbith* revealed distinctions in phenolic composition with inflorescence as the best source of bioactive compounds. There was no clear influence of ultrasound assisted extraction on the content of total phenolics, flavonoids and dihydrocinnamic acid derivatives. All examined extracts exhibited significant antioxidant activity examined through five different assays. Correlation analysis confirmed strong connection between phenolics (flavonoids and dihydrocinnamic acid derivatives, in particular) and several antioxidant assays such as CUPRAC and DPPH[•] assays.

Supplementary Materials: The following are available online at <https://sciforum.net/manuscripts/11947/slides.pdf>.

Author Contributions: Conceptualization, A.Ž.K. and S.N.K.; methodology, A.Ž.K.; formal analysis, D.D.M.; resources, S.J. and M.P.M.-J.; writing, S.Z.V.; writing—review and editing, S.N.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Acknowledgments: This work was supported by the Ministry of Education, Science, and Technological Development of the Republic of Serbia (Grant No. 451-03-9/2021-14/200116).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Alcántara, C.; Žugčić, T.; Abdelkebir, R.; García-Pérez, J.V.; Režek Jambrak, A.; Lorenzo, J.M.; Collado, M.C.; Granato, D.; Barba, F.J. Effects of ultrasound-assisted extraction and solvent on the phenolic profile, bacterial growth, and anti-inflammatory/antioxidant activities of Mediterranean olive and fig leaves extracts. *Molecules* **2020**, *25*, 1718. [[CrossRef](#)] [[PubMed](#)]
2. Berkani, F.; Serralheiro, M.L.; Dahmoune, F.; Ressaissi, A.; Kadri, N.; Remini, H. Ultrasound assisted extraction of phenolic compounds from a jujube by-product with valuable bioactivities. *Processes* **2020**, *8*, 1441. [[CrossRef](#)]
3. Ng, A.; Parker, M.L.; Parr, A.J.; Saunders, P.K.; Smith, A.C.; Waldron, K.W. Physicochemical characteristics of onion (*Allium cepa* L.) tissues. *J. Agric. Food Chem.* **2000**, *48*, 5612–5617. [[CrossRef](#)] [[PubMed](#)]
4. Santas, J.; Carbo, R.; Gordon, M.; Almajano, M. Comparison of the antioxidant activity of two Spanish onion varieties. *Food Chem.* **2008**, *107*, 1210–1216. [[CrossRef](#)]
5. Fraisse, D.; Felgines, C.; Texier, O.; Lamaison, J.-L. Caffeoyl derivatives: Major antioxidant compounds of some wild herbs of the Asteraceae family. *Food Nutr. Sci.* **2011**, *2*, 181–192. [[CrossRef](#)]
6. Gawron-Gzella, A.; Królikowska, A.; Pietrzak, M. Antioxidant activity of teas obtained from leaves of *Camellia sinensis* (L.) Kuntze in course of various production processes available on Polish market. *Herba Pol.* **2018**, *64*, 60–67. [[CrossRef](#)]
7. Rajurkar, S.N.; Hande, M.S. Estimation of phytochemical content and antioxidant activity of some selected traditional Indian medicinal plants. *Indian J. Pharm. Sci.* **2011**, *73*, 146–151. [[CrossRef](#)]
8. Prieto, P.; Pineda, M.; Aguilar, M. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific application to the determination of vitamin E. *Anal. Biochem.* **1999**, *269*, 337–341. [[CrossRef](#)] [[PubMed](#)]
9. Nibir, Y.M.; Sumit, A.F.; Akhand, A.A.; Ahsan, N.; Hossain, M.S. Comparative assessment of total polyphenols, antioxidant and antimicrobial activity of different tea varieties of Bangladesh. *Asian Pac. J. Tropic. Biomed.* **2017**, *7*, 352–357. [[CrossRef](#)]
10. Apak, R.; Güçlü, K.; Özyürek, M.; Çelik, S.E. Mechanism of antioxidant capacity assays and the CUPRAC (cupric ion reducing antioxidant capacity) assay. *Microchim. Acta* **2008**, *160*, 413–419. [[CrossRef](#)]
11. Sayed-Ahmad, B.; Talou, T.; Saad, Z.; Hijazi, A.; Merah, O. The Apiaceae: Ethnomedical family as source for industrial uses. *Ind. Crop. Prod.* **2017**, *109*, 661–671. [[CrossRef](#)]