



Yoghurt-Like Drink Enriched with Lactobacilli and Bifidobacteria [†]

Annalisa d'Amelio, Clelia Altieri  and Daniela Campaniello *

Department of Agriculture, Food, Natural Resources and Engineering (DAFNE), University of Foggia, 71122 Foggia, Italy; annalisa.damelio@unifg.it (A.d.); clelia.altieri@unifg.it (C.A.)

* Correspondence: daniela.campaniello@unifg.it

[†] Presented at the 4th International Electronic Conference on Foods, 15–30 October 2023; Available online: <https://foods2023.sciforum.net/>.

Abstract: To develop an enriched fermented milk, an experimental design was used in the first step by combining *Lactiplantibacillus plantarum* (isolated from oenological matrix) and *Bifidobacterium animalis* subsp. *lactis* strains, at different inoculum levels (4, 5 and 6 log CFU/mL) and temperatures (30, 35 and 40 °C) to study the effects on the acidification. Then, for the realization of the yoghurt-like drink, 5 log CFU/mL of both strains was inoculated in milk + 10% honey at 35 °C for 24 h; the fermented milk was stored at 4 °C for 50 days. The microbial concentrations were always >8 log CFU/mL.

Keywords: probiotics; fermented milk; experimental design

1. Introduction

The creation of a product requires a careful and scrupulous study of all the variables and an evaluation of how these can influence the final product. In the literature, many papers focus on the production of drinks with a probiotic or prebiotic added [1–3]. This type of product responds to the constant demand of consumers who are increasingly attentive to the nutritional and health characteristics of foods. Yoghurt is one of the oldest foods, and today, it has become a trendy product, in terms of nourishment and versatility, considered a healthy protein food and a valid alternative to milk [4]. Like other fermented drinks, yoghurt brings several benefits to our body; it has anti-inflammatory and purifying properties, the acidity of the yoghurt favours the growth of an intestinal bacterial biota capable of counteracting the putrefactive phenomena that occur in the intestine. Many functions are performed by the synergistic action of *Streptococcus thermophilus* and *Lactobacillus delbrueckii* subsp. *Bulgaricus* [4]; however, different bacteria (e.g., probiotics) can be added.

The aim of this work was the formulation of a yoghurt-like drink with Lactobacilli and Bifidobacteria, through the use of microorganisms isolated from oenological matrices of the Apulian territory; an experimental design, the Centroid, was used to optimize the product.

2. Materials and Methods

Microorganisms and Milk: *Bifidobacterium animalis* subsp. *lactis* DSM 10140 (Deutsche Sammlung von Mikroorganismen und Zellkulturen's collection, Braunschweig, Germany) and one strain of *Lactiplantibacillus plantarum* coded as 33 isolated from Apulian (Italy) oenological matrix, were used in this research. The strains were stored at −20 °C in MRS broth (Oxoid, Milan, Italy) and MRS Broth + 0.5% cysteine, for *Lactiplantibacillus* and *Bifidobacterium*, respectively, with 33% (V/V) of sterile glycerol (J.T. Baker, Milan, Italy). Before each assay, the bacteria were grown twice in the same medium at 30 (Lpb. *plantarum*) and 37 °C (*B. animalis* subsp. *lactis*) for 24 h; then, the cultures were centrifuged at 4000× g for 10 min at 4 °C, the supernatant was discarded, and the pellet re-suspended in distilled water. Apulian fresh whole pasteurized homogenized cow's milk (3.2 g/L protein; 4.9 g/L carbohydrates; 3.6 g/L fats) purchased from a local dealer was used.



Citation: d'Amelio, A.; Altieri, C.; Campaniello, D. Yoghurt-Like Drink Enriched with Lactobacilli and Bifidobacteria. *Biol. Life Sci. Forum* **2023**, *26*, 34. <https://doi.org/10.3390/Foods2023-15099>

Academic Editor: Antonio Bevilacqua

Published: 14 October 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Experimental design—Centroid: The optimization of the production of the yoghurt-like drink was performed according to a centroid design [5,6]. In this research, the independent variables were the inoculum level of *Lpb. plantarum* 33 (L) and of *B. animalis* subsp. *lactis* (B), and temperature (T) (Table 1).

Table 1. Combination, coded levels and values of Centroid.

Combinations	Coded Levels			Values		
	B *	L **	T ***	B	L	T
A	1	0	0	6	4	30
B	0	1	0	4	6	30
C	0	0	1	4	4	40
D	0.5	0.5	0	5	5	30
E	0.5	0	0.5	5	4	35
F	0	0.5	0.5	4	5	35

* B = *Bifidobacterium animalis* subsp. *lactis*, CFU/mL; ** L = *Lactiplantibacillus plantarum* 33, CFU/mL; *** T = temperature, °C.

According to the design, *Lpb. plantarum* and *B. animalis* subsp. *lactis* were inoculated to 4, 5 and 6 log CFU/mL in 100 mL of pasteurized milk supplemented with honey at 10% (V/V); then, the samples were incubated at 30, 35 or 40 °C for 7 days. For each combination, microbiological sampling was carried out and the acidification was periodically monitored through pH measuring using a pH-meter (Crison, Barcelona, Spain). Each analysis was performed in duplicate.

Modeling: Microbial growth data (log CFU/mL), were plotted on an Excel worksheet and processed as mean value $\pm$ standard deviation.

Concerning pH, data were modeled as acidification (Δ pH), i.e., pH decrease referred to the beginning of the experiment; then, it was used as a dependent variable for a primary modeling through the “lag-exponential model” of van Gerwen and Zwietering [7] and Baty and Delignette-Muller [8] to obtain α , the time before the beginning of the acidification kinetic (h); $d_{\max}$ is the maximal acidification rate (1/h), and Δ pH $_{\max}$ is the maximum level of acidification.

Successively, Δ pH $_{\max}$ e $d_{\max}$ were used as input values for a multiple regression approach; temperature and inoculum levels of both strains were used as independent variables. The analysis was performed through the software Statistica for Windows (StatSoft, Tulsa, OK, United States). The model was built by using the option “quadratic,” for the evaluation of the individual (“*B. animalis* subsp. *lactis* inoculum level”, “*Lpb. plantarum* 33 inoculum level” and “temperature”) and interactive effects (“*B. animalis* subsp. *lactis* inoculum level * *Lpb. plantarum* 33 inoculum level”; “*B. animalis* subsp. *lactis* inoculum level * temperature,”; and “temperature * *Lpb. plantarum* 33 inoculum level”). Finally, the effect of each independent variable on the fitting parameters of the acidification kinetic (Δ pH $_{\max}$ and $d_{\max}$) was evaluated through the individual desirability functions, as reported by Speranza et al. [3].

Yoghurt-like realization: an amount of 5 log CFU/mL of *Lpb. plantarum* 33 and *B. animalis* subsp. *lactis* was inoculated in a 10 mL sample composed of 9 mL of milk and 1 mL honey and incubated at 35 °C for 24 h; then, the samples were stored at 4 °C. Microbiological analyses and pH measurements were periodically assessed. For microbiological analyses, the following media were used: MRS Agar acidified to pH 5 for *Lpb. plantarum* 33, and MRS Agar added with a solution of 4 antibiotics (Paromomycin sulphate 0.01 g, Neomycin sulphate 0.005 g, Lithium 0.15 g, Nalidixic acid 0.75 g) for *B. animalis* subsp. *lactis*. The experiments were repeated twice on two independent samples [9,10].

3. Results

The microbial strains used, *B. animalis* subsp. *lactis* DSM 10140 and *Lpb. plantarum* 33 were preliminary analysed to evaluate their acidifying capability and subjected to adaptation trials to improve their performances [11]; they were then inoculated in milk + honey at 10% (data not yet published). Successively, the product optimization was realised through an experimental design (Centroid).

3.1. Centroid

Table 2 shows the combinations of the experimental design and the performances of both strains referred to $\Delta\text{pH}_{\text{max}}$, d_{max} and α . $\Delta\text{pH}_{\text{max}}$ varied from a minimum value of 2.20 (combination C) to a maximum of 2.59 (combination E); the acidification rate (d_{max}) ranged from 0.08 to 0.19 g^{-1} . The parameter α was statistically significant only for combinations A, B, C and D, ranging from 3.53 ± 0.61 (combination A) to 6.59 ± 1.15 days. The high standard error for B, C and D combinations suggests that acidification could occur even more rapidly.

Table 2. Acidification of *Lpb. plantarum* 33 and *B. animalis* subsp. *lactis*: fitting parameters of the “lag-exponential equation” (mean $\pm$ standard error).

Combinations				
	$\Delta\text{pH}_{\text{max}}$	d_{max}	α	R
A	2.38 ± 0.01	0.08 ± 0.00	3.53 ± 0.61	0.999
B	2.24 ± 0.05	0.09 ± 0.00	5.19 ± 2.10	0.994
C	2.20 ± 0.07	0.19 ± 0.02	5.34 ± 2.00	0.980
D	2.34 ± 0.03	0.10 ± 0.00	6.59 ± 1.15	0.996
E	2.59 ± 0.06	0.11 ± 0.01	-	0.995
F	2.52 ± 0.07	0.11 ± 0.00	-	0.994

$\Delta\text{pH}_{\text{max}}$ = maximum acidification (decrease in pH at the end of the assay); d_{max} = maximal acidification rate ($\Delta\text{pH}/\text{day}$); α = the time before the beginning of the acidification kinetic (d); R = regression coefficient.

Successively, $\Delta\text{pH}_{\text{max}}$ and d_{max} were used as dependent variables for a multiple regression approach. The first output was the table of the standardized effects, showing the statistical weight and the significance of each individual (inoculum of both microorganisms and temperature) and interactive factors.

Table 3 shows the standardized effects of the three variables on $\Delta\text{pH}_{\text{max}}$. As individual term, *B. animalis* showed the greatest effect (73.814), followed by *Lpb. plantarum* 33 (69.330) and temperature (68.251), while the interaction “*Lpb. plantarum*–temperature” was the most significant, followed by “*B. animalis* subsp. *lactis*–temperature”. Concerning d_{max} , the maximum effect was attributable to the temperature followed by the level of inoculum of *Lpb. plantarum* and *B. animalis* strains. The “*Lpb. plantarum*–temperature” and “*B. animalis* subsp. *lactis*–temperature” interactions were also significant.

Table 3. Standardized effects of the inoculum of *B. animalis* subsp. *lactis* DSM 10140 [B], *Lpb. plantarum* [L] and temperature [T] on the maximum acidification ($\Delta\text{pH}_{\text{max}}$) and on the acidification rate (d_{max}).

EFFECTS	$\Delta\text{pH}_{\text{max}}$	d_{max}
[B]	73.814	12.746
[L]	69.330	13.429
[T]	68.251	28.331
[B] $\times$ [L]	- (A)	-
[B] $\times$ [T]	7.385	−2.999
[L] $\times$ [T]	7.614	−3.275
R ² (B)	0.901	0.930

(A) Not significant; (B) determination coefficient adjusted for the multiple regression.

Other outputs of centroid are the ternary plots; Figure 1a,b show the triangular surface for the effects of *B. animalis* subsp. *lactis* (B), *Lpb. plantarum* (L) and temperature (T) on the maximum acidification ($\Delta\text{pH}_{\text{max}}$) (Figure 1a) and on the acidification rate (d_{max}) (Figure 1b). The model predicted the maximum acidification ($\Delta\text{pH}_{\text{max}} = 2.5$) when either *Lpb. plantarum* or *B. animalis* and temperature were at the coded levels 0.5 (that is the microorganisms at 5 log CFU/mL and temperature at 35 °C) (Figure 1a). Concerning d_{max} (Figure 1b) the model predicted the highest values ($d_{\text{max}} = 0.18$) at the coded level 1 of the temperature (40 °C).




Figure 1. Triangular graph relating to the effects of *B. animalis* subsp. *lactis* (B), *Lpb. plantarum* (L) and temperature (T) on the $\Delta\text{pH}_{\text{max}}$ (a) and d_{max} (b).

Triangular surfaces provide a quantitative assessment of interactive effects, but not of each individual term. This type of evaluation is possible through the desirability profile.

The desirability profile of $\Delta\text{pH}_{\text{max}}$ (Figure 2a) shows that both for *B. animalis* subsp. *lactis* and *Lpb. plantarum*, the increase in the inoculum level negatively affected acidification with a decreased performance of the strains at coded levels 0.75 or 1 (corresponding to 5.50 and 6 log CFU/mL). Concerning temperature, $\Delta\text{pH}_{\text{max}}$ was minimal at 30 °C and also decreased at coded levels 0.75 and 1 (corresponding to 37.5 and 40 °C).




Figure 2. Prediction (upper side) and desirability profiles (down side) for the effect of inoculum level (log CFU/mL) of *B. animalis* subsp. *lactis* (B), *Lpb. plantarum* (L) and temperature as °C (T) on $\Delta\text{pH}_{\text{max}}$ (a) and d_{max} (b).

The desirability profile of the acidification rate ($d_{\max}$) (Figure 2b) shows that both *B. animalis* subsp. *lactis* and *Lpb. plantarum* 33 reduced the acidification rate at the inoculum coded level 1 (corresponding to 6 log CFU/mL). With regard to temperature, a positive correlation was observed: as temperature increased (for values coded as 0.75 and 1 = 37.5 and 40 °C, respectively), $d_{\max}$ increased.

As shown by the desirability profiles (Figure 2a,b), the temperature acted differently on the maximum acidification and on the rate: in relation to the $\Delta pH_{\max}$, the increase in temperature led to a reduction in acidification; instead, in the case of $d_{\max}$, the increase in temperature was positively correlated with acidification rate. This phenomenon known as parameters uncoupling [12] highlights that increasing the temperature increases the acidification rate; however, temperature cannot reach the maximum values because a reduction in acidification capability could occur.

3.2. Yoghurt-Like Drink

The main result of optimization of Centroid design was the best combination for obtaining the final product; as a result of modelling, the conditions were identified by the software as follows: 5 log CFU/mL of *Lpb. plantarum* 33 and of *B. animalis* subsp. *lactis* 10,140, and temperature at 35 °C for 24 h. The concentration of both strains reached >8 log CFU/mL. Then, the samples were stored at 4 °C for 50 days and both strains were able to survive and never resulted below the criticality threshold (7 log CFU/mL) (Table 4). Furthermore, the pH did not undergo significant changes during storage (initial pH = 4.84; final pH = 4.22). Finally, the texture, odor and color of the yoghurt-like drink were optimal.

Table 4. Cellular concentration (log CFU/mL) *Lpb. plantarum* 33 and *B. animalis* subsp. *lactis* DSM 10140 inoculated into the yoghurt-like drink and stored at a temperature of 4 °C. Lowercase letters indicate statistically significant differences along the column (effect of time for each strain); capital letters refer to the differences between the 2 strains at the same time of analysis (one-way ANOVA and Tukey's test, $p < 0.05$). Data are expressed as mean values $\pm$ standard deviation.

Time (Days)	33	10,140
0	8.78 $\pm$ 0.00 ^{a,A}	8.40 $\pm$ 0.00 ^{a,B}
1	8.74 $\pm$ 0.01 ^{a,A}	8.90 $\pm$ 0.02 ^{b,A}
4	8.90 $\pm$ 0.03 ^{a,A}	8.85 $\pm$ 0.00 ^{b,A}
6	8.93 $\pm$ 0.02 ^{a,A}	8.72 $\pm$ 0.03 ^{b,A}
19	8.88 $\pm$ 0.03 ^{a,A}	8.78 $\pm$ 0.02 ^{b,A}
26	8.85 $\pm$ 0.00 ^{a,A}	8.88 $\pm$ 0.03 ^{b,A}
33	8.52 $\pm$ 0.02 ^{b,A}	8.60 $\pm$ 0.00 ^{a,A}
40	8.74 $\pm$ 0.01 ^{a,A}	8.88 $\pm$ 0.02 ^{b,A}
50	8.74 $\pm$ 0.04 ^{a,A}	8.88 $\pm$ 0.03 ^{b,A}

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/Foods2023-15099/s1>.

Author Contributions: Conceptualization, D.C.; methodology, D.C.; investigation, A.d., C.A. and D.C.; data curation, D.C.; writing—original draft preparation, A.d.; writing—review and editing, D.C. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Apulia Region-REFIN, grant number B765D6AC Soluzioni microbiche modello per la bioeconomia e la bioindustria agroalimentare pugliese (Model Microbial Solutions for the Apulian Bioeconomy and Agro-Food Bioindustry).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this article are available on request.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Mudgil, P.; Barak, S. Dairy-Based Functional Beverages. In *Milk-Based Beverages*; Grumezescu, A.M., Holban, A.M., Eds.; Woodhead Publishing: Cambridge, UK, 2019; Volume 9, pp. 67–93.
2. Perotti, M.C.; Bergamini, C.V.; Vénica, C.I.; Vélez, M.A.; Wolf, I.V.; Hynes, E. Production of Functional Milk-Based Beverages. In *Milk-Based Beverages*; Grumezescu, A.M., Holban, A.M., Eds.; Woodhead Publishing: Cambridge, UK, 2019; Volume 9, pp. 173–238.
3. Speranza, B.; Bevilacqua, A.; Campaniello, D.; Sinigaglia, M.; Musaico, D.; Corbo, M.R.; Lamacchia, C. The Impact of Gluten Friendly Flour on the Functionality of an Active Drink: Viability of *Lactobacillus acidophilus* in a Fermented Milk. *Front. Microbiol.* **2018**, *9*, 2042. [[CrossRef](#)] [[PubMed](#)]
4. Fisberg, M.; Machado, R. History of yogurt and current patterns of consumption. *Nutr. Rev.* **2015**, *73* (Suppl. 1), 4–7. [[CrossRef](#)] [[PubMed](#)]
5. Bevilacqua, A.; Corbo, M.R.; Sinigaglia, M. Design of experiments: A powerful tool in food microbiology. In *Current Research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology*; Mendez-Vilas, A., Ed.; Formatex Research Center: Badajoz, Spain, 2010; pp. 1419–1429.
6. Bevilacqua, A.; Sinigaglia, M. Food shelf life and safety: Challenge tests, prediction and mathematical tools. In *Application of Alternative Food-Preservation Technologies to Enhance Food Safety and Stability*; Bevilacqua, A., Corbo, M.R., Sinigaglia, M., Eds.; Bentham Publisher: Sharja, United Arab Emirates, 2010; pp. 161–187.
7. van Gerwen, S.J.; Zwietering, M.H. Growth and inactivation models to be used in quantitative risk assessments. *J. Food Prot.* **1998**, *61*, 1541–1549. [[CrossRef](#)] [[PubMed](#)]
8. Baty, F.; Delignette-Muller, M.L. Estimating the bacterial lag time: Which model, which precision? *Int. J. Food Microbiol.* **2004**, *91*, 261–277. [[CrossRef](#)]
9. Ghasempour, Z.; Alizadeh, M.; Bari, M.R. Optimisation of probiotic yoghurt production containing Zedo gum. *Int. J. Dairy Technol.* **2012**, *65*, 118–125. [[CrossRef](#)]
10. Al-Sahlan, S.T.G.; Khassaf, W.H.; Niamah, A.K.; Abd Al-Manhel, A.J. Date juice addition to bio-yogurt: The effects on physicochemical and microbiological properties during storage, as well as blood parameters in vivo. *J. Saudi Soc. Agric. Sci.* **2023**, *22*, 71–77. [[CrossRef](#)]
11. Perricone, M.; Corbo, M.R.; Sinigaglia, M.; Speranza, B.; Bevilacqua, A. Viability of *Lactobacillus reuteri* in fruit juices. *J. Funct. Foods* **2014**, *10*, 421–426. [[CrossRef](#)]
12. Bevilacqua, A.; Corbo, M.R.; Mastromatteo, M.; Sinigaglia, M. Combined effects of pH, yeast extract, carbohydrates and diammonium hydrogen citrate on the biomass production and acidifying ability of a probiotic *Lactobacillus plantarum* strain, isolated from table olives, in a batch system. *World J. Microbiol. Biotechnol.* **2008**, *24*, 1721–1729. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.