



# Colorimetric Evaluation of Quinoa Flour Fermented by *Monascus purpureus* Enriched with Monosodium Glutamate and Sodium Chloride <sup>†</sup>

Evelyn Quispe-Rivera <sup>1</sup>, Franz Tucta-Huillca <sup>1</sup>, Vasco Cadavez <sup>2,3</sup>, Ursula Gonzales-Barron <sup>2,3</sup>  
and Marcial Silva-Jaimes <sup>1,\*</sup>

- <sup>1</sup> Laboratorio de Microbiología de Alimentos, Universidad Nacional Agraria La Molina (UNALM), Av. La Molina s/n La Molina, Lima 15024, Peru; emich.q.r@gmail.com (E.Q.-R.); tucta.h.f@gmail.com (F.T.-H.)
- <sup>2</sup> Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal; vcadavez@ipb.pt (V.C.); ubarron@ipb.pt (U.G.-B.)
- <sup>3</sup> Laboratório Associado para a Sustentabilidade e Tecnologia em Regiões de Montanha (SusTEC), Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal
- \* Correspondence: misilva@lamolina.edu.pe
- <sup>†</sup> Presented at the 4th International Electronic Conference on Foods, 15–30 October 2023; Available online: <https://foods2023.sciforum.net/>.

**Abstract:** In the present study, the colour and C:N ratio of quinoa flours pigmented with *M. purpureus* supplemented with monosodium glutamate and sodium chloride were evaluated during 14 days of fermentation. The best values of L\*, a\* and b\* corresponded to the eighth day. This research showed that the pigmented flour produced by solid state fermentation of quinoa by *M. purpureus* showing variations in the colour red along with the C:N ratio during the fermentation time, resulting in a product with good visual sensory attributes that can be used to develop new naturally pigmented products with potential functional characteristics.

**Keywords:** *Chenopodium quinoa*; CIELAB; pseudocereal; carbon:nitrogen ratio



**Citation:** Quispe-Rivera, E.; Tucta-Huillca, F.; Cadavez, V.; Gonzales-Barron, U.; Silva-Jaimes, M. Colorimetric Evaluation of Quinoa Flour Fermented by *Monascus purpureus* Enriched with Monosodium Glutamate and Sodium Chloride. *Biol. Life Sci. Forum* **2023**, *26*, 46. <https://doi.org/10.3390/Foods2023-15082>

Academic Editor: Efstathios Giaouris

Published: 14 October 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The source of natural pigments from microorganisms such as bacteria, algae and filamentous fungi are good alternatives because manipulating them is more controllable and they are suitable for large-scale production. Of the pigment-producing microorganisms, fungi are considered to be the best source due to their ability to synthesise more soluble and stable pigments [1].

*Monascus* is a fungus widely used in Asian countries because it produces secondary metabolites of interest in the food industry, the most important of which are the red pigments that serve to colour foods and improve their appearance [2,3]. It also has good antioxidant, anti-inflammatory and anticarcinogenic properties, which is why it has received widespread attention in solid fermentation studies [3].

A nitrogen source, such as monosodium glutamate, is exploited by *Monascus*, which is used to evaluate the pigment synthesis kinetics and mycelial development of the fungus, where its addition in solid state fermentation enhances the production of water-soluble pigments [4,5]. Additionally, sodium chloride, a common salt compound, produces significant changes in red pigment production and growth of *Monascus* [6]. Therefore, it was considered important in this research to evaluate the colour of quinoa flour fermented by *M. purpureus* supplemented with monosodium glutamate and sodium chloride.

## 2. Materials and Methods

### 2.1. Strain *Monascus purpureus* CECT 2955

The *M. purpureus* strain was acquired from the Spanish Type Culture Collection (CECT). It was previously resuspended and seeded in PDA (potato dextrose agar) at 30 °C for 7 days, then seeded in QFA (quinoa flour agar) at pH 6, then incubated at 30 °C for 7 days. A stock inoculum containing a suspension of  $1.0 \times 10^6$  spores/mL was prepared and used for solid state fermentation [7].

### 2.2. Fermentation of Quinoa Grains by *M. purpureus*

Quinoa was used as substrate where 30 g of quinoa grains, NaCl at 0.05, 0.10, 0.20 and 0.40% (*w/w*) and monosodium glutamate at 1.0% (*w/w*) were added to 25 mL of distilled water per flask; it was sterilised in an autoclave (PRESOCLAVE III 80, J.P. SELECTA, s.a., Barcelona, Spain) at 121 °C for 15 min. Fermentation of the quinoa grains was carried out by adding 1 mL of *M. purpureus* spore suspension to the sterile substrate previously cooled to room temperature. It was placed in an incubator (ILW, Pol Eko, Poland) at 30 °C for 0, 2, 4, 6, 8, 10, 12 and 14 days. At the end of the fermentation time, each treatment was dried at 65 °C until a constant weight was reached. It was then milled to obtain pigmented quinoa flour.

### 2.3. Colorimetric Analysis of CIELAB

The evaluations were carried out in the CIELAB ( $L^*$ ,  $a^*$ ,  $b^*$ ) colour space [8] where the colour of the red fermented quinoa flours was analysed at room temperature with a CM-5 colorimeter (Minolta Camera Co., Osaka, Japan), with a D65 light source and an observation angle of 10°. A standard white plate was used for calibration before measurements were taken. One reading was taken per sample placed inside the plate for four replicates per treatment.

### 2.4. Determination of the C:N Ratio

The carbon and nitrogen content were determined by the Walkley–Black and Kjeldahl methods, respectively, where the samples were the fermented quinoa flours previously dried for analysis. The results were divided to find the C:N ratio. The analysis was performed for days 0, 2, 4, 6, 8, 10, 12 and 14 at four levels of sodium chloride.

### 2.5. Statistical Analysis

For the statistical analysis, the software R (version 4.1.0, R Foundation for Statistical Computing, Vienna, Austria) was used. An  $8 \times 4$  factorial arrangement was used with the independent variables being the days of fermentation (0, 2, 4, 6, 8, 10, 12 and 14) and the concentration of sodium chloride (0.05, 0.01, 0.20 and 0.40%).

## 3. Results and Discussions

The determination of  $L^*$ ,  $a^*$  and  $b^*$  values for flour samples fermented by *M. purpureus* using sodium chloride and monosodium glutamate during a 14-day incubation period are shown in Table 1. The values obtained for the lightness ( $L^*$ ) ranged from 45.95 to 64.24, showing that on days 8, 10, 12 and 14, the lowest values were obtained as  $45.95 \pm 2.334$ ,  $46.33 \pm 1.950$ ,  $46.13 \pm 2.819$  and  $46.97 \pm 2.335$ , respectively, the above results showed no significant differences. The lightness of the flour samples decreases due to the fermentative process of the fungus where the production of secondary metabolites such as pigments increases. This production of pigments, mainly red, also caused  $a^*$  values to vary between 6.52 and 19.79 belonging to the red colour range ( $+a^*$ ), where the highest values were obtained on days 6, 8 and 10 with a mean of  $19.33 \pm 1.026$ ,  $19.79 \pm 1.064$  and  $19.17 \pm 0.985$ , respectively, with no significant differences between them. These values are favoured by the nitrogen source (monosodium glutamate) which is related to a higher production of red pigments [9], solid state fermentation is also considered to improve the production of red pigments [4,10]. On the other hand, the values obtained for  $b^*$  were located in the yellow

colour zone (+b\*) between 18.95 and 27.78, with the lowest values on days 12 and 14, with a mean of  $18.75 \pm 0.770$  and  $18.95 \pm 0.999$ , respectively, showing no significant differences, where the initial yellow colour of the quinoa began to decrease due to the production of pigments in the *Monascus* fermentation.

**Table 1.** Colorimetric characteristics of *M. purpureus* pigments on different fermentation days.

Fermentation Time <sup>1</sup> (Day)	CIELAB Colour System <sup>2</sup>		
	L*	a*	b*
0	64.24 ± 1.457 <sup>e</sup>	6.52 ± 0.290 <sup>a</sup>	27.72 ± 0.205 <sup>e</sup>
2	61.34 ± 1.650 <sup>d</sup>	7.80 ± 0.408 <sup>b</sup>	27.78 ± 0.632 <sup>e</sup>
4	55.93 ± 2.200 <sup>c</sup>	14.25 ± 0.868 <sup>c</sup>	25.65 ± 0.984 <sup>d</sup>
6	49.14 ± 2.725 <sup>b</sup>	19.33 ± 1.026 <sup>ef</sup>	21.99 ± 1.442 <sup>c</sup>
8	45.95 ± 2.334 <sup>a</sup>	19.79 ± 1.064 <sup>f</sup>	20.28 ± 0.857 <sup>b</sup>
10	46.33 ± 1.950 <sup>a</sup>	19.17 ± 0.985 <sup>ef</sup>	19.99 ± 1.079 <sup>b</sup>
12	46.13 ± 2.819 <sup>a</sup>	18.50 ± 1.211 <sup>de</sup>	18.75 ± 0.770 <sup>a</sup>
14	46.97 ± 2.335 <sup>a</sup>	17.93 ± 0.865 <sup>d</sup>	18.95 ± 0.999 <sup>a</sup>

<sup>1</sup> Fermentation day with four levels of sodium chloride concentrations. <sup>2</sup> Mean ± standard deviation (SD). Black letters (a–f) represent statistically significant differences ( $p < 0.05$ ).

The values indicated did not show significant differences between the results of L\*, a\* and b\*; a\* being the most important within the results as it is related to the red colour. It is necessary to consider the fermentation time in the production of this sensory attribute, so day 8 was considered as the minimum time to stop the fermentation of the quinoa grains, obtaining a substrate with good colour.

The data obtained in the fermentation at day 8 with sodium chloride at different percentages were analysed and are shown in Table 2. The data for lightness (L\*) were found in a range from 44.09 to 48.48, the concentrations of 0.10, 0.20, 0.40% with a value of  $44.09 \pm 3.146$ ,  $45.70 \pm 0.890$  and  $45.53 \pm 1.615$  being the ones that presented a lower value without presenting significant differences. The results obtained for a\* were positive; that is to say that they are oriented to the red colour and showed no significant differences for any of the concentrations used, such as 0.05, 0.10, 0.20 and 0.40% of sodium chloride, obtaining  $19.85 \pm 1.174$ ,  $19.51 \pm 0.198$ ,  $20.47 \pm 0.289$  and  $19.35 \pm 1.780$ , respectively. The results obtained for b\* show that at 0.05, 0.20 and 0.40% sodium chloride were the lowest values with a mean of  $19.90 \pm 0.775$ ,  $19.56 \pm 0.349$  and  $20.39 \pm 0.638$ , respectively, with no significant differences.

**Table 2.** Colour characteristics of *M. purpureus* pigments at different sodium chloride concentrations on the eighth day of fermentation.

Sodium Chloride (%)	CIELAB Colour System <sup>1</sup>			C:N Ratio
	L*	a*	b*	
0.05	48.48 ± 0.713 <sup>b</sup>	19.85 ± 1.174 <sup>a</sup>	19.90 ± 0.775 <sup>a</sup>	11.31 ± 0.258 <sup>a</sup>
0.1	44.09 ± 3.146 <sup>a</sup>	19.51 ± 0.198 <sup>a</sup>	21.26 ± 0.592 <sup>b</sup>	11.95 ± 0.313 <sup>b</sup>
0.2	45.70 ± 0.890 <sup>ab</sup>	20.47 ± 0.289 <sup>a</sup>	19.56 ± 0.349 <sup>a</sup>	12.56 ± 0.199 <sup>c</sup>
0.4	45.53 ± 1.615 <sup>ab</sup>	19.35 ± 1.780 <sup>a</sup>	20.39 ± 0.638 <sup>ab</sup>	13.14 ± 0.248 <sup>d</sup>

<sup>1</sup> Mean ± standard deviation (SD). Black letters (a–d) represent statistically significant differences ( $p < 0.05$ ).

The values shown in Table 2 indicate that sodium chloride concentration did not affect the red colour (+a\*), but did affect lightness (L\*) and yellow colour (+b\*). The variations in L\* and b\* were the result of the effect of osmotic stress on the pigment production of *M. purpureus* [6]. Considering that the research is focused on the production of a red flour with fermentation supplements, the lowest concentration of sodium chloride (0.05%) was considered due to its cost-performance.

At a concentration of 0.05% sodium chloride on day 8 the lowest C:N ratio was produced with a mean value of  $11.31 \pm 0.258$ ; the C:N ratio is of great importance for pigment production considering that an amount higher than 20 could give a greater intensity to the red colouring, although the production of the dyes will also depend on the strain and other factors [11].

#### 4. Conclusions

This research showed that the pigmented flour produced by the solid state fermentation of quinoa by *M. purpureus* supplemented with monosodium glutamate and sodium chloride showed variation in the days of fermentation with respect to the colour red, with the eighth day being the appropriate time to stop fermentation, obtaining the values of  $L^*$  ( $48.48 \pm 0.713$ ),  $a^*$  ( $19.85 \pm 1.174$ ),  $b^*$  ( $19.90 \pm 0.775$ ) and C:N ( $11.31 \pm 0.258$ ) and resulting in a product with a good visual sensory attribute that can be used to develop new naturally pigmented products with possible functional characteristics.

**Author Contributions:** Conceptualization, U.G.-B. and V.C.; methodology, E.Q.-R., F.T.-H. and V.C.; software, E.Q.-R. and V.C.; validation, V.C.; formal analysis, E.Q.-R., U.G.-B. and V.C.; investigation, E.Q.-R. and F.T.-H.; resources, M.S.-J.; data curation, E.Q.-R. and V.C.; writing—original draft preparation, E.Q.-R.; writing—review and editing, V.C.; visualization, E.Q.-R., U.G.-B. and V.C.; supervision, M.S.-J., U.G.-B. and V.C.; project administration, M.S.-J.; funding acquisition, M.S.-J. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by CONCYTEC-PROCIENCIA under Basic Research Project 2019-01 [contract 383-2019- FONDECYT]. The Laboratorio de Microbiología de Alimentos UNALM, Laboratorio de Biotecnología Ambiental-Biorremediación UNALM and Centro de Investigación de Montaña (CIMO).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Summary data available upon request.

**Conflicts of Interest:** The authors declare no conflict of interest.

#### References

- Salim, R.; Fadel, M.; Youssef, Y.; Taie, H.; Abosereh, N.; El-Sayed, G.; Marzouk, M. A local *Talaromyces atroroseus* TRP-NRC isolate: Isolation, genetic improvement, and biotechnological approach combined with LC/HRESI-MS characterization, skin safety, and wool fabric dyeing ability of the produced red pigment mixture. *J. Genet. Eng. Biotechnol.* **2022**, *20*, 1–22. [\[CrossRef\]](#) [\[PubMed\]](#)
- Da Costa, J.; De Oliveira, C.; Vendruscolo, F. Cheese whey as a potential substrate for *Monascus* pigments production. *AIMS Agric. Food* **2020**, *5*, 785–798. [\[CrossRef\]](#)
- Gong, P.; Shi, R.; Liu, Y.; Luo, Q.; Wang, C.; Chen, W. Recent advances in *Monascus* pigments produced by *Monascus purpureus*: Biosynthesis, fermentation, function, and application. *LWT* **2023**, *185*, 115162. [\[CrossRef\]](#)
- Babitha, S.; Soccol, C.R.; Pandey, A. Jackfruit Seed—A novel substrate for the production of *Monascus* pigments through solid-state fermentation. *Food Technol. Biotechnol.* **2006**, *44*, 465–471.
- Mehri, D.; Perendeci, N.; Goksungur, Y. Utilization of whey for red pigment production by *Monascus purpureus* in submerged fermentation. *Fermentation* **2021**, *7*, 75. [\[CrossRef\]](#)
- Babitha, S.; Soccol, C.; Pandey, A. Effect of stress on growth, pigment production and morphology of *Monascus* sp. in solid cultures. *J. Basic Microbiol.* **2007**, *47*, 118–126. [\[CrossRef\]](#) [\[PubMed\]](#)
- Quispe-Rivera, E.; Tucta-Huillca, F.; Silva-Jaimes, M.; Gonzales-Barron, U.; Cadavez, V. Optimization of Pigment Extraction from Quinoa Flour Fermented by *Monascus purpureus* Supplemented with Sodium Chloride. *Biol. Life Sci. Forum* **2022**, *18*, 67. [\[CrossRef\]](#)
- Huang, R.; Huang, K.; Guan, X.; Li, S.; Cao, H.; Zhang, Y.; Lao, X.; Bao, Y.; Wang, J. Effect of defatting and extruding treatment on the physicochemical and storage properties of quinoa (*Chenopodium quinoa* Wild) flour. *LWT* **2021**, *147*, 111612. [\[CrossRef\]](#)
- Chatterjee, S.; Maity, S.; Chattopadhyay, P.; Sarkar, A.; Laskar, S.; Sen, S. Characterization of red pigment from *Monascus* in submerged culture red pigment from *Monascus purpureus*. *J. Appl. Sci. Res.* **2009**, *5*, 2102–2108.

10. De Oliveira, F.; Rocha, I.; Pinto, D.; Ventura, S.; Dos Santos, A.; Crevelin, E.; Ebinuma, V. Identification of azaphilone derivatives of *Monascus* colorants from *Talaromyces amestolkiae* and their halochromic properties. *Food Chem.* **2022**, *372*, 131214. [[CrossRef](#)] [[PubMed](#)]
11. Said, F.; Brooks, J.; Chisti, Y. Optimal C: N ratio for the production of red pigments by *Monascus ruber*. *World J. Microbiol. Biotechnol.* **2014**, *30*, 2471–2479. [[CrossRef](#)] [[PubMed](#)]

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.