



Proceeding Paper

# Divergent Impacts of Moderate and Severe Drought on the Antioxidant Response of *Calendula officinalis* L. Leaves and Flowers <sup>†</sup>

María Rita Guzman <sup>1,\*</sup> and Isabel Marques <sup>2,\*</sup>

<sup>1</sup> Estacion Biológica La Ceiba, Chisec 16013, Guatemala

<sup>2</sup> Forest Research Center (CEF) & Associate Laboratory TERRA, Instituto Superior de Agronomia (ISA), Universidade de Lisboa (UL), Tapada da Ajuda, 1349-017 Lisbon, Portugal

\* Correspondence: mrguzman@gmail.com (M.R.G.); isabelmarques@isa.ulisboa.pt (I.M.)

<sup>†</sup> Presented at the 3rd International Electronic Conference on Agronomy, 15–30 October 2023; Available online: <https://iecag2023.sciforum.net/>.

**Abstract:** We studied the impacts of moderate and severe drought on different pot marigold (*Calendula officinalis* L.) genotypes, evaluating the antioxidant performance of leaves and flowers concerning the levels of proline and malondialdehyde, the activity of antioxidant enzymes (catalase, peroxidase, and ascorbate peroxidase), as well as impacts on flower production. Overall, we found high resilience to moderate drought. However, the severe drought significantly affected flower production, despite the high level of antioxidants, proline, and malondialdehyde. Results also indicate significant variation in drought tolerance among pot marigolds, providing an opportunity to identify valuable tolerance traits.

**Keywords:** abiotic stress; antioxidants; drought; flower tolerance; marigold

## 1. Introduction

Drought stress is a major abiotic stress, limiting crop production and yield [1,2]. Drought decreases photosynthesis and chlorophyll synthesis, and alters nutrient metabolism, ion uptake, and translocation, ultimately limiting plant vegetative growth. Drought also has an impact on flower number and size, affecting the viability and durability of flowers [3]. The abortion of reproductive organs is a major limiting factor for flower production under water stress. Thus, understanding its impacts has become a major aim for the sustainability of the floricultural industry.

Drought triggers the overproduction of reactive oxygen species (ROS), which are toxic and cause damage to proteins, lipids, carbohydrates, and DNA, leading to oxidative stress and, ultimately, plant death [4,5]. To cope with the effects of drought, plants activate the antioxidant system that includes a variety of ROS scavengers, such as superoxide dismutase (SOD), ascorbate peroxidase (APX), peroxidase (POX), and catalase (CAT), as well as non-enzymatic metabolites such as carotenoids, flavonoids, and proline [6]. Malondialdehyde (MDA) produced by membrane lipids in response to ROS, is often used as a drought indicator to evaluate the degree of membrane damage and the level of drought tolerance since plants with low amounts of MDA are generally considered to be more tolerant to drought [7]. Nevertheless, the antioxidant response is highly dependent on the level and duration of drought and the species, and even varies within genotypes of the same species [8–10].

*Calendula officinalis* L. (pot marigold or calendula), from the plant family Asteraceae, is a widely used plant in ornamental horticulture, as well as in traditional healing treatments, due to its wide range of secondary metabolites, flavonoids, and carotenoids content [11]. It has recently been proposed as an oilseed crop since the oil from its seeds has high



**Citation:** Guzman, M.R.; Marques, I. Divergent Impacts of Moderate and Severe Drought on the Antioxidant Response of *Calendula officinalis* L. Leaves and Flowers. *Biol. Life Sci. Forum* **2023**, *27*, 53. <https://doi.org/10.3390/IECAG2023-16636>

Academic Editor: Gianni Bellocchi

Published: 15 December 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

amounts of fatty acids [11]. Nevertheless, it is highly affected by drought, although some genotypes show some tolerance to water scarcity [12]. For instance, drought caused a significant decline in the number of leaves and leaf area in *Calendula* cv. Orange King [13]. In measurements performed under field conditions, drought increased the activity of CAT, POX, and APX, as well as the levels of MDA and leaf and root proline while reducing seed yield in several pot marigold genotypes [12]. Nevertheless, high variation in responses was found between genotypes [12]. Drought can be a significant constraint in this species, but studies performed under controlled conditions remain scarce. Consequently, this study aims to understand the impacts of moderate and severe drought in four different pot marigold genotypes under a controlled environment. Specifically, we aim to understand the response of the pot marigold antioxidant machinery to drought, namely, the levels of proline and malondialdehyde, and the enzymatic activity of catalase (CAT), peroxidase (POX), and ascorbate peroxidase (APX) in the leaves and flowers. We also aim to detect the impact of drought on the production of flowers.

## 2. Material and Methods

### 2.1. Plant Experimental Design

The antioxidant performance of *Calendula officinalis* was tested in four different genotypes: cv. Indian Prince, Golden Emperor, Orange Prince, and Sun Glow. Seeds were grown in 2 L capacity pots in a controlled environmental chamber under a long-day photoperiod (16 h light), at a temperature of 23/1 °C (light/dark period) and relative humidity of 72–76%. Plants were watered every two days using the Hoagland nutritive solution. One-month-old *Calendula* plants were subjected to different water treatments (control, 100% field capacity (FC); moderate drought, 60% FC; and severe drought, 35% FC) and followed for three weeks [12]. Each treatment consisted of 10 biological replications per cultivar.

### 2.2. Non-Enzymatic Activities in Leaves and Flowers

Proline was determined according to acid–ninhydrin and toluene methods [14]. Absorbance was determined at 520 nm. Results are expressed in micrograms of proline per gram of dry weight. Lipid peroxidation was quantified according to [15] and measured in terms of malondialdehyde content (MDA). Absorbance was measured at 532 nm. Results are expressed as nmol of MDA per gram of dry weight.

### 2.3. Antioxidative Enzyme Activities in Leaves and Flowers

The activity of catalase (CAT) was determined in a 1.5 mL reaction mixture with 50 mM K-phosphate buffer (pH 7.0), 10 mM H<sub>2</sub>O<sub>2</sub>, and the enzyme following [16], and measured at 240 nm. The results were expressed in CAT mg<sup>-1</sup> of protein. Peroxidase activity (POX) was determined as described by [17], measuring the absorbance at 430 nm. Results are expressed in units mg<sup>-1</sup> of protein. Ascorbate peroxidase (APX) was determined according to [18], measured at 290 nm. Results are expressed in APX mg<sup>-1</sup> of protein. Data are expressed in dry weight (DW).

### 2.4. Impacts of Drought on the Production of Flowers

For each genotype and experimental treatment, we quantified the total number of flowers produced. The dry matter content was acquired by drying samples to a constant weight using a thermo-ventilated oven at 65 °C.


### 2.5. Statistical Analysis

Mean values ( $\pm$ SE) were calculated from 10 replicates per cultivar using IBM SPSS v.22. To analyze the effects of salinity, we used a multivariate ANOVA or a *t*-test after checking the homogeneity of variance using Levene's Test for Equality of Variances. Significant differences between means were also subjected to Tukey's test for post hoc comparisons (at the 1% significance level).

### 3. Results and Discussion

#### 3.1. Non-Enzymatic Activities in Leaves and Flowers

Moderate drought had no significant effects on the levels of proline in the leaves of the *Calendula* genotypes ( $F_{2,11} = 0.341$ ,  $p = 0.871$ ), while some slight significant increases were already reported in the flowers of Golden Emperor and Sun Glow (respectively,  $F_{2,11} = 11.872$ ,  $p < 0.05$ ;  $F_{2,11} = 10.201$ ,  $p < 0.05$ ; Figure 1A). In contrast, severe drought significantly increased the levels of proline in all genotypes, either considering leaves or flowers (Figure 1A). The accumulation of proline under drought stress has been reported in many plant species [19], including *Calendula* [12]. Proline is mainly synthesized in the leaves and transported to other areas to balance osmotic pressure and scavenge ROS, allowing plants to cope with drought [20]. Under stress, proline also maintains cell turgor or osmotic balance, stabilizes membranes, and prevents oxidative bursts in plants. Thus, the increase found in this study suggests its role in protecting *Calendula* cultivars from the effects of severe drought.



**Figure 1.** Effects of moderate drought (MD) and severe drought (SD) related to control conditions recorded in the leaves and flowers of four *Calendula officinalis* genotypes. (A) Increases in proline content related to control conditions ( $\mu\text{g g DW}^{-1}$ ). (B) Increases in malondialdehyde levels related to control conditions (MDA;  $\text{nmol g DW}^{-1}$ ). Mean values  $\pm$  SE ( $n = 10$ ). Asterisks indicate significant differences between MD or SD and control conditions for the same species ( $t$ -test at  $p < 0.001$ ), considering only leaves or flowers.

The levels of MDA showed no significant differences between the control conditions and moderate drought when recorded in the leaves ( $F_{2,11} = 0.121$ ,  $p < 0.05$ ; Figure 1B). However, slight increases were already felt in the flowers of the Golden Emperor and Sun Glow genotypes (Figure 1B). The highest increase in MDA was recorded under severe drought, in all genotypes, especially in the Golden Emperor and Sun Glow genotypes (Figure 1B). ROS induces lipid peroxidation, giving rise to MDA, an indicator of membrane damage, especially during stress. Overall, the more the plant is stressed, the higher its MDA content. Thus, this stress marker indicates that the flowers of some *Calendula* genotypes were already affected by moderate drought, while severe drought had strong negative effects on lipid peroxidation in the leaves and flowers of all genotypes. The level of MDA can be used in future studies to evaluate the degree of plasma membrane damage and the ability of *Calendula* plants to tolerate drought stress.

#### 3.2. Antioxidative Enzyme Activities in Leaves and Flowers

Under moderate drought, the enzymatic activities of catalase (CAT), peroxidase (POX), and ascorbate peroxidase (APX) measured in the leaves showed no significant differences from control conditions in all genotypes (Table 1). Nevertheless, the flowers of some genotypes already showed a significant increase in the activity of CAT (Golden Emperor, Orange Prince, and Sun Glow) and POX (Indian Prince) under moderate conditions (Table 1).

**Table 1.** Enzyme activities of catalase (CAT), peroxidase (POX), and ascorbate peroxidase (APX), expressed as mg of protein/DW under control water conditions, moderate drought (MD), and severe drought (SD) in four *Calendula officinalis* genotypes. Results indicate measurements in leaves/flowers. Results are expressed as means  $\pm$  SE (n = 10). Different superscripts indicate significant differences between water levels for the same species (ANOVA followed by a Tukey test at  $p < 0.001$ ) considering only leaves or flowers.

	Control	MD	SD
<b>CAT</b>			
Indian Prince	11.23 $\pm$ 2.21 <sup>a</sup> /8.14 $\pm$ 1.17 <sup>a</sup>	11.66 $\pm$ 3.44 <sup>a</sup> /8.11 $\pm$ 1.19 <sup>a</sup>	28.37 $\pm$ 4.22 <sup>b</sup> /10.14 $\pm$ 1.29 <sup>a</sup>
Golden Emperor	14.25 $\pm$ 2.74 <sup>a</sup> /7.89 $\pm$ 1.15 <sup>a</sup>	14.23 $\pm$ 3.01 <sup>a</sup> /8.55 $\pm$ 2.20 <sup>b</sup>	21.99 $\pm$ 4.05 <sup>b</sup> /10.31 $\pm$ 2.17 <sup>c</sup>
Orange Prince	12.07 $\pm$ 2.22 <sup>a</sup> /7.99 $\pm$ 1.11 <sup>a</sup>	12.03 $\pm$ 2.66 <sup>a</sup> /9.01 $\pm$ 1.25 <sup>b</sup>	16.28 $\pm$ 3.31 <sup>b</sup> /14.20 $\pm$ 1.99 <sup>c</sup>
Sun Glow	10.03 $\pm$ 2.00 <sup>a</sup> /6.45 $\pm$ 1.11 <sup>a</sup>	10.20 $\pm$ 2.11 <sup>a</sup> /7.25 $\pm$ 1.19 <sup>b</sup>	17.25 $\pm$ 3.07 <sup>b</sup> /9.29 $\pm$ 2.33 <sup>c</sup>
<b>POX</b>			
Indian Prince	1.24 $\pm$ 0.22 <sup>a</sup> /0.33 $\pm$ 0.07 <sup>a</sup>	1.51 $\pm$ 0.44 <sup>a</sup> /1.03 $\pm$ 0.98 <sup>b</sup>	3.25 $\pm$ 0.22 <sup>b</sup> /2.07 $\pm$ 0.01 <sup>c</sup>
Golden Emperor	0.98 $\pm$ 0.21 <sup>a</sup> /0.41 $\pm$ 0.11 <sup>a</sup>	1.03 $\pm$ 0.11 <sup>a</sup> /0.48 $\pm$ 0.67 <sup>a</sup>	2.71 $\pm$ 0.56 <sup>b</sup> /1.06 $\pm$ 0.23 <sup>b</sup>
Orange Prince	1.08 $\pm$ 0.22 <sup>a</sup> /0.37 $\pm$ 0.13 <sup>a</sup>	1.13 $\pm$ 0.66 <sup>a</sup> /0.41 $\pm$ 0.21 <sup>a</sup>	2.08 $\pm$ 0.71 <sup>b</sup> /1.03 $\pm$ 0.01 <sup>b</sup>
Sun Glow	1.12 $\pm$ 0.19 <sup>a</sup> /0.56 $\pm$ 0.212 <sup>a</sup>	1.18 $\pm$ 0.18 <sup>a</sup> /0.23 $\pm$ 2.21 <sup>a</sup>	2.01 $\pm$ 0.98 <sup>b</sup> /1.20 $\pm$ 0.21 <sup>b</sup>
<b>APX</b>			
Indian Prince	9.44 $\pm$ 2.01 <sup>a</sup> /4.55 $\pm$ 1.04 <sup>a</sup>	9.51 $\pm$ 2.03 <sup>a</sup> /4.23 $\pm$ 2.01 <sup>a</sup>	18.23 $\pm$ 3.05 <sup>b</sup> /5.20 $\pm$ 2.24 <sup>b</sup>
Golden Emperor	10.21 $\pm$ 2.33 <sup>a</sup> /3.03 $\pm$ 1.89 <sup>a</sup>	10.25 $\pm$ 2.26 <sup>a</sup> /2.99 $\pm$ 2.28 <sup>a</sup>	16.39 $\pm$ 2.56 <sup>b</sup> /4.56 $\pm$ 1.98 <sup>b</sup>
Orange Prince	11.23 $\pm$ 2.04 <sup>a</sup> /6.21 $\pm$ 2.23 <sup>a</sup>	10.99 $\pm$ 3.01 <sup>a</sup> /6.15 $\pm$ 1.99 <sup>a</sup>	14.99 $\pm$ 2.27 <sup>b</sup> /8.05 $\pm$ 2.01 <sup>b</sup>
Sun Glow	8.05 $\pm$ 2.31 <sup>a</sup> /4.02 $\pm$ 1.89 <sup>a</sup>	8.12 $\pm$ 2.01 <sup>a</sup> /4.11 $\pm$ 2.17 <sup>a</sup>	11.13 $\pm$ 2.27 <sup>b</sup> /6.73 $\pm$ 2.35 <sup>b</sup>

The activities of all enzymes increased significantly under severe drought in all genotypes and considering both leaves and flowers (Table 1). However, significant variation in enzyme activities was found between genotypes (always  $p > 0.05$ ). The highest enzymatic activities in the leaves were reported in the genotype Indian Prince under severe drought, while the highest enzymatic values were recorded in the flowers of the genotype Orange Prince, also under severe drought (Table 1). An increase in the activity of antioxidant enzymes was also found in other *Calendula* genotypes subjected to drought, although high variation was found between genotypes [12]. Together with the increase in the content of proline, the higher antioxidant enzyme activities found under drought suggest a good antioxidant mechanism to cope with drought. Altogether, these enzymatic and non-enzymatic components help to reduce the oxidative stress in *Calendula* triggered by drought, as reported in other plants [18,21]. *Calendula* plants can keep ROS under control through this efficient and versatile scavenging system. This would help to protect cellular structures and functions as well as to maintain water balance and the efficiency of physiological processes.

### 3.3. Impacts of Drought on the Production of Flowers

Although flowers are crucial for the floriculture industry, studies on understanding the impacts of abiotic stresses, especially concerning *Calendula* species, are remarkably scarce. In this study, flower production decreased by 5.01% in Indian Prince, 4.39% in Golden Emperor, 6.99% in Orange Prince, and 6.81% in Sun Glow under moderate drought. However, a harsh effect was felt under severe drought, decreasing the production of flowers by 23.67% in Indian Prince, 25.66% in Golden Emperor, 37.53% in Orange Prince, and 39.88% in Sun Glow. In general, drought decreases flower production in many species [22] since flower development and the related reproductive processes are very sensitive to stress, and also because resources are allocated to plant survival under stress conditions. Therefore, it is not surprising to find that drought also has a strong impact on *Calendula* plants. However, as impacts vary between genotypes, it is crucial to conduct further tests on drought tolerance by screening additional *Calendula* plants. Understanding the effects of stress on flower development or abortion would help to develop high-yield cultivars that can cope with environmental changes using traditional and molecular breeding approaches.

#### 4. Conclusions

*Calendula* plants showed high resilience to moderate drought, in contrast with severe drought, which had a harsh impact on most genotypes. The levels of proline and MDA can be used in future studies as stress markers to understand the impacts of drought on these plants. The antioxidant machinery studied here increased under the harsh drought effect but did not prevent negative effects on flower production, which was significantly affected by drought. As drought showed a negative impact on flower production, future studies should focus on understanding its effects on flower development and fertility. Additionally, high variation was found between pot marigold genotypes, suggesting differences in drought tolerance, which can be used to screen useful tolerance traits. Apart from adjustments in the antioxidant system, it should be noted that drought tolerance depends on additional plant features, which should be measured to characterize the severity of drought. Useful traits for future studies should include, for example, net photosynthesis, the abundance of osmoprotectants, ABA content, and membrane integrity. It would also be useful to identify genes that respond to abiotic stresses in *Calendula* plants.

**Author Contributions:** Conceptualization, I.M.; methodology, I.M.; formal analysis, M.R.G.; investigation, M.R.G.; data curation, M.R.G.; writing—original draft preparation, M.R.G. and I.M.; writing—review and editing, M.R.G. and I.M.; supervision, I.M.; project administration, I.M.; funding acquisition, I.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received national funds through the FCT—Fundação para a Ciência e a Tecnologia, I.P., Portugal, through the research unit UIDB/00239/2020 (CEF), and under the Scientific Employment Stimulus—Individual Call (CEEC Individual)—2021.01107.CEECIND/CP1689/CT0001 (IM).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** The original contributions presented in the study are included in the article, further inquiries can be directed to the corresponding author.

**Conflicts of Interest:** The authors declare no conflicts of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of the data; in the writing of the manuscript; or in the decision to publish the results.

#### References

1. Fernandes, I.; Marques, I.; Paulo, O.S.; Batista, D.; Partelli, F.L.; Lidon, F.C.; Damatta, F.M.; Ramalho, J.C.; Ribeiro-Barros, A.I. Understanding the impact of drought in *Coffea* genotypes: Transcriptomic analysis supports a common high resilience to moderate water deficit but a genotype dependent sensitivity to severe water deficit. *Agronomy* **2021**, *11*, 2255. [[CrossRef](#)]
2. Fahad, S.; Bajwa, A.A.; Nazir, U.; Anjum, S.A.; Farooq, A.; Zohaib, A.; Sadia, S.; Nasim, W.; Adkins, S.; Saud, S.; et al. Crop production under drought and heat stress: Plant responses and management options. *Front. Plant Sci.* **2017**, *8*, 1147. [[CrossRef](#)] [[PubMed](#)]
3. Kuppler, J.; Wieland, J.; Junker, R.R.; Ayasse, M. Drought-induced reduction in flower size and abundance correlates with reduced flower visits by bumble bees. *AoB Plants* **2021**, *13*, plab001. [[CrossRef](#)] [[PubMed](#)]
4. Chaves, M.M.; Oliveira, M.M. Mechanisms underlying plant resilience to water deficits: Prospects for water-saving agriculture. *J. Exp. Bot.* **2004**, *55*, 2365–2384. [[CrossRef](#)] [[PubMed](#)]
5. Lawas, L.M.F.; Shi, W.; Yoshimoto, M.; Hasegawa, T.; Hinch, D.K.; Zuther, E.; Jagadish, S.V.K. Combined drought and heat stress impact during flowering and grain filling in contrasting rice cultivars grown under field conditions. *Field Crop. Res.* **2018**, *229*, 66–77. [[CrossRef](#)]
6. Gill, S.S.; Tuteja, N. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol. Biochem.* **2010**, *48*, 909–930. [[CrossRef](#)] [[PubMed](#)]
7. Ma, J.; Du, G.; Li, X.; Zhang, C.; Guo, J. A major locus controlling malondialdehyde content under water stress is associated with Fusarium crown rot resistance in wheat. *Mol. Genet. Genom.* **2015**, *290*, 1955–1962. [[CrossRef](#)] [[PubMed](#)]
8. Hasanuzzaman, M.; Raihan, M.R.H.; Masud, A.A.C.; Rahman, K.; Nowroz, F.; Rahman, M.; Nahar, K.; Fujita, M. Regulation of reactive oxygen species and antioxidant defense in plants under salinity. *Int. J. Mol. Sci.* **2021**, *22*, 9326. [[CrossRef](#)]
9. Vinci, G.; Marques, I.; Rodrigues, A.P.; Martins, S.; Leitão, A.E.; Smedo, M.C.; Silva, M.J.; Lidon, F.C.; DaMatta, F.M.; Ribeiro-Barros, A.I.; et al. Protective Responses at the Biochemical and Molecular Level Differ between a *Coffea arabica* L. Hybrid and Its Parental Genotypes to Supra-Optimal Temperatures and Elevated Air [CO<sub>2</sub>]. *Plants* **2022**, *11*, 2702. [[CrossRef](#)]

10. Marques, I.; Fernandes, I.; Paulo, O.S.; Batista, D.; Lidon, F.C.; Partelli, F.; DaMatta, F.M.; Ribeiro-Barros, A.I.; Ramalho, J.C. Overexpression of Water-Responsive Genes Promoted by Elevated CO<sub>2</sub> Reduces ROS and Enhances Drought Tolerance in Coffea Species. *Int. J. Mol. Sci.* **2023**, *24*, 3210. [[CrossRef](#)]
11. Ahmad, S.; Ahmed, W.; Mihoub, A.; Jamal, A.; Farhan Saeed, M.; Masood, N.; Radicetti, E.; Fawad, M.; Nicola, S. Developmental, Phytochemical and Enzymatic Changes in Pot Marigold (*Calendula officinalis* L.) cvs. Hybrid and French with Salicylic Acid (SA) and Polyamine Spermidine (SP) Foliar Spray. *Agronomy* **2023**, *13*, 191. [[CrossRef](#)]
12. Ghadyeh Zarrinabadi, I.; Razmjoo, J.; Abdali Mashhadi, A.; Karim mojeni, H.; Boroomand, A. Physiological response and productivity of pot marigold (*Calendula officinalis*) genotypes under water deficit. *Ind. Crops Prod.* **2019**, *139*, 111488. [[CrossRef](#)]
13. Akhtar, G.; Faried, H.N.; Razzaq, K.; Ullah, S.; Wattoo, F.M.; Shehzad, M.A.; Sajjad, Y.; Ahsan, M.; Javed, T.; Dessoky, E.S.; et al. Chitosan-Induced Physiological and Biochemical Regulations Confer Drought Tolerance in Pot Marigold (*Calendula officinalis* L.). *Agronomy* **2022**, *12*, 474. [[CrossRef](#)]
14. Khedr, A.H.A.; Abbas, M.A.; Abdel Wahid, A.A.; Quick, W.P.; Abogadallah, G.M. Proline induces the expression of salt-stress-responsive proteins and may improve the adaptation of *Pancreaticum maritimum* L. to salt-stress. *J. Exp. Bot.* **2003**, *54*, 2553–2562. [[CrossRef](#)] [[PubMed](#)]
15. De Azevedo Neto, A.D.; Prisco, J.T.; Enéas-Filho, J.; De Abreu, C.E.B.; Gomes-Filho, E. Effect of salt stress on antioxidative enzymes and lipid peroxidation in leaves and roots of salt-tolerant and salt-sensitive maize genotypes. *Environ. Exp. Bot.* **2006**, *56*, 87–94. [[CrossRef](#)]
16. Chrysargyris, A.; Tzionis, A.; Xylia, P.; Tzortzakis, N. Effects of salinity on tagetes growth, physiology, and shelf life of edible flowers stored in passive modified atmosphere packaging or treated with ethanol. *Front. Plant Sci.* **2018**, *871*, 1765. [[CrossRef](#)] [[PubMed](#)]
17. Tarchoune, I.; Sgherri, C.; Izzo, R.; Lachaal, M.; Ouergui, Z.; Navari-Izzo, F. Antioxidative responses of *Ocimum basilicum* to sodium chloride or sodium sulphate salinization. *Plant Physiol. Biochem. PPB* **2010**, *48*, 772–777. [[CrossRef](#)]
18. Zhu, Z.; Wei, G.; Li, J.; Qian, Q.; Yu, J. Silicon alleviates salt stress and increases antioxidant enzymes activity in leaves of salt-stressed cucumber (*Cucumis sativus* L.). *Plant Sci.* **2004**, *167*, 527–533. [[CrossRef](#)]
19. Elsayed, A.I.; El-Hamahmy, M.A.M.; Rafudeen, M.S.; Mohamed, A.H.; Omar, A.A. The impact of drought stress on antioxidant responses and accumulation of flavonolignans in milk thistle (*Silybum marianum* (L.) Gaertn.). *Plants* **2019**, *8*, 611. [[CrossRef](#)]
20. Cao, Y.; Luo, Q.; Tian, Y.; Meng, F. Physiological and proteomic analyses of the drought stress response in *Amygdalus Mira* (Koehne) Yü et Lu roots. *BMC Plant Biol.* **2017**, *17*, 53. [[CrossRef](#)]
21. Dumanović, J.; Nepovimova, E.; Natić, M.; Kuča, K.; Jačević, V. The Significance of Reactive Oxygen Species and Antioxidant Defense System in Plants: A Concise Overview. *Front. Plant Sci.* **2021**, *11*, 552969. [[CrossRef](#)] [[PubMed](#)]
22. Descamps, C.; Boubnan, N.; Jacquemart, A.L.; Quinet, M. Growing and Flowering in a Changing Climate: Effects of Higher Temperatures and Drought Stress on the Bee-Pollinated Species *Impatiens glandulifera* Royle. *Plants* **2021**, *10*, 988. [[CrossRef](#)] [[PubMed](#)]

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.