Results of Oilseed Poppy Breeding and Production Potential of New Varieties [†]

Jozef Fejér ^{1,*} , Ľuboš Nastišin ² and Janka Majdanová ³

¹ Department of Ecology (DE), Faculty of Humanities and Natural Sciences (FHNS), University of Prešov (UP), 17. November St., SK-080 01 Prešov, Slovakia

² Research and Breeding Station Malý Šariš, Research Institute of Plant Production (RIPP) Piešťany, National Agricultural and Food Centre (NAFC) Nitra, Malý Šariš 221, SK-080 01 Prešov, Slovakia; lubos.nastisin@nppc.sk

³ Testing Station Beluša, Department of Variety Testing, The Central Control and Testing Institute in Agriculture, Továrenská 421/2, SK-018 61 Beluša, Slovakia; janka.majdanova@uksup.sk

* Correspondence: jozef.fejer@unipo.sk

[†] Presented at the 1st International Electronic Conference on Agronomy, 3–17 May 2021; Available online: <https://iecag2021.sciforum.net/>.

Abstract: Breeding of poppy (*Papaver somniferum* L.) for food use of poppy seeds has a long tradition in Slovakia. Conventional breeding methods have brought several successes in the form of registration of new varieties with better yield parameters. In the breeding process, the methods of crossing and selection of offspring were used. In the years 2016–2018, the testing of selected poppy seeds was performed at the Central Control and Testing Institute in Agriculture with the aim to register new varieties. Testing was carried out according to the valid methodology in five localities of the Slovak Republic. The experiments were based on the block method on plots with an area of 10 m² in four replications. The obtained poppy seed yield and empty dry capsules were compared to the parameters of standard—control varieties ('Opal' and 'Major'). In the year 2019, four new food poppy varieties—'Azurit', 'MS Diamant', 'MS Topas', and 'MS Zafir'—were registered. On average, the obtained poppy seed yield during three years from five experimental localities ranged from 1.77 t·ha⁻¹ ± 0.16 ('Azurit') to 1.90 t·ha⁻¹ ± 0.17 ('MS Diamant'). For comparison, the average yield of control varieties was 1.76 t·ha⁻¹ ± 0.12. An important additional indication in evaluation of the variety economic value is the yield of empty dry capsules. This ranged from 0.98 t·ha⁻¹ ± 0.12 and 0.98 t·ha⁻¹ ± 0.09 ('Azurit' and 'MS Diamant', respectively) to 1.09 t·ha⁻¹ ± 0.21 and 1.09 t·ha⁻¹ ± 0.16 ('MS Topas' and 'MS Zafir', respectively). The average dry capsule yield of control varieties was 0.93 t·ha⁻¹ ± 0.13. Statistical evaluation by analysis of variance did not reveal significant differences in seed yield between the tested and control varieties ($p = 0.661$). The difference was significant in the yield of dry capsules ($p = 0.048$). The locality had a statistically significant effect on the yield of seeds and capsules ($p < 0.001$). A statistically significant difference on the yield of seeds ($p = 0.050$) and capsules ($p < 0.001$) was also found in the experimental year. New poppy varieties showed a generally higher seed yield potential compared to standard varieties. Based on the knowledge on soil and climatic conditions of individual experimental localities and test years, it is possible to choose a suitable variety for specific growing conditions in order to be able to maximize its genetic potential.

Keywords: *Papaver somniferum* L.; poppy seeds; poppy capsules; variety; yield

Citation: Fejér, J.; Nastišin, Ľ.; Majdanová, J. Results of Oilseed Poppy Breeding and Production Potential of New Varieties. *Biol. Life Sci. Forum* **2021**, *3*, 14. <https://doi.org/10.3390/IECAG2021-09681>

Academic Editor: Peter Langridge

Published: 1 May 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Poppy (*Papaver somniferum* L.) is a traditional crop of Slavic people. The consumption of poppy seed is deeply rooted in the culture of the people of Central Europe, including Slovakia. Poppy breeding in Slovakia has undergone a long-term development. Since the beginning of this creative activity (dating from 1948), the main goal in the creation of new

varieties has been focused on varieties of the universal type, i.e., varieties with a high yield of high-quality poppy seeds for food use and poppy straw with a high morphine content for the pharmaceutical industry. With regard to the negative correlations between seed yield and morphine content, as well as the preference for seed yield as the main product over the by-product poppy straw, varieties of this type were not registered in Slovakia until 2019. With regard to the interest in varieties producing quality poppy seeds and the interest in the quality of poppy straw by the pharmaceutical industry, breeding is currently divided into two directions—food-type varieties and industrial-type varieties.

This study provides an overview of the results in the breeding of poppy food varieties over the last ten years (although this activity has been ongoing since 1961 in Slovakia) at the Research and Breeding Station (Malý Šariš, RIPP, NAFC), and the results of cooperation between this institution and the Prešov University in Prešov.

2. Material and Methods

2.1. Breeding Methods

Methods of collecting landraces poppy populations, and of crossing domestic and foreign varieties and genotypes were used in the breeding of new varieties [1–4]. Subsequently, selection and evaluation of lines was performed in accordance with breeding objectives [5,6].

2.2. Plant Material

The evaluated varieties MS Topas, MS Zafir, MS Diamant and control varieties Opal and Major were bred on Research and Breeding Station Malý Šariš (NAFC Nitra, RIPP, Piešťany, Slovakia). This institution is also the owner/maintainer of the varieties. The breeder and the owner/maintainer of the Azurit variety is the University of Prešov in Prešov, Slovakia.

2.3. Establishment, Conduct and Evaluation of Field Experiments

Field experiments with poppy were carried out on plots with an area of 10 m² in four repetitions, at five localities in Slovakia during the years 2016–2018. The establishment and evaluation were carried out according to the methodology of CCTIA [7] and TG 166/4 [8]. Four new varieties were evaluated and compared to the standard varieties Major and Opal (year of registration, 2002 and 1995, resp.). These varieties were classified as standard comparison and stable varieties by The Central Control and Testing Institute in Agriculture, Slovakia.

3. Results

On average, the obtained poppy seed yield during three years from five experimental localities ranged from $1.77 \pm 0.16 \text{ t} \cdot \text{ha}^{-1}$ (Azurit) to $1.90 \pm 0.17 \text{ t} \cdot \text{ha}^{-1}$ (MS Diamant). For comparison, the average yield of control varieties was $1.76 \pm 0.12 \text{ t} \cdot \text{ha}^{-1}$ (Table 1). The statistically significant differences in seed yield ($p = 0.661$) between tested and control varieties were not found. The yield of dry empty capsules is presented in Table 2. It ranged from $0.98 \pm 0.12 \text{ t} \cdot \text{ha}^{-1}$ and $0.98 \pm 0.09 \text{ t} \cdot \text{ha}^{-1}$ ('Azurit' and 'MS Diamant', respectively) to $1.09 \pm 0.21 \text{ t} \cdot \text{ha}^{-1}$ and $1.09 \pm 0.16 \text{ t} \cdot \text{ha}^{-1}$ ('MS Topas' and 'MS Zafir', respectively). The average dry capsule yield of control varieties was $0.93 \pm 0.13 \text{ t} \cdot \text{ha}^{-1}$. Statistically significant differences in capsule yield were found ($p = 0.048$). In the past, poppy straw from such varieties has been used to extract alkaloids. Nowadays, the poppy straw from such type of varieties is unsuitable for pharmaceutical processing due to its low alkaloid content.

Table 1. Seed yield ($\text{t}\cdot\text{ha}^{-1}$) average values of the five experimental localities.

Variety	Year			Average	S.D.	
	2016	2017	2018			
Azurit	1.85	1.58	1.88	1.77	±	0.68
MS Topas	2.01	1.71	*	1.86	±	0.71
MS Zafir	1.99	1.72	*	1.85	±	0.75
MS Diamant	*	1.78	2.02	1.90	±	0.61
Major Control 1	1.84	1.68	1.91	1.81	±	0.71
Opal Control 2	1.82	1.61	1.70	1.71	±	0.56
Average C1 and C2	1.83	1.65	1.81	1.76	±	0.10

* = not evaluated; S.D. = standard deviation; C1 = Major, C2 = Opal.

Table 2. Dry empty capsules yield ($\text{t}\cdot\text{ha}^{-1}$) average values of the five experimental localities.

Variety	Year			Average	S.D.	
	2016	2017	2018			
Azurit	1.11	0.87	0.95	0.98	±	0.33
MS Topas	1.23	0.94	*	1.09	±	0.29
MS Zafir	1.21	0.98	*	1.09	±	0.35
MS Diamant	*	0.92	1.04	0.98	±	0.26
Major Control 1	1.14	0.93	1.00	1.02	±	0.29
Opal Control 2	0.95	0.77	0.81	0.84	±	0.18
Average C1 and C2	1.05	0.85	0.91	0.93	±	0.10

* = not evaluated; S.D. = standard deviation; C1 = Major, C2 = Opal.

A statistically significant influence of the experimental localities on the yield of seeds and capsules was found ($p < 0.001$). Furthermore, the statistical evidence of the experimental year on these parameters was found (yield of seeds $p = 0.050$; yield of dry capsules $p < 0.001$).

4. Conclusions

Newly registered food-type varieties demonstrated a higher poppy seed yield compared to standard control varieties. The significant influence of the experimental locality and the experimental year on the evaluated parameters was observed. Based on the known climatic and soil conditions of the experimental localities, it is possible to elaborate a recommendation for the zoning of suitable varieties to specific growing conditions.

Supplementary Materials: The poster presentation is available online at <https://www.mdpi.com/article/10.3390/IECAG2021-09681/s1>.

Author Contributions: Conceptualization, J.F., L.N. and J.M.; methodology J.F., L.N. and J.M.; formal analysis, J.F., L.N. and J.M.; writing—original draft preparation J.F., and L.N.; writing—review and editing, J.F. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: No additional data available.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Boháč, J. *Špeciálne šľachtenie poľnohospodárskych plodín [Special Breeding of Agricultural Crops]*; VŠP [Slovak University of Agriculture]: Nitra, Slovakia, 1986; pp. 171–177.
2. Fejer, J. Breeding Aspects of Slovakian Poppy Plants. In *Acta Horticulturae*; ISHS Secretariat: Leuven, Belgium, 2014; No. 1036; pp. 101–105. ISBN 978-94-62610-23-1.
3. Vašák, J. (Ed.) Mák [Poppy]. In *Šlechtění a odrůdy [Breeding and Varieties]*; Český mák a ČZÚ v Praze [Czech University of Life Sciences Prague]: Praha, Czech Republic, 2010; pp. 65–89.
4. Voškeruša, J.; Kratochvíl, V.; Martinek, V.; Mráz, F.; Němcová, D.; Žídková, A.; Derco, M.; Šinský, T.; Horel, F. *Pěstování olejnin v ČSSR. [Growing oilseeds in the Czechoslovak Socialist Republic]*; Státní zemědělské nakladatelství Praha [State Agricultural Publishing House Prague: Prague, Czechoslovak Socialist Republic, 1965; pp. 178–227.
5. Fejer, J. *Morfologicko—Biologická diverzita druhu mak siaty (Papaver somniferum L.) a jej hodnotenie [Morphological and Biological Diversity of Poppy Species and Its Evaluation]*; Prešovská univerzita v Prešove [University of Prešov in Prešov], Grafotlač Prešov: Prešov, Slovakia, 2015; 117p, ISBN 978-80-555-1334-8.
6. Fejer, J.; Brezinová, B.; Havrlentová, M. *Variabilita krajových populácií maku siateho (Papaver somniferum L.) a potenciál ich využitia pre rozšírenie genetickej diverzity pri tvorbe nových odrôd [Variability of Landraces Populations of Poppy (Papaver somniferum L.) and the Potential of Their Use for the Enlargement of Genetic Diversity in the Creation of New Varieties]*, 1. vyd. [1st ed.]; Vydavateľstvo Prešovskej Univerzity [University of Prešov Publishing]: Prešov, Slovakia, 2018; 102p, ISBN 978-80-555-2032-2.
7. *Methodology of CCTIA for Tests of Economic Value of Poppy*; Central Control and Testing Institute in Agriculture in Bratislava, Department of Variety Testing: Bratislava, Slovakia; 22p, internal document, unpublished.
8. TG/166/4. *Guidelines for the Conduct of Tests (GCT) for Distinctness, Uniformity and Stability (DUS)*; Opium/Seed Poppy (Papaver somniferum L.): Geneva, Switzerland, 2014; 31p.