



Abstract

Silicon Differentially Modulates Phenolic Metabolism and Antioxidant Capacity of Barley Cultivars Subjected to Phosphorus Deficiency[†]

Paula Cartes^{1,2}, Isis Vega¹ and Sofia Pontigo^{1,2,*} 

- ¹ Center of Plant, Soil Interaction and Natural Resources Biotechnology, Scientific and Technological Bioresource Nucleus (BIOREN-UFRO) Universidad de La Frontera, P.O. Box 54-D, Temuco 4780000, Chile; paula.cartes@ufroterra.cl (P.C.); i.vega01@ufroterra.cl (I.V.)
- ² Departamento de Ciencias Químicas y Recursos Naturales, Facultad de Ingeniería y Ciencias Universidad de La Frontera, P.O. Box 54-D, Temuco 4780000, Chile
- * Correspondence: sofia.pontigo@ufroterra.cl
- † Presented at the 1st International Electronic Conference on Agronomy, 3–17 May 2021. Available online: <https://iecag2021.sciforum.net/>.

Abstract: Current evidence shows that silicon (Si) can alleviate multiple plant stresses by inducing the antioxidant defense and phenolic metabolism of plants. Nevertheless, the mechanisms underlying these responses remain unclear. We investigated the Si effect on the phenolic metabolism of two barley cultivars differing in their tolerance to phosphorus (P) deficiency (cv. Sebastian, P-deficiency-tolerant and cv. Traveler, P-deficiency-sensitive). Plants were hydroponically grown with P (0, 0.01 or 0.1 mM P; applied as Na₂HPO₄) in combination with Si (0, 1 or 2 mM Si; applied as Na₂SiO₃). At harvest, total phenols, antioxidant capacity, individual phenolics, and the gene expression of enzymes involved in the synthesis of soluble phenolic compounds (including phenylalanine ammonia lyase (*HvPAL*) and chalcone synthase (*HvCHS*)) were analyzed in shoots. In cv. Sebastian grown without P, Si reduced both total phenols and antioxidant capacity to levels comparable to plants supplied with an optimal P dose. In contrast, increasing Si doses triggered an enhancement of total phenols and antioxidant ability in cv. Traveler cultivated in the absence of P. Seven flavonoids were identified, with the most relevant being derivatives of apigenin and luteonarin. Although we did not observe a clear effect of Si on the content of individual phenolics in cv. Sebastian exposed to P stress, an increment in the concentration of apigenin–pentoxide–hexoside was detected in cv. Traveler as a consequence of Si application to P-deficient plants. Differential expression of genes associated with the synthesis of phenolics was also induced by Si under P-stress. The transcript level of *HvPAL* diminished in cv. Sebastian and augmented in cv. Traveler due to Si supply under P limitation. Likewise, Si decreased *HvCHS* expression in cv. Sebastian grown without P, whereas an up-regulation of *HvCHS* was observed when Si was applied to cv. Traveler grown with low P. Acknowledgments. FONDECYT Regular Project N°1201257 and FONDECYT Postdoctoral Project N°3200901.

Keywords: silicon; P stress; antioxidant phenols; phenylalanine ammonia lyase gene; chalcone synthase gene; barley



Citation: Cartes, P.; Vega, I.; Pontigo, S. Silicon Differentially Modulates Phenolic Metabolism and Antioxidant Capacity of Barley Cultivars Subjected to Phosphorus Deficiency. *Biol. Life Sci. Forum* **2021**, *3*, 53. <https://doi.org/10.3390/IECAG2021-09707>

Academic Editor: Youssef Rouphael

Published: 1 May 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Supplementary Materials: The poster presentation is available online at <https://www.mdpi.com/article/10.3390/IECAG2021-09707/s1>.

Conflicts of Interest: The authors declare no conflict of interest.