



Article

Glutathione Is Involved in Resistance of Oilseed Rape to Powdery Mildew [†]

Elena Mikhaylova * , Michael Shein and Valentin Alekseev

Institute of Biochemistry and Genetics UFRC RAS, Prospekt Oktyabrya 71, 450054 Ufa, Russia; mikeshenoda@yandex.ru (M.S.); valentin-1994@yandex.ru (V.A.)

* Correspondence: mikhele@list.ru

[†] Presented at the 1st International Electronic Conference on Plant Science, 1–15 December 2020; Available online: <https://iecps2020.sciforum.net/>.

Abstract: Oilseed rape (*Brassica napus*) is an economically important crop. In temperate climate powdery mildew *Erysiphe cruciferaum* can reduce its yields dramatically. Nevertheless, cultivars resistant to this fungal disease have not been selected yet. Plants, grown in the field and in the light chamber, were infected with powdery mildew. After 30 days there were severe signs of infection on most of the plants, however several individuals were less susceptible to it. Leaves of *B. napus* were used to measure the level of GSH, GSSG and GST by *spectrophotometric* assay. The level of total glutathione was higher in plants with increased resistant to powdery mildew than in both severely infected plants and control plants that were not subjected to the infection. The GST activity in resistant and control plants was at the same level, however in susceptible plants it was lower. Therefore, oilseed rape with naturally increased glutathione content can be used in the selection of resistant cultivars. GST and GSH genes should be also considered as targets for genetic engineering and genome editing.

Keywords: *Brassica napus*; oilseed rape; rapeseed; powdery mildew; *Erysiphe cruciferaum*; GST; GSH; GSSG; oxidative stress



Citation: Mikhaylova, E.; Shein, M.; Alekseev, V. Glutathione Is Involved in Resistance of Oilseed Rape to Powdery Mildew. *Biol. Life Sci. Forum* **2021**, *4*, 28. <https://doi.org/10.3390/IECPS2020-08749>

Academic Editor: Yoselin Benitez-Alfonso

Published: 1 December 2020

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Brassica napus L. (oilseed rape, rapeseed) is used as a source of vegetable oil and protein, animal feed and biodiesel. It is one of the most important crops in moderate climate zone, due to its remarkable cold tolerance. However, oilseed rape is highly susceptible to fungal diseases, including powdery mildew *Erysiphe cruciferaum* [1,2].

Powdery mildew is an obligate biotrophic pathogen, most common in the end of the growing season [1]. Infection can reduce the yield of rapeseed and its quality, causing chlorosis, necrosis and dehydration. Resistance to this pathogen is very rare in *B. napus* and was only achieved by hybridization with distant relatives [3]. Commercial cultivars resistant to this fungal disease have not been selected yet. The impossibility to grow *E. cruciferaum* in culture media apart from the plant complicates associated studies.

Plant physiological reactions to powdery mildew infection are poorly studied. Glutathione plays an important role in plant resistance to oxidative stress. *Reduced Glutathione* (GSH) is involved in detoxification of ROS and toxic molecules, catalyzed by glutathione S-transferases (GSTs). Glutathione disulfide (GSSG) is formed during this process [4]. Induction of the level of GSH and GST upon exposure to stress provides better protection of the plant cell. Although GSTs are usually associated with tolerance to heavy metals [5], there are data on the role of several GST genes in resistance to other stress factors such as extreme temperatures [6] and fungal diseases [7–10]. Therefore, it is supposed that described mechanisms can also work during plant–fungus interactions, they have never been studied in oilseed rape infected by *E. cruciferaum* and the actual level of GSTs and glutathione in healthy and infected plants have never been evaluated.

2. Experiments

2.1. Biological Material

B. napus of cultivar “Ratnik” were used in the experiments. Seeds were sown in the experimental plot in natural conditions and in the laboratory in vessels filled with commercial soil (Geolia, Russia) under 10,000 lux generated by LED grow light.

E. cruciferaum was isolated from the infected *B. napus* plants and distinguished on the basis of the sequencing of ITS DNA marker. Leaves of the experimental plants were rubbed with the leaf of the infected plant to spread infection. Control plants remained untreated.

After 30 days, 42 plants with severe signs of infection and 30 control plants, from both the laboratory and field experiments, were used for spectrophotometric GSH, GSSG and GST assay. Each plant with increased resistance to powdery mildew was studied individually.

2.2. Spectrophotometric Assay

Fresh leaves of *B. napus* were homogenized and used to measure the level of GSH, GSSG and GST by spectrophotometric assay simultaneously. Levels of glutathione and glutathione disulfide were determined using o-phthalaldehyde (79,760, Sigma-Aldrich, St. Louis, MO, USA) as a fluorescent reagent; a derivatization of GSH to prevent GSH autooxidation was performed using N-ethylmaleimide (E1271, Sigma-Aldrich) [11]. GST activity was measured using the model substrate 1-chloro-2,4-dinitrobenzene (138630, Sigma-Aldrich) [12]. All measurements were performed in 6 replications in a 96-well plate using a Perkin Elmer LS 55 Luminescence Spectrometer (US).

3. Results

Most of the *B. napus* plants demonstrated severe signs of infection within 30 days after treatment, however several individuals were less susceptible to it (Figure 1). In general, 12 plants with increased resistance to powdery mildew were found and examined. Out of them, six plants were cultivated in the field, and six in the laboratory.




Figure 1. Rapeseed, susceptible (A) and resistant (B) to powdery mildew.

There was a difference in GSH and GSSG content and GST activity not only between infected and control plants, but also between plants cultivated in laboratory and in the field. All measured parameters were increased in the latter. Therefore, the results were evaluated separately for laboratory and field experiment.

Resistant plants had extremely high glutathione content in comparison with susceptible plants (6–7 times higher in several plants; 2–3 times higher on average) (Figure 2); however the activity of GST never varied that much (Figure 3). In resistant and control plants it was at the same level, however in susceptible plants it was no more than three times lower. In untreated plants GST activity was stable and did not vary significantly, however some plants had increased glutathione levels.




Figure 2. Relative content of glutathione in the studied plants. Plants grown in the field (1–6) and in the lab (7–12) are highlighted in green and purple, respectively.




Figure 3. Relative content of GST in studied plants. Plants grown in the field (1–6) and in the lab (7–12) are highlighted in green and purple, respectively.

The GSH:GSSG level in most of the susceptible plants was lower than in resistant plants, which is a sign of heavier stress.

4. Discussion

Our results suggest that an increased level of glutathione can give plants an advantage during infection with powdery mildew. The analysis of control plants showed that there are individuals with a physiologically high level of glutathione, and they could be the ones showing resistance to the pathogen. These untreated plants also demonstrated higher a GSH:GSSG level, which is characteristic for the lower oxidative stress.

However, there are some data on the increased expression of *GST* genes in plants exposed to fungal infection [7–10] in wheat, tomato and barley; in our experiments there was no significant difference between control and resistant plants in *GST* activity. Many of the susceptible plants had reduced *GST* activity; therefore, this protein is also of research interest.

5. Conclusions

Oilseed rape with naturally increased glutathione content can be used in the selection of resistant cultivars. *GST* and *GSH* genes should be also considered as targets for genetic engineering and genome editing.

Author Contributions: E.M. conceived and designed the experiments, analyzed the data and wrote the paper; M.S. and V.A. performed the experiments. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Council on grants of the President of the Russian Federation MK-1146.2020.11.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data supporting the findings of this study are available within the article.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Uloth, M.B.; You, M.P.; Barbetti, M.J. Plant age and ambient temperature: Significant drivers for powdery mildew (*Erysiphe cruciferarum*) epidemics on oilseed rape (*Brassica napus*). *Plant Pathol.* **2018**, *67*, 445–456. [\[CrossRef\]](#)
2. Alkooranee, J.T.; Liu, S.; Aledan, T.R.; Yin, Y.; Li, M. First report of powdery mildew caused by *Erysiphe cruciferarum* on *Brassica napus* in China. *Plant Dis.* **2015**, *99*, 1651. [\[CrossRef\]](#)
3. Gong, Q.; Dai, C.Y.; Zhang, X.H.; Wang, X.L.; Huang, Z.; Xu, A.X.; Dong, J.G.; Yu, C.Y. Towards breeding of rapeseed (*Brassica napus*) with alien cytoplasm and powdery mildew resistance from Ethiopian mustard (*Brassica carinata*). *Breed. Sci.* **2020**, 20017. [\[CrossRef\]](#) [\[PubMed\]](#)
4. Gullner, G.; Komives, T. Defense reactions of infected plants: Roles of glutathione and glutathione S-transferase enzymes. *Acta Phytopathol. Entomol. Hung.* **2006**, *41*, 3–10. [\[CrossRef\]](#)
5. Kuluev, B.R.; Berezhneva, Z.A.; Mikhaylova, E.V.; Postrigan, B.N.; Knyazev, A.V. Productivity and stress-tolerance of transgenic tobacco plants with constitutive expression of rapeseed glutathione synthetase gene BnGSH. *Ecol. Genet.* **2017**, *15*, 12–19. [\[CrossRef\]](#)
6. Kouno, T.; Ezaki, B. Multiple regulation of Arabidopsis AtGST11 gene expression by four transcription factors under abiotic stresses. *Physiol. Plant.* **2013**, *148*, 97–104. [\[CrossRef\]](#) [\[PubMed\]](#)
7. Mauch, F.; Dudler, R. Differential induction of distinct glutathione-S-transferases of wheat by xenobiotics and by pathogen attack. *Plant Physiol.* **1993**, *102*, 1193–1201. [\[CrossRef\]](#) [\[PubMed\]](#)
8. Wang, J.M.; Liu, H.Y.; Xu, H.M.; Li, M.; Kang, Z.S. Analysis of differential transcriptional profiling in wheat infected by *Blumeria graminis* f. sp. *tritici* using GeneChip. *Mol. Biol. Rep.* **2012**, *39*, 381–387. [\[CrossRef\]](#) [\[PubMed\]](#)
9. Pei, D.; Ma, H.; Zhang, Y.; Ma, Y.; Wang, W.; Geng, H.; Wu, J.; Li, C. Virus-induced gene silencing of a putative glutathione S-transferase gene compromised Ol-1-mediated resistance against powdery mildew in tomato. *Plant Mol. Biol. Rep.* **2011**, *29*, 972–978. [\[CrossRef\]](#)
10. Hossain, M.A.; Mostofa, M.G.; Diaz-Vivancos, P.; Burritt, D.J.; Fujita, M.; Tran, L.S.P. (Eds.) *GlutaThione in Plant Growth, Development, and Stress Tolerance*; Springer International Publishing: Cham, Switzerland, 2017. [\[CrossRef\]](#)

-
11. Hissin, P.J.; Hilf, R. Fluorometric determination of glutathione using o-phthaldialdehyde. *Anal. Biochem.* **1976**, *74*, 214–226. [[CrossRef](#)]
 12. Sappl, P.G.; Carroll, A.J.; Clifton, R.; Lister, R.; Whelan, J.; Harvey Millar, A.; Singh, K.B. The Arabidopsis glutathione transferase gene family displays complex stress regulation and co-silencing multiple genes results in altered metabolic sensitivity to oxidative stress. *Plant J.* **2009**, *58*, 53–68. [[CrossRef](#)] [[PubMed](#)]