



Abstract

# Comparative Study on the Physicochemical Properties of Extruded Fortified Rice Kernels Produced from Different Rice Varieties with Their Corresponding Rice Varieties <sup>†</sup>

A. Nithya \*, Chandrakant Genu Dalbhagat and Hari Niwas Mishra

Agricultural and Food Engineering Department, Indian Institute of Technology Kharagpur, Kharagpur 721302, West Bengal, India

\* Correspondence: micnithya@gmail.com

<sup>†</sup> Presented at the 2nd International Electronic Conference on Foods—Future Foods and Food Technologies for a Sustainable World, 15–30 October 2021; Available online: <https://foods2021.sciforum.net/>.

**Abstract:** Broken rice is a by-product of rice milling industries that could be refabricated to produce fortified rice kernels (FRK), a novel health product. FRK are manufactured by blending broken rice flour with a vitamin-mineral premix and extruding it to rice-shaped kernels. These FRK are then mixed with raw rice in a specific ratio to produce micronutrient fortified rice, which reduces the cost of rice fortification. For this purpose, the FRK produced from a rice variety must match the properties of the same rice variety. Therefore, the study aimed to collect the broken rice of different varieties (raw and parboiled) and produce FRK thereof. In this study, four parboiled and three raw rice varieties were collected and processed to produce FRK. The FRK's apparent amylose content (AAC), color, density, functional properties, and cooking time were compared with that of the corresponding native rice variety. The AAC of the native rice varieties was in the range of 20% to 24%. After extrusion, the AAC of the FRK (18.02% to 22.79%) decreased. The lightness, redness, and yellowness values of the FRK varied between 68 to 77, 0.9 to 3, and 16.02 to 19.06, respectively. The total color change ( $\Delta E$ ) of FRK were < 3.6 for the raw rice, while were 7.19–10.29 for parboiled rice. The difference between the bulk density of all varieties of raw rice and FRK was < 9%. The water absorption index (WAI), water solubility index (WSI), and cooking time of the FRK were lower than that of the native rice. This study showed that there were significant differences between the properties of FRK of different rice varieties. Further studies on the optimization of the processing conditions for rice varieties based on amylose: amylopectin ratio would help the production of FRK that matches the characteristics of native rice variety, thus facilitating sustainable production of low-cost novel health food.

**Keywords:** fortified rice kernels; by-product utilization; physicochemical properties; extrusion; fortification



**Citation:** Nithya, A.; Dalbhagat, C.G.; Mishra, H.N. Comparative Study on the Physicochemical Properties of Extruded Fortified Rice Kernels Produced from Different Rice Varieties with Their Corresponding Rice Varieties. *Biol. Life Sci. Forum* **2021**, *6*, 114. <https://doi.org/10.3390/Foods2021-11065>

Academic Editor: Diego A. Moreno

Published: 14 October 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/Foods2021-11065/s1>, Poster: Comparative study on the physicochemical properties of extruded fortified rice kernels produced from different rice varieties with their corresponding rice varieties.

**Author Contributions:** Conceptualization, A.N., C.G.D. and H.N.M.; methodology, A.N., C.G.D. and H.N.M.; validation, A.N., C.G.D. and H.N.M.; formal analysis, A.N.; investigation, A.N.; resources, H.N.M.; data curation, A.N.; writing—original draft preparation, A.N.; writing—review and editing, A.N., C.G.D. and H.N.M.; visualization, A.N. and H.N.M.; supervision, H.N.M.; project administration, H.N.M.; funding acquisition, H.N.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** The research was funded by Department of Biotechnology (DBT), Government of India, New Delhi. Sanction order No. BT/PR27470/PFN/20/1336/2018 Dated: 29/03/2019.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Data is contained within the article or Supplementary Material.

**Acknowledgments:** The authors gratefully acknowledge the financial support of the Department of Biotechnology (DBT), Government of India, New Delhi, for the 'FRK' project.

**Conflicts of Interest:** The authors have declared no conflict of interest for this article.