

Proceeding Paper

Identification of Potential Protein Biomarkers in a Depressed Chinese Malaysian University Student Using Liquid Chromatography-Tandem Mass Spectrometry [†]

Sin Yee Yap ¹, Chai Nien Foo ^{1,2,*}, Yang Mooi Lim ^{1,3}, Foong Leng Ng ^{1,4}, Pek Yee Tang ⁵, Jagjit Kaur Najar Singh ^{1,6}, Sherina Mohd-Sidik ⁷ and Kai-Shuen Pheh ⁸

- ¹ Centre for Cancer Research, M. Kandiah Faculty of Medicine and Health Sciences, Universiti Tunku Abdul Rahman, PT21144, Jalan Sungai Long, Bandar Sungai Long, Kajang 43000, Malaysia; sinyeeyap24@utar.my (S.Y.Y.); ymlim@utar.edu.my (Y.M.L.); ngfl@utar.edu.my (F.L.N.); jagjitk@utar.edu.my (J.K.N.S.)
 - ² Department of Population Medicine, M. Kandiah Faculty of Medicine and Health Sciences, Universiti Tunku Abdul Rahman, PT21144, Jalan Sungai Long, Bandar Sungai Long, Kajang 43000, Malaysia
 - ³ Department of Pre-Clinical Science, M. Kandiah Faculty of Medicine and Health Sciences, Universiti Tunku Abdul Rahman, PT21144, Jalan Sungai Long, Bandar Sungai Long, Kajang 43000, Malaysia
 - ⁴ Department of Traditional Chinese Medicine, M. Kandiah Faculty of Medicine and Health Sciences, Universiti Tunku Abdul Rahman, PT21144, Jalan Sungai Long, Bandar Sungai Long, Kajang 43000, Malaysia
 - ⁵ Department of Mechatronics and Biomedical Engineering, Lee Kong Chian Faculty of Engineering and Science, Universiti Tunku Abdul Rahman, PT21144, Jalan Sungai Long, Bandar Sungai Long, Kajang 43000, Malaysia; tangpy@utar.edu.my
 - ⁶ Department of Nursing, M. Kandiah Faculty of Medicine and Health Sciences, Universiti Tunku Abdul Rahman, PT21144, Jalan Sungai Long, Bandar Sungai Long, Kajang 43000, Malaysia
 - ⁷ Department of Psychiatry, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, UPM Serdang, Seri Kembangan 43400, Malaysia; sherina@upm.edu.my
 - ⁸ Department of Psychology and Counselling, Faculty of Arts and Social Science, Universiti Tunku Abdul Rahman, Jalan Universiti, Bandar Barat, Kampar 31900, Malaysia; phehks@utar.edu.my
- * Correspondence: foocn@utar.edu.my
- [†] Presented at the 2nd International Electronic Conference on Biomedicines, 1–31 March 2023; Available online: <https://ecb2023.sciforum.net/>.



Citation: Yap, S.Y.; Foo, C.N.; Lim, Y.M.; Ng, F.L.; Tang, P.Y.; Najar Singh, J.K.; Mohd-Sidik, S.; Pheh, K.-S. Identification of Potential Protein Biomarkers in a Depressed Chinese Malaysian University Student Using Liquid Chromatography-Tandem Mass Spectrometry. *Med. Sci. Forum* **2023**, *21*, 10. <https://doi.org/10.3390/ECB2023-14089>

Academic Editor: Masaru Tanaka

Published: 9 June 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Depression is a serious psychological disorder with high prevalence rates, especially among university students. Serum proteins related to the immune system and oxygen and lipid transfer could have contributing roles in the development of depression and could act as biomarkers for depression. Currently, there is a lack of accurate biological methods that can be used to diagnose depression. Biomarkers could be an inexpensive and convenient way to predict depression and understand its pathophysiology. This study aimed to screen the serum proteome profile of a depressed student for the identification of potential depression biomarkers. A Malaysian private university student who was recruited from the pre-test study ($n = 10$) was further analyzed for serum proteome due to the fact that he was depressed, with scores of 15 out of 27 on the Patient Health Questionnaire (PHQ-9). After depleting the high-abundance proteins from the serum sample, liquid chromatography-tandem mass spectrometry (LC-MS/MS) was performed to identify the expressed proteins. A total of 224 proteins were identified. Globins, globulins, apolipoproteins and glycoproteins were most commonly detected. Here, we show the potential biomarkers that can be used to identify depression vulnerable individuals. These findings may be relevant to the development of new diagnostic and treatment strategies. However, further studies with larger sample sizes and healthy controls are needed to confirm the role of these candidate biomarkers for the prediction and diagnosis of depression.

Keywords: depression; depressive disorder; serum proteome; proteomics; biomarker; protein markers; identification; university student; case study; LCMS

1. Introduction

Mental health disorders are among the biggest health issues affecting Malaysians [1]. National Health and Morbidity Survey 2019 reported that nearly half a million of Malaysians suffered from depression [2]. According to a study in 2020, 42.3% of Malaysian adults had depression [3]. University students are more vulnerable to depression compared to other populations. A study in 2018 showed that 29.4% of university students in Malaysia were reported to have depression [4]. A pilot study conducted by the authors showed that Malaysian university students had a prevalence of depression of 33.8% [5]. Depression can cause a series of emotional and physical problems that interfere with a student's ability to work, eat, sleep and enjoy life. Resources provided by universities to combat mental health problems are limited, and this has indirectly led to the increasing prevalence of depression among students.

For the past few decades, researchers have attempted to link diagnostic and prognostic biomarkers for psychiatric disorders including depression, schizophrenia, and bipolar disorder. Bodily fluids such as blood, urine, and cerebral spinal fluid are some of the easily accessible sources to detect these psychiatric biomarkers [6]. For example, serum S100B was found to be lower in medicated young patients, compared with those who were drug-free, and healthy controls [7]. In addition, decreased serum total proteins, albumin, and globulin were found to be associated with depressive severity in schizophrenia [8]. In a recent study conducted among Chinese older adults, higher levels of immunoglobulin A, lower levels immunoglobulin M and complement C3 were found in depressed group [9]. The discovery of these biomarkers could provide better insights into mental illness mechanisms and aid the diagnosis of psychiatric disorders, as well as accelerating the development of targeted therapy tailored to each individual. This study aimed to screen the serum proteome profile of a depressed student for the identification of a potential serum protein marker panel for the detection of depression.

2. Methods

Written consent was obtained prior to the data collection. Patient health questionnaire-9 (PHQ-9) [10] was used to assess the presence of depression. Blood samples were collected in a red vacutainer with no anticoagulants, and the separated serum was stored at -80°C before analysis. The serum was analyzed by means of QT of liquid chromatography with tandem mass spectrometry (LC-MS/MS). Agilent MassHunter data acquisition software (Agilent Technologies Inc, Santa Clara, CA, USA) and PEAKS 7.0 software (Bioinformatics Solutions Inc, Waterloo, ON, Canada) were used for protein profiling.

3. Results

Globulins were most commonly detected, followed by globins, apolipoproteins, glycoproteins, inhibitory proteins, complement proteins, binding proteins, ceruloplasmin, albumin, and luminal proteins, as shown in Table 1.

Table 1. List of identified protein biomarkers.

Type	Protein	−10lgP
Globulins	Alpha-2-macroglobulin	209.22
	Serotransferrin	170.6
	Immunoglobulin	166.94~64.69
	Vitamin D-binding protein	158.89
	Prothrombin	145.5
	Angiotensin	104.31
	Protein AMBP	72.23
	Transthyretin	61.89

Table 1. Cont.

Type	Protein	−10lgP
Globins	Haptoglobin	127.26
	Hemoglobin subunit beta	87.75
	Hemoglobin alpha	53.68
Apolipoproteins	Hemoglobin C	53.68
	Apolipoprotein B	218.02
	Apolipoprotein H	134.17
	Apolipoprotein A	132.28~71.93
	Apolipoprotein C	70.65~37.24
	Apolipoprotein E	55.66
	Apolipoprotein D	24.17
Glycoproteins	Beta-2-glycoprotein	134.17
	Alpha-2-HS-glycoprotein	132.92
	Hemopexin	124.8
	Vitronectin	117.28
	Antithrombin	102.28
	Alpha-1B-glycoprotein	99.58
	Afamin	88.28
	Alpha-1-acid glycoprotein	87.65~74.17
	Fibronectin	87.14
	Histidine-rich glycoprotein	82.56
	Clusterin	75.45
	Complement factor H	61.76
	Complement C3	257.29
	C4a anaphylatoxin	201.95
	C3/C5 convertase	103.31
Complement proteins	Complement factor B	103.31
	Complement component C6	63.39
Binding proteins	Hepatitis B virus receptor binding protein	162.98
	Epididymis secretory sperm binding protein	160.87~99.58
	Gelsolin	95.4
	Actin-depolymerizing factor	95.4
	Retinol-binding protein	81.8
Inhibitory proteins	C4b-binding protein alpha chain	81.16
	serpins	113.58~65.31
	Inter-alpha-trypsin inhibitor heavy chain H2	125.8
	Inter-alpha-trypsin inhibitor heavy chain H1	120.32
	Inter-alpha-trypsin inhibitor heavy chain H4	113.23
	(ITIH4) protein	
	Kininogen-1	108.14
Other	Bradykinin	108.14
	Ceruloplasmin	141.67
	Albumin	235.66~225.4
	Luminal proteins	169.56~133.23

Note: −10lgP represents the level of significance.

4. Discussions

The functions of these identified proteins are mostly related to our immune system and lipid transport. The contributing biological cause behind the link between the immune system and depression could be due to inflammation. When we are sick, our immune system creates an inflammatory response that makes us feel sad, irritable or unmotivated in order to keep us in bed in order for our body to devote body resources to fight and heal. When inflammatory cytokines in the body reach a certain threshold, our brain will initiate its own inflammatory response, and cause the macrophages to pump out cytokines that not only attack invaders such germs, but can also harm healthy tissues throughout body and brain [11]. Cytokines can also alter the neurotransmitter systems involved in the development of depression [12]. Meta-analysis in 2019 demonstrated alterations of cytokines levels in patients with antidepressant outcomes [13].

C-reactive protein (CRP) is an inflammation protein marker, and it increases during infection [14]. Numerous studies correlated elevated CRP levels with the presence of depression and its severity. A meta-analysis consisting of 30 studies with 11,813 participants showed that the presence of low-grade inflammation and elevated CRP levels were found in depressed patients [15]. Additionally, higher CRP levels were associated with depressive symptoms from Netherlands Study of Depression and Anxiety (NESDA) and UK Biobank cohorts [16].

Neurotransmitter systems can be affected by inflammation via the reduction in monoamine synthesis. Monoamine neurotransmitters such as dopamine, norepinephrine, and serotonin are primarily associated with the development of depression [17]. Increased dopamine signaling was found to be associated with lower levels of depressive symptoms among Asian samples [18]. However, a contradictory result was found among Caucasian samples in the same study [18]. In addition, neurotransmitters such as glutamate and GABA can also be affected by inflammation. A study on depressed adolescents found that higher levels of pro-inflammatory cytokines are associated with higher levels of glutamate [19]. Increased body levels of glutamic acid could lead to hyperglutamatergic neurotransmission, which may result in the occurrence of depression [20]. Another study showed that patients with a first episode of depression had significantly decreased glutamate and increased GABA levels compared to healthy controls [21].

Serum lipids may be linked to depression via the alteration of serotonin metabolism, the neurotransmitter in our brain that regulates our mood. Cholesterol has a crucial role in brain development and in neuron-to-neuron signaling [22]. Past studies showed that low-density lipoprotein (LDL) cholesterol levels are associated with depression [23–25]. LDL cholesterol can reduce the availability of serotonin and increase depression risk [26] by directly impairing the function of the serotonin 1A receptor in the brain. Activation of this receptor is associated with many other neurotransmitters related to the recovery and repair of neurons, as well as depression [27]. Daut and Fonken [28] suggested that alterations in the serotonin system may disrupt circadian rhythms and increase a person's susceptibility to depression. A recent study found a rapid response to selective serotonin reuptake inhibitors (SSRIs) in post-COVID-19 depressed patients, and their findings suggest that SSRIs could be an effective depression treatment option for the neuroinflammation triggered by SARS-CoV-2 [29]. However, there is a recent review that opposes the idea that serotonin causes depression. The researchers found no consistent evidence showing an association between serotonin and depression [30].

5. Conclusions

In short, most protein markers identified in this study were related to inflammation and lipid transport. The discovery and identification of easily accessible depression biomarkers such as serum proteins could enhance our understanding of the pathophysiology of depression, as well as providing possible treatment targets for depression. Future studies with healthy controls are much needed to further confirm the role of these biomarkers in depression. Future studies could also look into the relationship between these inflammation markers and dietary patterns, so that we can modify our food intake for the early prevention of depression.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ECB2023-14089/s1>.

Author Contributions: Conceptualization, C.N.F., Y.M.L., F.L.N. and P.Y.T.; methodology, C.N.F., Y.M.L., F.L.N. and S.Y.Y.; validation, C.N.F., Y.M.L., F.L.N., S.Y.Y., K.-S.P. and S.M.-S.; formal analysis, C.N.F. and S.Y.Y.; investigation, C.N.F., Y.M.L., F.L.N. and S.Y.Y.; data curation, S.Y.Y.; writing—original draft preparation, S.Y.Y.; writing—review and editing, C.N.F., Y.M.L., F.L.N., S.Y.Y., K.-S.P., S.M.-S., P.Y.T. and J.K.N.S.; supervision, C.N.F., Y.M.L., and F.L.N.; project administration, C.N.F., Y.M.L., F.L.N. and S.Y.Y.; funding acquisition, C.N.F., Y.M.L., F.L.N., P.Y.T., S.M.-S. and J.K.N.S. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by the Universiti Tunku Abdul Rahman Research Fund (UTARRF), (Project number IPSR/RMC/UTARRF/2019-C2/F01 and IPSR/RMC/UTARRF/2022-C1/F02).

Institutional Review Board Statement: The study was conducted according to the Declaration of the UTAR Research Ethics and Code of Conduct guidelines, Code of Practice for Research Involving Humans, and approved by the UTAR Scientific and Ethical Review Committee (SERC) (U/SERC/194/2022).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Not applicable.

Acknowledgments: The authors thank the student studied, without whom the research would not be possible.

Conflicts of Interest: The authors declare no conflict of interest. The funder had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Chua, S.N. The Economic Cost of Mental Disorders in Malaysia. *Lancet Psychiatry* **2020**, *7*, e23. [\[CrossRef\]](#) [\[PubMed\]](#)
2. Institute for Public Health (IPH). *National Health and Morbidity Survey (NHMS) 2019: Non-Communicable Diseases, Healthcare Demand, and Health Literacy*; Institute for Public Health (IPH): Oakland, CA, USA, 2020.
3. Perveen, A.; Hamzah, H.; Ramlee, F.; Othman, A.; Minhad, M. Mental Health and Coping Response among Malaysian Adults during COVID-19 Pandemic Movement Control Order. *J. Crit. Rev.* **2020**, *7*, 2020.
4. Islam, M.A.; Low, W.Y.; Tong, W.T.; Choo, C.W.Y.; Abdullah, A. Factors Associated with Depression among University Students in Malaysia: A Cross-Sectional Study | KNE Life Sciences. In Proceedings of the 2nd International Meeting of Public Health 2016 (IMOPH), Depok, Indonesia, 19–20 November 2016; pp. 415–427. [\[CrossRef\]](#)
5. Yap, S.Y.; Foo, C.N.; Lim, Y.M.; Ng, F.L.; Mohd-Sidik, S.; Tang, P.Y.; Najjar Singh, J.K.; Pheh, K.-S. Traditional Chinese Medicine Body Constitutions and Psychological Determinants of Depression among University Students in Malaysia: A Pilot Study. *Int. J. Environ. Res. Public Health* **2021**, *18*, 5366. [\[CrossRef\]](#) [\[PubMed\]](#)
6. García-Gutiérrez, M.S.; Navarrete, F.; Sala, F.; Gasparyan, A.; Austrich-Olivares, A.; Manzanares, J. Biomarkers in Psychiatry: Concept, Definition, Types and Relevance to the Clinical Reality. *Front. Psychiatry* **2020**, *11*, 432. [\[CrossRef\]](#)
7. Rajewska-Rager, A.; Dmitrzak-Weglarz, M.; Kapelski, P.; Lepczynska, N.; Pawlak, J.; Twarowska-Hauser, J.; Skibinska, M. Longitudinal Assessment of S100B Serum Levels and Clinical Factors in Youth Patients with Mood Disorders. *Sci. Rep.* **2021**, *11*, 11973. [\[CrossRef\]](#) [\[PubMed\]](#)
8. Yin, X.; Cai, Y.; Zhu, Z.; Zhai, C.; Li, J.; Ji, C.; Chen, P.; Wang, J.; Wu, Y.; Chan, R.; et al. Associations of Decreased Serum Total Protein, Albumin, and Globulin with Depressive Severity of Schizophrenia. *Front. Psychiatry* **2022**, *13*, 957671. [\[CrossRef\]](#)
9. Sun, Z.; Lin, J.; Zhang, Y.; Yao, Y.; Huang, Z.; Zhao, Y.; Zhang, P.; Fu, S. Association between Immunoglobulin A and Depression in Chinese Older Adults: Findings from a Cross-Sectional Study. *Immun. Ageing* **2022**, *19*, 21. [\[CrossRef\]](#)
10. Kroenke, K.; Spitzer, R.L.; Williams, J.B. The PHQ-9: Validity of a Brief Depression Severity Measure. *J. Gen. Intern. Med.* **2001**, *16*, 606–613. [\[CrossRef\]](#)
11. Kany, S.; Vollrath, J.T.; Relja, B. Cytokines in Inflammatory Disease. *Int. J. Mol. Sci.* **2019**, *20*, 6008. [\[CrossRef\]](#)
12. Afridi, R.; Suk, K. Neuroinflammatory Basis of Depression: Learning From Experimental Models. *Front. Cell. Neurosci.* **2021**, *15*, 691067. [\[CrossRef\]](#)
13. Liu, J.J.; Wei, Y.B.; Strawbridge, R.; Bao, Y.; Chang, S.; Shi, L.; Que, J.; Gadad, B.S.; Trivedi, M.H.; Kelsoe, J.R.; et al. Peripheral Cytokine Levels and Response to Antidepressant Treatment in Depression: A Systematic Review and Meta-Analysis. *Mol. Psychiatry* **2020**, *25*, 339–350. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Sproston, N.R.; Ashworth, J.J. Role of C-Reactive Protein at Sites of Inflammation and Infection. *Front. Immunol.* **2018**, *9*, 754. [\[CrossRef\]](#) [\[PubMed\]](#)
15. Osimo, E.F.; Baxter, L.J.; Lewis, G.; Jones, P.B.; Khandaker, G.M. Prevalence of Low-Grade Inflammation in Depression: A Systematic Review and Meta-Analysis of CRP Levels. *Psychol. Med.* **2019**, *49*, 1958–1970. [\[CrossRef\]](#)

16. Milaneschi, Y.; Kappelmann, N.; Ye, Z.; Lamers, F.; Moser, S.; Jones, P.B.; Burgess, S.; Penninx, B.W.J.H.; Khandaker, G.M. Association of Inflammation with Depression and Anxiety: Evidence for Symptom-Specificity and Potential Causality from UK Biobank and NESDA Cohorts. *Mol. Psychiatry* **2021**, *26*, 7393–7402. [\[CrossRef\]](#)
17. Shao, X.; Zhu, G. Associations Among Monoamine Neurotransmitter Pathways, Personality Traits, and Major Depressive Disorder. *Front. Psychiatry* **2020**, *11*, 381. [\[CrossRef\]](#)
18. Avinun, R.; Nevo, A.; Radtke, S.R.; Brigidi, B.D.; Hariri, A.R. Divergence of an Association between Depressive Symptoms and a Dopamine Polygenic Score in Caucasians and Asians. *Eur. Arch. Psychiatry Clin. Neurosci.* **2020**, *270*, 229–235. [\[CrossRef\]](#)
19. Ho, T.C.; Teresi, G.I.; Segarra, J.R.; Ojha, A.; Walker, J.C.; Gu, M.; Spielman, D.M.; Sacchet, M.D.; Jiang, F.; Rosenberg-Hasson, Y.; et al. Higher Levels of Pro-Inflammatory Cytokines Are Associated With Higher Levels of Glutamate in the Anterior Cingulate Cortex in Depressed Adolescents. *Front. Psychiatry* **2021**, *12*, 642976. [\[CrossRef\]](#) [\[PubMed\]](#)
20. Kumar, P.; Kraal, A.Z.; Prawdzik, A.M.; Ringold, A.E.; Ellingrod, V. Dietary Glutamic Acid, Obesity, and Depressive Symptoms in Patients With Schizophrenia. *Front. Psychiatry* **2021**, *11*, 620097. [\[CrossRef\]](#)
21. Draganov, M.; Vives-Gilabert, Y.; de Diego-Adeliño, J.; Vicent-Gil, M.; Puigdemont, D.; Portella, M.J. Glutamatergic and GABA-Ergic Abnormalities in First-Episode Depression. A 1-Year Follow-up 1H-MR Spectroscopic Study. *J. Affect. Disord.* **2020**, *266*, 572–577. [\[CrossRef\]](#)
22. Gliozzi, M.; Musolino, V.; Bosco, F.; Scicchitano, M.; Scarano, F.; Nucera, S.; Zito, M.C.; Ruga, S.; Carresi, C.; Macrì, R.; et al. Cholesterol Homeostasis: Researching a Dialogue between the Brain and Peripheral Tissues. *Pharmacol. Res.* **2021**, *163*, 105215. [\[CrossRef\]](#)
23. Wagner, C.J.; Musenbichler, C.; Böhm, L.; Färber, K.; Fischer, A.-I.; von Nippold, F.; Winkelmann, M.; Richter-Schmidinger, T.; Mühle, C.; Kornhuber, J.; et al. LDL Cholesterol Relates to Depression, Its Severity, and the Prospective Course. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* **2019**, *92*, 405–411. [\[CrossRef\]](#) [\[PubMed\]](#)
24. Kim, E.J.; Hong, J.; Hwang, J.-W. The Association between Depressive Mood and Cholesterol Levels in Korean Adolescents. *Psychiatry Investig.* **2019**, *16*, 737–744. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Jo, S.-Y.; Kwon, Y.-J.; Cho, A.-R. Serum Low-Density Lipoprotein Cholesterol Levels and Depressive Mood in Korean Adults: A Nationwide Population-Based Study. *Korean J. Fam. Med.* **2022**, *43*, 63–68. [\[CrossRef\]](#) [\[PubMed\]](#)
26. Han, A.L. Association between Lipid Ratio and Depression: A Cross-Sectional Study. *Sci. Rep.* **2022**, *12*, 6190. [\[CrossRef\]](#)
27. Vahid-Ansari, F.; Albert, P.R. Rewiring of the Serotonin System in Major Depression. *Front. Psychiatry* **2021**, *12*, 2275. [\[CrossRef\]](#)
28. Daut, R.A.; Fonken, L.K. Circadian Regulation of Depression: A Role for Serotonin. *Front. Neuroendocrinol.* **2019**, *54*, 100746. [\[CrossRef\]](#)
29. Mazza, M.G.; Zanardi, R.; Palladini, M.; Rovere-Querini, P.; Benedetti, F. Rapid Response to Selective Serotonin Reuptake Inhibitors in Post-COVID Depression. *Eur. Neuropsychopharmacol.* **2022**, *54*, 1–6. [\[CrossRef\]](#)
30. Moncrieff, J.; Cooper, R.E.; Stockmann, T.; Amendola, S.; Hengartner, M.P.; Horowitz, M.A. The Serotonin Theory of Depression: A Systematic Umbrella Review of the Evidence. *Mol. Psychiatry* **2022**, 1–14. [\[CrossRef\]](#)

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.