

Association between Air Pollutants and Cancer Incidence Rates in Japan: An Ecological Study [†]

Kohei Hasegawa ^{1,*} , Teruomi Tsukahara ^{1,2}  and Tetsuo Nomiyama ^{1,2} 

¹ Department of Preventive Medicine and Public Health, School of Medicine, Shinshu University, 3-1-1 Asahi, Matsumoto, Nagano 390-8621, Japan; tsukat@shinshu-u.ac.jp (T.T.); nomiyama@shinshu-u.ac.jp (T.N.)

² Department of Occupational Medicine, School of Medicine, Shinshu University, 3-1-1 Asahi, Matsumoto, Nagano 390-8621, Japan

* Correspondence: koheih@shinshu-u.ac.jp; Tel.: +81-263-37-2622; Fax: +81-263-37-3499

[†] Presented at the 3rd International Electronic Conference on Environmental Research and Public Health—Public Health Issues in the Context of the COVID-19 Pandemic, 11–25 January 2021; Available online: <https://ecerph-3.sciforum.net/>.

Abstract: We investigated the associations between the incidence rates of five typical cancer types and five air pollutants in 317 Japanese municipalities in 2017. We obtained the concentrations of the five air pollutants, i.e., sulfur dioxide (SO₂), nitric oxide (NO), nitric dioxide (NO₂), photochemical oxidants (O_x), and fine particulate matter (PM_{2.5}), from Japan's National Institute for Environmental Studies and calculated the yearly mean of each. We identified patients of both sexes with the five most common cancers in Japan's National Cancer Registry, which covers all cancer patients in the country. For males, we included prostate, stomach, colorectal, lung, and liver cancers. For females, we included breast, colorectal, lung, stomach, and uterine cancers. We calculated the Spearman's correlation coefficients between 25 pairs of air-pollutant concentrations and the age-standardized incidence rate of the cancer types for each sex. We used Poisson regression models to examine the dose–response relationships. We identified 11 significantly positive correlation coefficients at the Bonferroni-corrected alpha level for the five pollutants for the five cancers in both sexes. We observed significantly positive dose–response relationships between NO₂ and colorectal cancer and PM_{2.5} and lung cancer for both sexes. We also observed significant dose–response relationships between SO₂ and PM_{2.5} and liver cancer for males and between NO and lung cancer and NO₂ and breast cancer for females. We did not observe significant associations with prostate, stomach, or uterine cancer. Our findings support the concept that exposure to air pollutants increases cancer incidence rates.

Keywords: air pollution; environmental pollution; cancer; environmental epidemiology



Citation: Hasegawa, K.; Tsukahara, T.; Nomiyama, T. Association between Air Pollutants and Cancer Incidence Rates in Japan: An Ecological Study. *Med. Sci. Forum* **2021**, *4*, 15. <https://doi.org/10.3390/ECERPH-3-09054>

Academic Editor: Jon Øyvind Odland

Published: 11 January 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Air pollution is a primary public health issue worldwide [1]. Many studies have observed an association between long-term exposure to air pollution and the incidence of lung cancer [2]. In response, the International Agency for Research on Cancer (IARC) classified outdoor air pollution as a Group 1 carcinogen (carcinogenic to humans) in 2013 [3].

The relationships between air pollution and cancers other than lung cancer are less clear [2,4]. For example, colorectal cancer is one of the most prevalent cancers globally [5], but studies on the association between this cancer and air pollution are few in number and have obtained inconsistent results [4,6,7]. Laboratory studies have demonstrated a plausible association between air pollution and liver cancer [8], but there have been few epidemiological studies on this topic [9–11] or on cancers other than lung cancer in Japan.

Among the many air pollutants, the adverse effects of exposure to nitric oxide (NO) have rarely been examined compared to those of sulfur dioxide (SO₂), nitric oxide (NO_x),

nitric dioxide (NO₂), ozone (O₃), and particulate matter (PM) [2,12]. Nevertheless, laboratory studies implied that NO might play an essential role in cancer biology [13,14].

In Japan, the nationwide population-based cancer registry, i.e., the National Cancer Registry (NCR) was established in 2016 under the Cancer Registry Promotion Act of 2013 [15]. This law obligated all of Japan's hospitals and some clinics to report all cancer patients encountered therein. The NCR thus enabled the accurate capture of the nationwide cancer incidence for the first time in Japan.

By leveraging data from the NCR, we investigated the association between air pollutants and the incidence rates of five common cancers.

2. Methods

2.1. Study Period and Area

This study was a cross-sectional, ecological investigation. The study period was from 1 January to 31 December 2017. We included 424 municipalities with more than one monitoring station that measured all included air pollutants. We excluded 105 municipalities where all or part of the air pollutant data was missing. To reduce the risk of exposure misclassification, we also excluded two municipalities that were in the designated evacuation area after the Fukushima Daiichi Nuclear Power Plant accident in 2011 [16]. A final total of 317 municipalities were included in this study; they are located throughout the length of Japan.

2.2. Cancer Incidence Rates

We obtained anonymized NCR data from the National Cancer Center Japan (NCCJ). The data included the patients' five-year age group, sex, residence at the time of diagnosis, and the diagnosis coded in the International Classification of Diseases (ICD).

We set the outcome as the incidence of five most common cancer types in Japan [17]. For males, we included prostate (ICD10: C61, D075), stomach (C16, D002), colorectal (C18–C20, D010–D012), lung (C33–C34 or D021–D022), and liver cancers (C22–C24, D015). For females, we included breast (C50, D050–D059), colorectal, lung, stomach, and uterine cancer (C53–C55, D60–D73). Notably, unlike the previous report [17], we included *in situ* neoplasms (D00–D09) to increase the sensitivity of the analysis. We excluded patients whose age ($n = 8$) or sex ($n = 9$) data were missing.

We used the age-adjusted cancer incidence rates per million population as the target variables to adjust for the age structure differences between municipalities. We obtained the data of each municipality's population as of 1 January 2017 from the Basic Resident Registry. As the standard reference population, we used the 2000 World Standard Population values [18].

2.3. Concentrations of Air Pollutants

We obtained the monthly data of the concentrations of SO₂, NO, NO₂, photochemical oxidants (O_x), and fine particulate matter (PM_{2.5}) in 2017 from Japan's National Institute for Environmental Studies and calculated the yearly mean. As few monitoring stations in Japan measured O₃, we used O_x instead. We considered the data 'missing' when measurements were not available for >20% of days in a particular month. We calculated the geometric mean if data from more than one monitoring station were available in a single municipality.

2.4. Statical Analysis

To explore the associations between air pollutants and cancer, we calculated Spearman's correlation coefficients between the age-adjusted incidence rates and air pollutant concentrations for each sex. We tested a total of 50 hypotheses and used Bonferroni correction for multiple testing problems.

For the pairs of pollutant and cancer types that indicated a positive correlation at the Bonferroni-corrected alpha level, we used Poisson regression models to examine the dose-response [12,19]. We categorized the air pollutant concentrations into four quartiles.

We included four other air pollutants as confounders. We considered p -values < 0.05 significant. We performed all of the statistical analyses in R (ver. 4.0.3) [20].

3. Results and Discussion

We analyzed the cases of a total of 288,405 individuals diagnosed with cancer in Japan. Table 1 provides the numbers of cases and the age-adjusted incidence rates of each cancer type.

Table 1. The total number of cases and age-adjusted incidence rates per million population of each cancer type in the 317 Japanese municipalities in 2017.

Male			Female		
Cancer Type	No. of Cases	Age-Adjusted Incidence Rate ¹	Cancer Type	No. of Cases	Age-Adjusted Incidence Rate ¹
Prostate	35,247	55.71	Breast	42,823	99.32
Stomach	33,733	55.88	Colorectal	31,996	48.41
Colorectal	45,832	84.80	Lung	16,276	22.52
Lung	32,129	52.60	Stomach	15,250	20.40
Liver	14,614	24.04	Uterine	20,505	67.36

¹ Adjusted for the 2000 World Standard Population proportion.

Table 2 shows the air pollution levels by mean and quartiles and Spearman's correlation coefficients between air pollutants in the 317 Japanese municipalities in 2017. The mean concentrations of the air pollutants were as follows: SO₂, 1.52 ppb; NO, 2.31 ppb; NO₂, 9.34 ppb; O_x, 35.16 ppb; and PM_{2.5} 11.56 µg/m³. SO₂, NO, NO₂, and PM_{2.5} were positively correlated, but O_x showed negative correlations with the other pollutants.

Table 2. Air pollution levels and Spearman's correlation coefficients between air pollutants in 317 Japanese municipalities in 2017.

Air Pollutant, Unit	Mean (SD)	Quartile			Spearman's Correlation Coefficient					
		Q1	Q2	Q3	SO ₂	NO	NO ₂	O _x	PM _{2.5}	
SO ₂ , ppb	1.52 (1.03)	0.92	1.25	2.00	SO ₂	-	0.26 *	0.32 *	-0.11 *	0.25 *
NO, ppb	2.31 (1.59)	1.08	2.00	3.17	NO		-	0.82 *	-0.61 *	0.22 *
NO ₂ , ppb	9.34 (4.30)	5.96	8.67	12.50	NO ₂			-	-0.57 *	0.22 *
O _x , ppb	35.16 (3.20)	33.21	34.92	36.92	O _x				-	-0.01
PM _{2.5} , µg/m ³	11.56 (2.29)	10.11	11.62	13.06	PM _{2.5}					-

* Significant at the alpha level of 0.05.

Table 3 lists the Spearman's correlation coefficients between the age-adjusted cancer incidence rates and the air pollutant concentrations. A total of 18 correlation coefficients were significantly positive at an alpha level of 0.05, and 11 of them remained significant at the level of Bonferroni correlation of 0.001.

Table 3. Spearman's correlation coefficients between the age-adjusted cancer incidence of five typical cancer types and the air pollutants stratified by sex.

Cancer Type	Male					Cancer Type	Female				
	SO ₂	NO	NO ₂	O _x	PM _{2.5}		SO ₂	NO	NO ₂	O _x	PM _{2.5}
Prostate	0.01	0.09	0.15 ^a	-0.08	-0.14 ^a	Breast	0.19 ^b	0.34 ^b	0.39 ^b	-0.24 ^b	0.08
Stomach	-0.03	-0.12 ^a	-0.07	0.10	-0.04	Colorectal	0.05	0.25 ^b	0.30 ^b	-0.23 ^b	-0.03
Colorectal	0.02	0.14 ^a	0.19 ^b	-0.17 ^a	-0.03	Lung	0.11 ^a	0.20 ^b	0.18 ^a	-0.07	0.21 ^b
Lung	0.05	0.09	0.08	-0.03	0.26 ^b	Stomach	0.01	-0.06	-0.04	0.09	-0.12 ^a
Liver	0.20 ^b	-0.03	-0.11	0.12 ^a	0.42 ^b	Uterine	0.12 ^a	-0.12 ^a	-0.15 ^a	0.09	0.17 ^a

^a Significant at an alpha level of 0.05. ^b Significant at the Bonferroni-corrected alpha level of 0.001.

Table 4 provides the results of Poisson regression models of 11 pollutant–cancer pairs. We observed positive dose–response relationships between NO₂ and colorectal cancer and between PM_{2.5} and lung cancer for both sexes. For males, SO₂ and PM_{2.5} also showed positive dose–response relationships with liver cancer. For the females, NO and lung cancer and NO₂ and breast cancer also showed positive dose–response relationships.

Table 4. Poisson regression analysis of selected pairs between air pollutants and cancer types.

Cancer Type (Sex)	Liver (Male)		Breast (Female)			
	RR ^a (95% CI)	<i>p</i>	RR ^a (95% CI)	<i>p</i>		
SO ₂ conc., ppb						
≤0.92	1.00		1.00			
0.92–1.25	0.99 (0.92–1.05)	0.71	1.05 (1.01–1.08)	<0.01		
1.25–2.00	1.07 (1.00–1.14)	0.04	1.03 (1.00–1.07)	0.06		
>2.00	1.08 (1.01–1.15)	0.03	1.02 (0.99–1.06)	0.17		
<i>p</i> for trend		<0.01		0.25		
Cancer type (sex)	Breast (female)		Colorectal (female)		Lung (female)	
	RR ^a (95% CI)	<i>p</i>	RR ^a (95% CI)	<i>p</i>	RR ^a (95% CI)	<i>p</i>
NO conc., ppb						
≤1.08	1.00		1.00		1.00	
1.08–2.00	1.05 (1.01–1.08)	<0.01	0.97 (0.92–1.02)	0.19	1.02 (0.95–1.10)	0.56
2.00–3.17	1.03 (1.00–1.07)	0.06	1.01 (0.95–1.08)	0.67	1.08 (0.99–1.18)	0.09
>3.17	1.02 (0.99–1.06)	0.17	1.00 (0.92–1.08)	0.93	1.13 (1.01–1.26)	0.04
<i>p</i> for trend		0.25		0.76		0.03
Cancer type (sex)	Colorectal (male)		Breast (female)		Colorectal (female)	
	RR ^a (95% CI)	<i>p</i>	RR ^a (95% CI)	<i>p</i>	RR ^a (95% CI)	<i>p</i>
NO ₂ conc., ppb						
≤5.96	1.00		1.00		1.00	
5.96–8.67	0.98 (0.94–1.02)	0.25	1.01 (0.97–1.04)	0.76	1.13 (1.08–1.19)	<0.01
8.67–12.50	1.01 (0.97–1.05)	0.59	1.09 (1.05–1.14)	<0.01	1.12 (1.06–1.18)	<0.01
>12.50	1.11 (1.05–1.17)	<0.01	1.20 (1.14–1.26)	<0.01	1.24 (1.15–1.33)	<0.01
<i>p</i> for trend		<0.01		<0.01		<0.01
Cancer type (sex)	Lung (male)		Liver (male)		Lung (female)	
	RR ^a (95% CI)	<i>p</i>	RR ^a (95% CI)	<i>p</i>	RR ^a (95% CI)	<i>p</i>
PM _{2.5} conc., µg/m ³						
≤10.11	1.00		1.00		1.00	
10.11–11.62	1.01 (0.97–1.06)	0.63	1.05 (0.98–1.13)	0.13	0.97 (0.90–1.04)	0.37
11.62–13.06	1.05 (1.00–1.10)	0.05	1.17 (1.09–1.25)	<0.01	1.07 (1.00–1.15)	0.06
>13.06	1.12 (1.07–1.17)	<0.01	1.38 (1.29–1.48)	<0.01	1.10 (1.03–1.18)	<0.01
<i>p</i> for trend		<0.01		<0.01		<0.01

^a Adjusted for age and the other four air pollutants. CI: confidence interval, RR: relative risk.

In this ecological study, we examined the associations between air pollutants and cancer incidence rates in 317 Japanese municipalities, and our analyses revealed significant positive dose–response relationships in some pollutant–cancer pairs.

The association between lung cancer and air pollution is well established [2,21]. In accord with previous reports, we observed positive correlations between lung cancer and all included pollutants except O_x. Dose–response relationships between lung cancer and PM_{2.5} were present after adjusting for other air pollutants in both sexes. Our results thus provide further evidence of the associations between air pollutants—especially PM_{2.5}—and lung cancer.

As noted above, there have been few investigations of the relationship between air pollutants and cancer types other than lung cancer, but those investigations have suggested some associations. An analysis of the Cancer Prevention Study II results examined the associations between PM_{2.5}, NO₂, and O₃ and many cancer types; only the relationships between PM_{2.5} and kidney cancer, PM_{2.5} and bladder cancer, and NO₂ and colorectal cancer were significant [4]. A study of PM_{2.5} exposure in Taiwan reported that PM_{2.5} was associated with gastrointestinal, liver, and colorectal cancers but not stomach cancer [22]. Positive correlations between PM and lung, breast, uterine, and ovarian cancers but not other types of cancer were obtained in a study of PM exposure in Japan [23]. A 2018 review

demonstrated that breast cancer was associated with NO₂ and less clearly with PM_{2.5} [24]. The present study's results are similar to those of the previous studies in general, and some additional associations were revealed for the first time in the Japanese population.

Among the air pollutants, examinations of NO in epidemiological studies are scarce. We observed a positive dose–response relationship between NO and lung cancer in the female population in this study. An ecological study from Taiwan reported a positive association between NO and lung cancer of a specific cell type [12]. Further investigations are necessary to determine the above-described associations by using individual-level data.

There are several study limitations to address. This was a cross-sectional, ecological study. We observed significant correlations, but this does not establish an individual-level association or causation. We considered confounding by age and sex, but we did not consider other known confounders, such as smoking for lung cancer, human papillomavirus infection for cervical cancer, or other unknown confounders. Thus, the observed associations could be the results of factors other than air pollution. In the study's design, we assumed that the patients lived in the municipality for most of their lives, and that the yearly mean air pollutants concentrations in 2017 reasonably reflect the patients' long-term exposure. We observed strong positive correlations between the yearly mean concentration in 2017 and the 5-year mean concentration in 2012–2016 among 71 monitors (SO₂, $r = 0.86$; NO, $r = 0.94$; NO₂, $r = 0.98$; O_x, $r = 0.84$; PM_{2.5}, $r = 0.82$). However, this assumption may have caused exposure misclassifications and biased our results.

4. Conclusions

Our findings support the concept that exposure to air pollutants increases cancer incidence rates.

Author Contributions: Conceptualization, K.H., T.T. and T.N.; methodology, K.H.; software, K.H.; validation, T.T. and T.N.; formal analysis, K.H.; investigation, K.H.; resources, K.H.; data curation, K.H.; writing—original draft preparation, K.H.; writing—review and editing, T.T. and T.N.; supervision, T.T.; project administration, T.N. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Committee for Medical Ethics of Shinshu University School of Medicine (protocol code: 4700, date of approval: 10 April 2020).

Informed Consent Statement: Patient consent was waived due to the retrospective nature of the study and the use of anonymized aggregated data.

Data Availability Statement: The dataset analyzed during the current study is not publicly available due to data protection requirements from the NCCJ.

Acknowledgments: We thank the NCCJ for managing and providing the NCR data.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: A systematic analysis for the Global Burden of Disease Study 2017. *Lancet Lond. Engl.* **2018**, *392*, 1923–1994. [\[CrossRef\]](#)
2. Turner, M.C.; Andersen, Z.J.; Baccarelli, A.; Diver, W.R.; Gapstur, S.M.; Pope, C.A.; Prada, D.; Samet, J.; Thurston, G.; Cohen, A. Outdoor air pollution and cancer: An overview of the current evidence and public health recommendations. *CA. Cancer J. Clin.* **2020**, *70*, 460–479. [\[CrossRef\]](#)
3. Loomis, D.; Grosse, Y.; Lauby-Secretan, B.; Ghissassi, F.E.; Bouvard, V.; Benbrahim-Tallaa, L.; Guha, N.; Baan, R.; Mattock, H.; Straif, K. The carcinogenicity of outdoor air pollution. *Lancet Oncol.* **2013**, *14*, 1262–1263. [\[CrossRef\]](#)
4. Turner, M.C.; Krewski, D.; Diver, W.R.; Pope, C.A.; Burnett, R.T.; Jerrett, M.; Marshall, J.D.; Gapstur, S.M. Ambient Air Pollution and Cancer Mortality in the Cancer Prevention Study II. *Environ. Health Perspect.* **2017**, *125*, 087013. [\[CrossRef\]](#) [\[PubMed\]](#)
5. Bray, F.; Ferlay, J.; Soerjomataram, I.; Siegel, R.L.; Torre, L.A.; Jemal, A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J. Clin.* **2018**, *68*, 394–424. [\[CrossRef\]](#) [\[PubMed\]](#)

6. Wong, C.M.; Tsang, H.; Lai, H.K.; Thomas, G.N.; Lam, K.B.; Chan, K.P.; Zheng, Q.; Ayres, J.G.; Lee, S.Y.; Lam, T.H.; et al. Cancer Mortality Risks from Long-term Exposure to Ambient Fine Particle. *Cancer Epidemiol. Biomark. Prev. Publ. Am. Assoc. Cancer Res. Cosponsored Am. Soc. Prev. Oncol.* **2016**, *25*, 839–845. [[CrossRef](#)] [[PubMed](#)]
7. Ancona, C.; Badaloni, C.; Mataloni, F.; Bolignano, A.; Bucci, S.; Cesaroni, G.; Sozzi, R.; Davoli, M.; Forastiere, F. Mortality and morbidity in a population exposed to multiple sources of air pollution: A retrospective cohort study using air dispersion models. *Environ. Res.* **2015**, *137*, 467–474. [[CrossRef](#)] [[PubMed](#)]
8. Kim, J.W.; Park, S.; Lim, C.W.; Lee, K.; Kim, B. The Role of Air Pollutants in Initiating Liver Disease. *Toxicol. Res.* **2014**, *30*, 65–70. [[CrossRef](#)]
9. Pan, W.-C.; Wu, C.-D.; Chen, M.-J.; Huang, Y.-T.; Chen, C.-J.; Su, H.-J.; Yang, H.-I. Fine Particle Pollution, Alanine Transaminase, and Liver Cancer: A Taiwanese Prospective Cohort Study (REVEAL-HBV). *JNCI J. Natl. Cancer Inst.* **2016**, *108*. [[CrossRef](#)] [[PubMed](#)]
10. Pedersen, M.; Andersen, Z.J.; Stafoggia, M.; Weinmayr, G.; Galassi, C.; Sørensen, M.; Eriksen, K.T.; Tjønneland, A.; Loft, S.; Jaensch, A.; et al. Ambient air pollution and primary liver cancer incidence in four European cohorts within the ESCAPE project. *Environ. Res.* **2017**, *154*, 226–233. [[CrossRef](#)] [[PubMed](#)]
11. VoPham, T.; Bertrand, K.A.; Tamimi, R.M.; Laden, F.; Hart, J.E. Ambient PM_{2.5} air pollution exposure and hepatocellular carcinoma incidence in the United States. *Cancer Causes Control* **2018**, *29*, 563–572. [[CrossRef](#)] [[PubMed](#)]
12. Liaw, Y.-P.; Ting, T.-F.; Ho, C.-C.; Chiou, Z.-Y. Cell type specificity of lung cancer associated with nitric oxide. *Sci. Total Environ.* **2010**, *408*, 4931–4934. [[CrossRef](#)]
13. Lala, P.K.; Chakraborty, C. Role of nitric oxide in carcinogenesis and tumour progression. *Lancet Oncol.* **2001**, *2*, 149–156. [[CrossRef](#)]
14. Sanuphan, A.; Chunhacha, P.; Pongrakhananon, V.; Chanvorachote, P. Long-term nitric oxide exposure enhances lung cancer cell migration. *BioMed Res. Int.* **2013**, *2013*, 186972. [[CrossRef](#)]
15. Cancer Statistics in Japan '17. Available online: https://ganjoho.jp/en/professional/statistics/brochure/2017_en.html (accessed on 1 December 2020).
16. Transition of Evacuation Designated Zones. Available online: <https://www.pref.fukushima.lg.jp/site/portal-english/en03-08.html> (accessed on 1 December 2020).
17. Cancer Incidence of Japan 2017. Available online: <https://www.mhlw.go.jp/content/10900000/000624853.pdf> (accessed on 1 December 2020).
18. Ahmad, O.B.; Boschi-Pinto, C.; Lopez, A.D.; Murray, C.J.L.; Lozano, R.; Inoue, M. *Age Standardization of Rates: A New Who Standard*; World Health Organization: Geneva, Switzerland, 2000.
19. Huang, H.-H.; Huang, J.-Y.; Lung, C.-C.; Wu, C.-L.; Ho, C.-C.; Sun, Y.-H.; Ko, P.-C.; Su, S.-Y.; Chen, S.-C.; Liaw, Y.-P. Cell-type specificity of lung cancer associated with low-dose soil heavy metal contamination in Taiwan: An ecological study. *BMC Public Health* **2013**, *13*, 330. [[CrossRef](#)] [[PubMed](#)]
20. R Core Team. *R: A Language and Environment for Statistical Computing*; R Foundation for Statistical Computing: Vienna, Austria, 2019.
21. *Outdoor Air Pollution*; IARC Working Group on the Evaluation of Carcinogenic Risks to Humans (Ed.) IARC Monographs on the Evaluation of Carcinogenic Risks to Humans; International Agency for Research on Cancer, World Health Organization: Lyon, France, 2016; ISBN 978-92-832-0147-2.
22. Guo, C.; Chan, T.-C.; Teng, Y.-C.; Lin, C.; Bo, Y.; Chang, L.; Lau, A.K.H.; Tam, T.; Wong, M.C.S.; Qian Lao, X. Long-term exposure to ambient fine particles and gastrointestinal cancer mortality in Taiwan: A cohort study. *Environ. Int.* **2020**, *138*, 105640. [[CrossRef](#)] [[PubMed](#)]
23. Iwai, K.; Mizuno, S.; Miyasaka, Y.; Mori, T. Correlation between suspended particles in the environmental air and causes of disease among inhabitants: Cross-sectional studies using the vital statistics and air pollution data in Japan. *Environ. Res.* **2005**, *99*, 106–117. [[CrossRef](#)] [[PubMed](#)]
24. White, A.J.; Bradshaw, P.T.; Hamra, G.B. Air Pollution and Breast Cancer: A Review. *Curr. Epidemiol. Rep.* **2018**, *5*, 92–100. [[CrossRef](#)] [[PubMed](#)]