



Valuations on Structures More General Than Fields [†]

Alessandro Linzi

Center for Information Technologies and Applied Mathematics, University of Nova Gorica,
5000 Nova Gorica, Slovenia; alessandro.linzi@ung.si

[†] Presented at the 1st International Online Conference on Mathematics and Applications, 1–15 May 2023;
Available online: <https://iocma2023.sciforum.net/>.

Abstract: Valuation theory is an important area of investigation in algebra, with applications in algebraic geometry and number theory. In 1957, M. Krasner introduced hyperfields, which are field-like objects with a multivalued addition, to describe some structures arising naturally from valued fields. We wish to discuss the possibility of generalising the notion of valuation to the multivalued setting and the potential that this higher point of view has in the understanding of classical valuation theory. We will see that a valuation on a field K is nothing but a homomorphism of hyperfields from K onto a special type of hyperfield, which we call a (generalised) tropical hyperfield.

Keywords: hyperfield; valuation; valued field; tropical hyperfield

1. Introduction

Valuation theory has its origins in the study of p -adic numbers' structure, arising as the completion of the field of rational numbers with respect to the metrics naturally induced by prime numbers, p (see [1]). The explicit arithmetic criteria available on complete fields for deciding whether a rational number is representable by a quadratic form constitute the main reason for the success of Hasse's local–global principle [2]. This principle represents one of the first successful applications of valuation theory in the realm of algebraic number theory after the initial efforts of Kürschák [3] and Ostrowski [4–6] in establishing its foundations.

Further work by Ostrowski [7] set the ground for the great development of valuation theory by relating it to Galois theory. However, it is thanks to the ideas illustrated by Krull in [8] that the techniques of valuation theory were able to show their effectiveness in more and more areas of mathematics, other than number theory.

The general definition of valuation on a field formulated by Krull (which is the same as the definition we adopt below) is fundamentally related to the concept of *place*. The latter is essential for the understanding of many problems in algebraic geometry as resolutions of singularities and local uniformization (for more details, see, e.g., [9] and references therein). Moreover, the breakthrough theorem of Ax–Kochen [10–12] and Ershov [13] have inspired many mathematicians to investigate the model theory of valued fields, which generated many interesting and deep mathematical ideas and constructions.

Even though valuation theory has recorded much success in its history, as often happens in mathematics, many relevant problems remain open (e.g., local uniformization in positive characteristics or the decidability of the elementary theory of a Laurent series field over a field of prime cardinality). The challenges that these and other problems offer push us to the limits of the existing theories, where we imagine new approaches with the guidance of the patterns that we are able to recognise and explain.

Let K be a field and let Γ be a linearly ordered Abelian group (always written additively). A surjective map

$$v : K \rightarrow \Gamma \cup \{\infty\}$$

is called a (*Krull*) *valuation on K* if it satisfies the following properties:

V1. $v(x) = \infty$ if and only if $x = 0$.



Citation: Linzi, A. Valuations on Structures More General Than Fields. *Comput. Sci. Math. Forum* **2023**, *7*, 2. <https://doi.org/10.3390/IOCMA2023-14402>

Academic Editor: Connelly Yang

Published: 28 April 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

- V2. $v(xy) = v(x) + v(y)$.
- V3. $v(x + y) \geq \min\{v(x), v(y)\}$.

for all $x, y \in K$. Here, ∞ is a symbol such that $\gamma + \infty = \infty + \gamma = \infty > \gamma$ for all $\gamma \in \Gamma$. If a valuation v on a field K is given, then (K, v) is usually called a *valued field*. We shall denote Γ by vK and call it the *value group of (K, v)* . The *value $v(x)$* of $x \in K$ will often be written as vx if there is no risk of confusion. For simplicity, in the sequel, all valued fields are assumed to be *non-trivially valued*, i.e., satisfying $vK \neq \{0\}$. Under this assumption, vK , and thus, K , are infinite.

If (K, v) is a valued field and $\gamma \geq 0$ is a nonnegative element of the value group vK , then it is well-known that

$$U_v^\gamma := \{x \in K \mid v(x - 1) > \gamma\}$$

is a subgroup of $K^\times := K \setminus \{0\}$ with respect to multiplication. It is called the group of *principal units of level γ* in (K, v) . Consider the corresponding quotient group:

$$K_\gamma^\times := K^\times / U_v^\gamma = \{xU_v^\gamma \mid x \in K^\times\}$$

and introduce the following notation:

$$xU_v^\gamma \boxplus yU_v^\gamma := \{zU_v^\gamma \mid z = x' + y' \text{ for some } x' \in xU_v^\gamma \text{ and } y' \in yU_v^\gamma\}.$$

Consider also $0U_v^\gamma = \{0\}$ and set

$$K_\gamma := K_\gamma^\times \cup \{0U_v^\gamma\}.$$

We will sometimes abuse notation and also denote the element $0U_v^\gamma$ of K_γ by 0. Similarly, we will sometimes denote by 1 the element $1U_v^\gamma$ of K_γ . Motivations for these choices will be clarified later. It is natural to extend \boxplus to a function

$$\boxplus : K_\gamma \times K_\gamma \rightarrow \mathcal{P}(K_\gamma),$$

where $\mathcal{P}(K_\gamma)$ denotes the power set of K_γ . Such a function is sometimes called a *(binary) multivalued operation* on the set K_γ . Let us extend it further to a binary operation on $\mathcal{P}(K_\gamma)$ as follows. For an $A, B \subseteq K_\gamma$ set

$$A \boxplus B := \bigcup_{(a,b) \in A \times B} a \boxplus b,$$

we have the understanding that an empty union is equal to the empty set. By direct inspection, it is not difficult to verify that the following properties hold:

1. $a \boxplus b \neq \emptyset$ for all $a, b \in K_\gamma$ (i.e., \boxplus is a *hyperoperation*).
2. $(a \boxplus b) \boxplus c = a \boxplus (b \boxplus c)$ are sets for all $a, b, c \in K_\gamma$.
3. $a \boxplus b = b \boxplus a$ are sets for all $a, b \in K_\gamma$.
4. For all $a \in K_\gamma$, there exists a unique $a^- \in K_\gamma$ such that $0 \in a \boxplus a^-$.
5. If $c \in a \boxplus b$, then $a \in c \boxplus b^-$ for all $a, b, c \in K_\gamma$.
6. $K_\gamma \setminus \{0\}$ is an Abelian group with respect to some operation \cdot with a neutral element 1 and $a \cdot 0 = 0 \cdot a = 0$ for all $a \in K_\gamma$.
7. $c \cdot (a \boxplus b) = c \cdot a \boxplus c \cdot b$ are sets for all $a, b, c \in K_\gamma$, where $c \cdot (a \boxplus b) = \{c \cdot d \mid d \in a \boxplus b\}$.

Clearly, the operation \cdot in properties 6 and 7 above is the multiplication induced from the multiplicative structure of the field K .

The construction that we just described was first studied by M. Krasner in the paper [14] (the article is included in Krasner’s collected works ([15], pp. 413–490), where, inspired by the above observations, he also gave the following general definition:

Definition 1. Any structure $(F, \boxplus, \cdot, 0, 1)$, where \boxplus is a binary multivalued operation on F and \cdot is a binary operation on F , satisfying properties 1–7 above is called a hyperfield.

The notion of a hyperfield thus generalises the notion of a field by allowing the additive operation to be multivalued. Any field K is a hyperfield with respect to the hyperoperation $x \boxplus y := \{x + y\}$. Conversely, any hyperfield F whose multivalued operation \boxplus satisfies $|a \boxplus b| = 1$ for all $a, b \in F$ can be naturally regarded as a field. In this short paper, we will discuss the possibility of defining valuations on hyperfields and discuss some new possibilities that the resulting theory offers.

2. Valued Hyperfields

We begin this section with the following result of Krasner, which further motivates us to fulfil our aim of finding a suitable notion of valuation for hyperfields.

Proposition 1 ([14]). Let (K, v) be a valued field and let $\gamma \geq 0$ be a nonnegative element of vK . Then, $v_x = 0$ for all $x \in U_v^\gamma$. Moreover, the map

$$v_\gamma : K_\gamma \rightarrow vK \cup \{\infty\}$$

$$xU_v^\gamma \mapsto vx$$

is surjective and satisfies

1. $v_\gamma a = \infty$ if and only if $a = 0$ for all $a \in K_\gamma$.
2. $v_\gamma(a \cdot b) = v_\gamma a + v_\gamma b$ for all $a, b \in K_\gamma$.
3. If $c \in a \boxplus b$, then $v_\gamma c \geq \min\{v_\gamma a, v_\gamma b\}$ for all $a, b, c \in K_\gamma$.
4. If $c \in a \boxplus b$, then $d \in a \boxplus b$ if and only if $v_\gamma e > \gamma + \min\{v_\gamma a, v_\gamma b\}$ for every $e \in c \boxplus d^{-}$ and for all $a, b \in K_\gamma$.
5. If $0 \notin a \boxplus b$, then $c, d \in a \boxplus b$ implies $v_\gamma c = v_\gamma d$ for all $a, b \in K_\gamma$.

Proof. Since in the ordered Abelian group vK the operation is compatible with the order and

$$v(1) = v(1 \cdot 1) = v(1) + v(1),$$

$v(1) = 0$ must hold in any valued field (K, v) ; we deduce that $v(x^{-1}) = -vx$ also holds for all $x \in K^\times$. A similar reasoning yields $v(-1) = 0$ and $v(-x) = vx$ for all $x \in K$. Moreover, if $vx < vy$, then $v(x + y) = vx$. Indeed, if the latter would not hold, then we would find

$$vx = v((x + y) - y) \geq \min\{v(x + y), v(-y)\} = \min\{v(x + y), vy\} > vx,$$

a contradiction. Now let $x \in U_v^\gamma$, i.e., $v(x - 1) > \gamma$. If $vx > 0 = v(1)$, then $v(x - 1) = v(1) = 0 \leq \gamma$, and if $vx < 0$, then $v(x - 1) = vx < 0 \leq \gamma$. Therefore, $vx = 0$ is necessary for $v(x - 1) > \gamma$ to hold.

As a consequence, the map v_γ is well-defined, and Properties 1 and 2 can be verified directly from Properties V1 and V2 of v . For Property 3, assume that

$$zU_v^\gamma \in xU_v^\gamma \boxplus yU_v^\gamma$$

holds for some $x, y, z \in K$. By the definition of \boxplus , $z = xs + yt$ for some $s, t \in U_v^\gamma$; hence,

$$vz = v(xs + yt) \geq \min\{v(xs), v(yt)\} = \min\{vx + vs, vy + vt\} = \min\{vx, vy\}$$

follows by the V3 property of v since $vs = vt = 0$. For property 4, note that for all $s, t \in U_v^\gamma$, we have

$$v(s - t) = v(s - 1 - t + 1) \geq \min\{v(s - 1), v(1 - t)\} > \gamma.$$

Now, take $x, y \in K$ and assume without loss of generality that $vx \leq vy$. By the definition of \boxplus , for $z, z' \in K$, we have that

$$zU_v^\gamma, z'U_v^\gamma \in xU_v^\gamma \boxplus yU_v^\gamma$$

if and only if $z = xs + yt$ and $z' = xs' + yt'$ for some $s, t, s', t' \in U_v^\gamma$. We deduce that

$$v(zs'' - z't'') = v(x(ss'' - s't'') + y(ts'' - t't'')) > \gamma + \min\{vx, vy\}$$

holds for all $s'', t'' \in U_v^\gamma$ since $ss'', s't'', ts'', t't'' \in U_v^\gamma$. Conversely, if $v(z - z') > \gamma + \min\{vx, vy\} = \gamma + vx$ holds for some $z, z' \in K$ and

$$zU_v^\gamma \in xU_v^\gamma \boxplus yU_v^\gamma,$$

i.e., $z = xs + yt$ for some $s, t \in U_v^\gamma$, then

$$v((z - z')x^{-1}) = v(z - z') - vx > \gamma.$$

Therefore, for $s' := 1 + (z - z')x^{-1} \in U_v^\gamma$, we obtain that

$$z' = (1 + s' - s)x + yt.$$

Since $v(s' - s) > \gamma$ as we have already shown above, we deduce that

$$z'U_v^\gamma \in xU_v^\gamma \boxplus yU_v^\gamma.$$

It remains to be shown that Property 5 holds as well. For this, we take $x, y \in K$ and assume that

$$v(xt + ys) < v(xt' + ys')$$

for some $s, t, s', t' \in U_v^\gamma$. By Property 4, we obtain that

$$v(0 - (xs + yt)) = v(xs + yt) = v(xt + ys - (xt' + ys')) > \gamma + \min\{vx, vy\}.$$

By another application of Property 4, we obtain that $0 \in xU_v^\gamma \boxplus yU_v^\gamma$. \square

The similarity between Properties 1, 2 and 3 and the Properties V1, V2 and V3 motivate the following definition.

Definition 2. *If F is a hyperfield, Γ is an ordered Abelian group, and $v : F \rightarrow \Gamma \cup \{\infty\}$ a surjective map satisfying properties 1, 2, and 3 of Proposition 1 above, then we call v a valuation and (F, v) a valued hyperfield. We retain our notation and terminology as explained in the introduction for valuations on fields.*

Krasner also studied valued hyperfields, but his definition was more restrictive than the above one. He postulated that valuations on hyperfields additionally satisfy Property 4 (where the free variable γ is bounded existentially and required to be nonnegative in the value group) and Property 5. Even though Krasner’s definition captures some relatively well-behaved structures (cf., e.g., [16–19]), our choice can be motivated by the observation of many natural examples that do not fit Krasner’s definition. Among these examples, we present an important one below.

Example 1 (Generalised tropical hyperfield). *Let Γ be an ordered Abelian group and let ∞ be a symbol such that $\gamma + \infty = \infty + \gamma = \infty > \gamma$ for all $\gamma \in \Gamma$. For $\gamma, \delta \in \Gamma \cup \{\infty\}$ such that $\gamma \leq \delta$, let us denote by $[\gamma, \delta]$ the set consisting of all $\varepsilon \in \Gamma \cup \{\infty\}$ such that $\gamma \leq \varepsilon \leq \delta$. We consider the multivalued operation \boxplus defined on $\mathcal{T}(\Gamma) := \Gamma \cup \{\infty\}$ as follows:*

$$\gamma \boxplus \infty = \infty \boxplus \gamma = \{\gamma\} \quad (\gamma \in \mathcal{T}(\Gamma))$$

and

$$\gamma \boxplus \delta := \begin{cases} \{\min\{\gamma, \delta\}\} & \text{if } \gamma \neq \delta, \\ [\gamma, \infty] & \text{otherwise.} \end{cases} \quad (\gamma, \delta \in \Gamma)$$

It is not difficult to check that $\mathcal{T}(\Gamma)$ is a hyperfield, where the multiplication is given by the operation $+$ of Γ . The hyperfield $\mathcal{T}(\mathbb{R}, +, 0, >)$, where $<$ denotes the standard order of the real numbers, is known as the tropical hyperfield (see, e.g., Section 1 of [20]). We call the hyperfields of the form $\mathcal{T}(\Gamma)$ generalised tropical hyperfields.

The identity map on $\mathcal{T}(\Gamma)$ is a valuation on $\mathcal{T}(\Gamma)$ as it (almost trivially) satisfies Properties 1–3 of Proposition 1. Nevertheless, Property 4 does not hold in this case as $0 \in [0, \infty] = 0 \boxplus 0$, but for all $\delta > 0$, we have that $0 \boxplus \delta = \{0\}$, and $0 > \gamma + 0$ does not hold for any $\gamma \geq 0$.

3. Homomorphisms of Hyperfields

Similar to other algebraic structures, hyperfields can be arranged in a category. The standard choice for arrows in the category of hyperfields are the homomorphisms that are defined as follows:

Definition 3. Let F_1 and F_2 be hyperfields. A map $\sigma : F_1 \rightarrow F_2$ is called a homomorphism of hyperfields if the following properties hold:

- H1. $\sigma(a) = 0_2$ if and only if $a = 0_1$ for all $a \in F_1$.
- H2. $\sigma : F_1^\times \rightarrow F_2^\times$ is a homomorphism of groups.
- H3. $\sigma(a \boxplus_1 b) \subseteq \sigma(a) \boxplus_2 \sigma(b)$ for all $a, b \in F_1$.

The next observation provides an alternative definition of valuation on hyperfields and, in particular, on fields (cf. Example 1.8(2) of [20]). We leave the straightforward proof to the reader.

Proposition 2. Let F be a hyperfield and let Γ be an ordered Abelian group. Then, (F, v) is a valued hyperfield with $vF = \Gamma$ if and only if the map $v : F \rightarrow \mathcal{T}(\Gamma)$ is a surjective homomorphism of hyperfields.

Thus, if we think of valuations as surjective homomorphisms of hyperfields, then their properties can be reflected directly into the additive structure of the target hyperfield.

In [21], generalised tropical hyperfields are characterised as hyperfields $(T, \boxplus, \cdot, 0, 1)$ satisfying the following properties:

- (T1) For all $a, b \in T$, if $0 \notin a \boxplus b$, then $|a \boxplus b| = 1$, i.e., T is stringent.
- (T2) $0, 1 \in 1 \boxplus 1$.

It is interesting to note that many generalisations of the notion of valuation that appeared in the literature can analogously be understood as homomorphisms of hyperfields and, under this interpretation, correspond to less restrictive properties for their target hyperfield than (T1) and (T2). Below, we briefly present an example.

Example 2. Let L be a lattice that is also a group with respect to a compatible operation $+$ (i.e., L is an ℓ -group). Let ∞ be a new symbol such that $l + \infty = \infty + l = \infty > l$ for all $l \in L$ and define on $L \cup \{\infty\}$ the following multivalued operation:

$$l \boxplus \infty = \infty \boxplus l = \{l\} \quad (l \in L \cup \{\infty\})$$

and

$$n \boxplus m := \{l \in L \mid n \wedge l = l \wedge m = n \wedge m\} \quad (n, m \in L).$$

By some results of Nakano (see Theorem 1 of [22]), it follows that the resulting structure $\mathcal{N}(L)$ is a hyperfield (notice that since any ℓ -group is distributive, L is modular). We have that $\mathcal{N}(L)$ is stringent if and only if the order of L is linear, in which case L is an ordered Abelian group and $\mathcal{N}(L) = \mathcal{T}(\Gamma)$, as the reader can easily check. Now, it is not difficult to see that v is a lattice-

valuation of a field K onto L (in the sense of, e.g., [23]) if and only if v is a surjective homomorphism of the hyperfields $K \rightarrow \mathcal{N}(L)$. Thus, in this case, it is the property (T1) of the target hyperfield that is relaxed, while (T2) holds in $\mathcal{N}(L)$.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The author declares no conflict of interest.

References

- Hensel, K. *Theorie der algebraischen Zahlen*; B. G. Teubner: Leipzig, Germany, 1908.
- Hasse, H. Zur Theorie der quadratischen Formen, insbesondere ihrer Darstellbarkeitseigenschaften im Bereich der rationalen Zahlen und ihrer Einteilung in Geschlechter. Ph.D. Thesis, Philipps-Universität zu Marburg, Marburg, Germany, 1921.
- Kürschák, J. Über Limesbildung und allgemeine Körpertheorie. *J. Reine Angew. Math.* **1913**, *142*, 211–253. [[CrossRef](#)]
- Ostrowski, A. Über einige Fragen der allgemeinen Körpertheorie. *J. Reine Angew. Math.* **1913**, *143*, 255–284. [[CrossRef](#)]
- Ostrowski, A. Über einige Lösungen der Funktionalgleichung $\psi(x) \cdot \psi(x) = \psi(xy)$. *Acta Math.* **1916**, *41*, 271–284. [[CrossRef](#)]
- Ostrowski, A. Über sogenannte perfekte Körper. *J. Reine Angew. Math.* **1917**, *147*, 191–204. [[CrossRef](#)]
- Ostrowski, A. Untersuchungen zur arithmetischen Theorie der Körper. *Math. Z.* **1935**, *39*, 321–404. [[CrossRef](#)]
- Krull, W. Allgemeine Bewertungstheorie. *J. Reine Angew. Math.* **1932**, *167*, 160–196. [[CrossRef](#)]
- Vaquíe, M. Valuations and Local Uniformization. In *Singularity Theory and Its Applications*; Mathematical Society of Japan: Tokyo, Japan, 2006; Volume 43, pp. 477–527. [[CrossRef](#)]
- Ax, J.; Kochen, S. Diophantine problems over local fields. I. *Am. J. Math.* **1965**, *87*, 605–630. [[CrossRef](#)]
- Ax, J.; Kochen, S. Diophantine problems over local fields. II. A complete set of axioms for p -adic number theory. *Am. J. Math.* **1965**, *87*, 631–648. [[CrossRef](#)]
- Ax, J.; Kochen, S. Diophantine problems over local fields. III. Decidable fields. *Ann. Math.* **1966**, *83*, 437–456. [[CrossRef](#)]
- Eršov, J.L. On the elementary theory of maximal normed fields. *Dokl. Akad. Nauk SSSR* **1965**, *165*, 21–23.
- Krasner, M. Approximation des corps valués complets de caractéristique $p \neq 0$ par ceux de caractéristique 0. In *Colloque d'algèbre supérieure, tenu à Bruxelles du 19 au 22 décembre 1956*; Centre Belge de Recherches Mathématiques, Établissements Ceuterick, Louvain; Librairie Gauthier-Villars: Paris, France, 1957; pp. 129–206.
- Bélaïr, L.; Poizat, B. *Oeuvres Choisies de Marc Krasner*; Editions L'Harmattan: Paris, France, 2019.
- Mittas, J. Sur les hyperanneaux et les hypercorps. *Math. Balkanica* **1973**, *3*, 368–382.
- Tolliver, J. An equivalence between two approaches to limits of local fields. *J. Number Theory* **2016**, *166*, 473–492. [[CrossRef](#)]
- Lee, J. Hyperfields, truncated DVRs, and valued fields. *J. Number Theory* **2020**, *212*, 40–71. [[CrossRef](#)]
- Linzi, A.; Touchard, P. On the hyperfields associated to valued fields. *arXiv* **2022**, arXiv:2211.05082.
- Baker, M.; Lorscheid, O. Descartes' rule of signs, Newton polygons, and polynomials over hyperfields. *J. Algebra* **2021**, *569*, 416–441. [[CrossRef](#)]
- Linzi, A. Notes on valuation theory for Krasner hyperfields. *arXiv* **2023**, arXiv:2301.08639.
- Nakano, T. Rings and partly ordered systems. *Math. Z.* **1967**, *99*, 355–376. [[CrossRef](#)]
- Fuchs, L.; Salce, L. Modules over non-Noetherian domains. In *Mathematical Surveys and Monographs*; American Mathematical Society: Providence, RI, USA, 2001; Volume 84, 613p. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.