



The Odd Beta Prime-G Family of Probability Distributions: Properties and Applications to Engineering and Environmental Data [†]

Ahmad Abubakar Suleiman ^{1,2,*} , Hanita Daud ^{1,*}, Mahmud Othman ³, Aliyu Ismail Ishaq ⁴ , Rachmah Indawati ⁵, Mohd Lazim Abdullah ⁶ and Abdullah Husin ³

¹ Fundamental and Applied Sciences Department, Universiti Teknologi PETRONAS, Seri Iskandar 32610, Malaysia

² Department of Statistics, Aliko Dangote University of Science and Technology, Wudil 713281, Nigeria

³ Department of Information System, Universitas Islam Indragiri, Tembilahan 29281, Indonesia; mahmod.othman@unisi.ac.id (M.O.); abdialam@yahoo.com (A.H.)

⁴ Department of Statistics, Ahmadu Bello University, Zaria 810107, Nigeria; aiishaq@abu.edu.ng

⁵ Faculty of Public Health, Airlangga University, Surabaya 60115, Indonesia; rachmah.indawati@fkm.unair.ac.id

⁶ Management Science Research Group, Faculty of Ocean Engineering Technology and Informatics, Universiti Malaysia Terengganu, Kuala Nerus 21030, Malaysia; lazim_m@umt.edu.my

* Correspondence: ahmad_22000579@utp.edu.my (A.A.S.); hanita_daud@utp.edu.my (H.D.)

[†] Presented at the 1st International Online Conference on Mathematics and Applications, 1–15 May 2023;

Available online: <https://iocma2023.sciforum.net/>.

Abstract: In this work, a novel generalized family of distributions called the odd beta prime is introduced. The linear representations of the proposed family are obtained. The expressions for the moments, the moment-generating function, and entropy are derived. A three-parameter special sub-model of the proposed family called the odd beta prime exponential distribution is proposed. Finally, two real data sets are used to illustrate the usefulness and flexibility of the proposed distribution.

Keywords: odd beta prime generalized family; exponential distribution; T-X transformer; moments; moment-generating function; entropy



Citation: Suleiman, A.A.; Daud, H.; Othman, M.; Ishaq, A.I.; Indawati, R.; Abdullah, M.L.; Husin, A. The Odd Beta Prime-G Family of Probability Distributions: Properties and Applications to Engineering and Environmental Data. *Comput. Sci. Math. Forum* **2023**, *7*, 20. <https://doi.org/10.3390/IOCMA2023-14429>

Academic Editor: Antonio Di Crescenzo

Published: 28 April 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

A popular research field is the construction of novel approaches for extending the existing distributions. The two interesting approaches to expanding a probability distribution are the T-X technique proposed by [1] and modified by [2]. The cumulative distribution function (cdf) for the generalizing family of distributions using this approach is given as:

$$F(x) = \int_a^{W(G(x))} v(m) dm = V\{W(G(x))\}, \quad (1)$$

where $v(m)$ is the pdf of the random variable $M \in [a, b]$, such that $-\infty \leq a < b \leq \infty$, and $W(G(x))$ is a link function of any cdf of continuous distributions that take different forms. If we consider the odd function form, $W(G(x)) = \frac{G(x)}{1-G(x)}$, then the cdf of the T-X class will be:

$$F(x) = \int_0^{\frac{G(x)}{1-G(x)}} v(m) dm = V\left\{\frac{G(x)}{1-G(x)}\right\}. \quad (2)$$

Many authors constructed extended generalized families by using the T-X approach. For example, see beta-G [3], Kw-G type-1 [4], gamma-X [5], exponentiated T-X [1], Weibull-G [6], generalized odd Lindley-G [7], and Maxwell–Weibull [8].

In this study, we consider the odd function form, $W(G(x)) = \frac{G(x)}{1-G(x)}$. Additionally, we considered the beta prime distribution for $v(m) = \frac{1}{B(c,d)} \frac{x^{c-1}}{(1+x)^{c+d}}$, $x > 0$.

Therefore, we now define the odd beta prime-G family with cdf given as:

$$F(x) = \frac{B_{\frac{G(x;\varepsilon)}{1-G(x;\varepsilon)}}(c,d)}{B(c,d)}, \quad x \geq 0 \quad (3)$$

The probability distribution function (pdf) of odd beta prime-G family is:

$$f(x) = \frac{g(x;\varepsilon)}{B(c,d)[1-G(x;\varepsilon)]^2} \frac{\left(\frac{G(x;\varepsilon)}{1-G(x;\varepsilon)}\right)^{c-1}}{\left[1 + \left(\frac{G(x;\varepsilon)}{1-G(x;\varepsilon)}\right)\right]^{c+d}}; \quad x \in \mathbb{R}, \quad (4)$$

where $G(x;\varepsilon)$ is a cdf of a baseline distribution with parameter ε , $g(x;\varepsilon)$ is the pdf of the baseline distribution, and $c > 0$ and $d > 0$ are the shape parameters.

Here, we are motivated to propose a new flexible family of distribution called the odd beta prime generalized (OBP-G) family, which provides greater accuracy and flexibility in fitting real-life data.

This article unfolds as follows. In Section 2, linear representations of the proposed family are derived. Some statistical properties are studied and obtained in Section 3. A special sub-model of the proposed family is introduced in Section 4. In Section 5, the performance of the proposed distribution is illustrated via two applications to real data sets. Finally, Section 6 concludes the article.

2. Linear Representations

This section presents important linear representations of the OBP-G family density function defined in (4).

Let us consider the generalized Binomial expansion as follows:

$$(1+m)^{-\eta} = \sum_{i=0}^{\infty} \binom{-\eta}{i} M^i = \sum_{i=0}^{\infty} (-1)^i \binom{-\eta+i-1}{i} M^i. \quad (5)$$

Applying (5) into (4), we obtain:

$$\begin{aligned} f(x) &= \frac{g(x;\varepsilon)}{B(c,d)[1-G(x;\varepsilon)]^2} \left(\frac{G(x;\varepsilon)}{1-G(x;\varepsilon)}\right)^{c-1} \sum_{i=0}^{\infty} (-1)^i \binom{c+d+i-1}{i} \left(\frac{G(x;\varepsilon)}{1-G(x;\varepsilon)}\right)^i, \\ &= \frac{g(x;\varepsilon)}{B(c,d)} \sum_{i=0}^{\infty} (-1)^i \binom{c+d+i-1}{i} \frac{(G(x;\varepsilon))^{c+i-1}}{(1-G(x;\varepsilon))^{c+i+1}}. \end{aligned} \quad (6)$$

Using the generalized Binomial expansion for $|z| < 1$ yields:

$$(1-m)^{-n} = \sum_{j=0}^{\infty} \frac{\Gamma(n+j)}{j!\Gamma n} z^j. \quad (7)$$

Substituting (7) into (6), we obtain:

$$\begin{aligned} f(x) &= \frac{g(x;\varepsilon)}{B(c,d)} \sum_{i=0}^{\infty} (-1)^i \binom{c+d+i-1}{i} (G(x;\varepsilon))^{c+i-1} \sum_{j=0}^{\infty} \frac{\Gamma(c+i+j+1)}{j!\Gamma(c+i+1)} (G(x;\varepsilon))^j, \\ &= \frac{g(x;\varepsilon)}{B(c,d)} \sum_{i,j=0}^{\infty} \Psi_{i,j} (G(x;\varepsilon))^{c+i+j-1}, \end{aligned} \quad (8)$$

where $\Psi_{i,j} = (-1)^i \binom{c+d+i-1}{i} \frac{\Gamma(c+i+j+1)}{j!\Gamma(c+i+1)}$.

3. Statistical Properties

This section provides some statistical properties of the OBP-G family of distributions, such as moments, the moment-generating function, and entropy.

3.1. Moments

Suppose that X follows the OBP-G family, the r th moments of X is obtained as:

$$E(X^r) = \int_{-\infty}^{\infty} x^r f(x) dx, \quad (9)$$

where $f(x)$ is defined in Equation (8).

Substituting (8) into (9) gives the moments of the OBP-G as:

$$E(X^r) = \frac{1}{B(c, d)} \sum_{i,j=0}^{\infty} \Psi_{i,j} \int_{-\infty}^{\infty} x^r g(x; \varepsilon) (G(x; \varepsilon))^{c+i+j-1} dx. \quad (10)$$

3.2. Moment-Generating Function

Assuming that a random variable X follows the OBP-G family, the moment-generating function of X is given as:

$$M_X(t) = E(e^{tx}) = \int_{-\infty}^{\infty} e^{tx} f(x) dx, \quad (11)$$

Inserting (8) into (11), we have:

$$M_X(t) = \frac{1}{B(c, d)} \sum_{i,j=0}^{\infty} \Psi_{i,j} \int_{-\infty}^{\infty} e^{tx} g(x; \varepsilon) (G(x; \varepsilon))^{c+i+j-1} dx. \quad (12)$$

This can be expressed as:

$$M_X(t) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} E(X^k). \quad (13)$$

3.3. Entropy

Assuming that X is a random variable that follows the OBP-G family, the Rényi entropy [9] of X is expressed as:

$$R_{\alpha}(x) = \frac{1}{1-\alpha} \log \left[\int_{-\infty}^{\infty} f^{\alpha}(x) dx \right]; \quad \alpha > 0, \alpha \neq 1, x \in \mathbb{R}. \quad (14)$$

The integrand $f^{\alpha}(x)$ can be obtained as:

$$f^{\alpha}(x) = [f(x)]^{\alpha} = \left\{ \frac{g(x; \varepsilon)}{B(c, d)[1 - G(x; \varepsilon)]^2} \frac{\left(\frac{G(x; \varepsilon)}{1 - G(x; \varepsilon)} \right)^{c-1}}{\left[1 + \left(\frac{G(x; \varepsilon)}{1 - G(x; \varepsilon)} \right) \right]^{c+d}} \right\}^{\alpha}. \quad (15)$$

Therefore, (15) can be rewritten as:

$$f^{\alpha}(x) = \frac{g^{\alpha}(x; \varepsilon)}{B^{\alpha}(c, d)[1 - G(x; \varepsilon)]^{2\alpha}} \frac{\left(\frac{G(x; \varepsilon)}{1 - G(x; \varepsilon)} \right)^{\alpha(c-1)}}{\left[1 + \left(\frac{G(x; \varepsilon)}{1 - G(x; \varepsilon)} \right) \right]^{\alpha(c+d)}}. \quad (16)$$

Substituting (5) into (16), we have:

$$f^{\alpha}(x) = \frac{g^{\alpha}(x; \varepsilon)}{B^{\alpha}(c, d)[1 - G(x; \varepsilon)]^{2\alpha}} \left(\frac{G(x; \varepsilon)}{1 - G(x; \varepsilon)} \right)^{\alpha(c-1)} \sum_{l=0}^{\infty} (-1)^l \binom{\alpha(c+d)+l-1}{l} \left(\frac{G(x; \varepsilon)}{1 - G(x; \varepsilon)} \right)^l. \quad (17)$$

4. The Odd Beta Prime Exponential Distribution

This section develops a new probability distribution referred to as the odd beta prime exponential (OBPE) distribution as a sub-model of the proposed family.

Let X be a random variable with exponential distribution; the cdf and pdf are, respectively, as follows:

$$G(x) = 1 - e^{-bx}, \quad x \geq 0, \quad (18)$$

$$g(x) = be^{-bx}, \quad x \geq 0, \quad (19)$$

where $b > 0$ is the rate parameter.

Hence, the cdf and pdf of the OBPE distribution can be obtained by inserting (18) and (19) into (3) and (4), respectively, as follows:

$$F(x) = \frac{B_{\frac{1-e^{-bx}}{e^{-bx}}}(c, d)}{B(c, d)}, \quad x \geq 0. \quad (20)$$

$$f(x) = \frac{b(1 - e^{-bx})^{c-1}}{B(c, d)e^{-bcx} \left[1 + \left(\frac{1-e^{-bx}}{e^{-bx}} \right) \right]^{c+d}}, \quad x \geq 0. \quad (21)$$

5. Applications

In this section, we analyze two real data sets involving engineering and environment to evaluate the applicability of the OBPE distribution.

5.1. The Airborne Communications Transceiver Data

These engineering data were discussed in [10], and they represent the repair times of 46 failures (in hours) of an airborne communications transceiver.

Here, we will compare the fits of the OBPE with the gamma-exponentiated exponential (GEE) in [11] and the beta-exponential (BE) in [12].

We considered the following criteria to compare these distributions: the values of the negative log-likelihood ($-\hat{\ell}$), Akaike information criteria (AIC), Bayesian information criteria (BIC), Cramer-von Mises (CM), and Anderson-Darling (AD). The smaller the values of these statistics, the better the fit to the data [13,14].


The maximum likelihood estimates (MLEs), standard errors (SEs), $-\hat{\ell}$, AIC, BIC, CM, and AD statistics for the OBPE, GEE, and BE are presented in Table 1. From the results in Table 1, it is clear that the OBPE distribution provides a better fit for the data, having smallest values of $-\hat{\ell}$, AIC, BIC, CM, and AD, and could be selected as a more appropriate model than other models. Figure 1 depicts the estimated pdfs and cdfs of the fitted distributions. It is clear from these plots that the OBPE describes the data in a more appropriate way than other competing models.

5.2. Exceedances of Wheaton River Flood Data

These environmental data were analyzed by [15], and they represent the exceedances of flood peaks (in m^3/s) of the Wheaton River near Carcross in Yukon Territory, Canada. The data consist of 72 exceedances for the years 1958–1984, rounded to one decimal place.

Table 1. MLEs with corresponding SEs (in parentheses), and some statistical measures of competing models for the airborne communications transceiver data.


Distribution	MLE and SE in ()			$-\hat{\ell}$	AIC	BIC	CM	AD
OBPE	$c = 0.6583$ (0.1625)	$d = 1.1018$ (0.1149)	$b = 1.4863$ (4.5831)	100.0163	204.032	207.689	0.054	0.337
GEE	$\alpha = 0.9323$ (0.1701)	$\theta = 0.2585$ (0.0615)	$\lambda = 0.3685$ (0.7650)	104.9309	213.861	217.519	0.175	1.103
BE	$a = 2.6732$ (0.4920)	$b = 2.0190$ (0.2613)	$\lambda = 1.1455$ (0.4016)	128.48	260.960	264.617	0.409	3.007

**Figure 1.** The estimated pdfs of the OBPE and other competing models for the airborne communications transceiver data.

Additionally, it is clear from the results in Table 2 and the illustrations in Figure 2 that the OBPE provides a better fit for these data than other competing fitted models.

Table 2. MLEs with corresponding SEs (in parentheses), and some statistical measures of competing models for the exceedances of Wheaton River flood data.

Distribution	MLE and SE in ()			$-\hat{\ell}$	AIC	BIC	CM	AD
OBPE	$c = 1.2126$ (0.1166)	$d = 6.7584$ (1.1659)	$b = 1.6077$ (0.3743)	257.839	519.6782	524.231	0.232	1.472
GEE	$\alpha = 10.6378$ (1.3609)	$\theta = 6.6058$ (0.6475)	$\lambda = 2.9650$ (0.4475)	279.958	563.9169	568.470	0.330	2.478
BE	$a = 12.2041$ (1.4391)	$b = 12.2115$ (1.0176)	$\lambda = 5.1753$ (0.2352)	282.334	568.669	573.223	0.469	2.891

**Figure 2.** The estimated pdfs of the OBPE and other competing models for the exceedances of Wheaton River flood data.

6. Conclusions

A new family of life distributions called the odd beta prime-G family was introduced. Some statistical properties of the new family, including moments, the moment-generating function, and entropy, were derived. A special sub-model of the newly proposed family called the odd beta prime exponential distribution was developed, and two real applications were analyzed to demonstrate the flexibility of the new distribution. Empirically, it was

proven that the proposed model can provide a better fit for modeling data than the other competing life distributions.

Author Contributions: Conceptualization, A.A.S., M.O. and H.D.; methodology, A.A.S., M.O., A.I.I. and M.L.A.; software, A.A.S. and A.I.I.; validation, A.A.S., M.O., R.I., H.D., R.I. and A.H.; supervision, M.O. and H.D.; formal analysis, A.A.S., M.O., A.I.I. and M.L.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Yayasan Universiti Teknologi PETRONAS (YUTP) with 492 cost center 015LC0-401.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors would like to thank the Universiti Teknologi PETRONAS for providing support for this project.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Alzaatreh, A.; Lee, C.; Famoye, F. A new method for generating families of continuous distributions. *Metron* **2013**, *71*, 63–79. [\[CrossRef\]](#)
2. Aljarrah, M.A.; Lee, C.; Famoye, F. On generating TX family of distributions using quantile functions. *J. Stat. Distrib. Appl.* **2014**, *1*, 1–17. [\[CrossRef\]](#)
3. Eugene, N.; Lee, C.; Famoye, F. Beta-normal distribution and its applications. *Commun. Stat.-Theory Methods* **2002**, *31*, 497–512. [\[CrossRef\]](#)
4. Cordeiro, G.M.; de Castro, M. A new family of generalized distributions. *J. Stat. Comput. Simul.* **2011**, *81*, 883–898. [\[CrossRef\]](#)
5. Torabi, H.; Montazeri, N.H. The logistic-uniform distribution and its applications. *Commun. Stat.-Simul. Comput.* **2014**, *43*, 2551–2569. [\[CrossRef\]](#)
6. Bourguignon, M.; Silva, R.B.; Cordeiro, G.M. The Weibull-G family of probability distributions. *J. Data Sci.* **2014**, *12*, 53–68. [\[CrossRef\]](#)
7. Afify, A.Z.; Cordeiro, G.M.; Maed, M.E.; Alizadeh, M.; Al-Mofleh, H.; Nofal, Z.M. The generalized odd Lindley-G family: Properties and applications. *An. Acad. Bras. Ciências* **2019**, *91*, e20180040. [\[CrossRef\]](#) [\[PubMed\]](#)
8. Ishaq, A.I.; Abiodun, A.A. The Maxwell–Weibull distribution in modeling lifetime datasets. *Ann. Data Sci.* **2020**, *7*, 639–662. [\[CrossRef\]](#)
9. Rényi, A. On measures of entropy and information. In Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, Berkeley, CA, USA, 20 June–30 July 1961; Volume 1, pp. 547–561.
10. Refaie, M.K. Burr X exponentiated exponential distribution. *J. Stat. Appl.* **2018**, *1*, 71–88.
11. Ristić, M.M.; Balakrishnan, N. The gamma-exponentiated exponential distribution. *J. Stat. Comput. Simul.* **2012**, *82*, 1191–1206. [\[CrossRef\]](#)
12. Nadarajah, S.; Kotz, S. The beta exponential distribution. *Reliab. Eng. Syst. Saf.* **2006**, *91*, 689–697. [\[CrossRef\]](#)
13. Singh, V.V.; Suleman, A.A.; Ibrahim, A.; Abdullahi, U.A.; Suleiman, S.A. Assessment of probability distributions of groundwater quality data in Gwale area, north-western Nigeria. *Ann. Optim. Theory Pract.* **2020**, *3*, 37–46.
14. Abdullahi, U.A.; Suleiman, A.A.; Ishaq, A.I.; Usman, A.; Suleiman, A. The Maxwell–Exponential Distribution: Theory and Application to Lifetime Data. *J. Stat. Model. Anal.* **2021**, *3*, 65–80. [\[CrossRef\]](#)
15. Choulakian, V.; Stephens, M.A. Goodness-of-fit tests for the generalized Pareto distribution. *Technometrics* **2001**, *43*, 478–484. [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.