

Supplementary materials

Thermodormancy and germination response to temperature of *Pyrus ussuriensis* seeds

Zhuolin Liu^{1,2}, Mingyue Li^{1,2}, Meiru Zhu^{1,2}, Rosana López³, Roberto L Salomón³, Peng Zhang^{1,2,*}

¹ College of Forestry, Northeast Forestry University, Harbin 150040, China; liuzhuolin2023@163.com (Z. L.); mingyue09222022@163.com (M. L.); zhumeiru2022@163.com (M. Z.); zhangpeng@nefu.edu.cn (P. Z.)

² Key Laboratory of Sustainable Forest Ecosystem Management—Ministry of Education, Northeast Forestry University, Harbin 150040, China.; liuzhuolin2023@163.com (Z. L.); mingyue09222022@163.com (M. L.); zhumeiru2022@163.com (M. Z.); zhangpeng@nefu.edu.cn (P. Z.)

³ Universidad Politécnica de Madrid, Department of Natural Systems and Resources, Research group FORESCENT, Ciudad Universitaria s/n, 28040 Madrid, Spain; rosana.lopez@upm.es (R. L.); roberto.salomon@upm.es (R. S.)

* Correspondence: zhangpeng@nefu.edu.cn

Figure S1. Photographs of the germination behaviour of dormant, non-dormant and thermodormant *P. ussuriensis* seeds in gibberellic acid at low (A) and high (B) concentrations and in abscisic acid at low (C) and high (D) concentrations.




Table S1. Degrees of freedom and p values of the two-way analysis of variance of the effect of dormancy state, solution treatment, and its interaction on the germination percentage of isolated embryos of *P. ussuriensis* seeds.

Source of Variation	df	p value
Dormancy state	2	< 0.001
Solution	5	< 0.001
Dormancy state:Solution	10	> 0.05

Table S2. Degrees of freedom and p values of the two-way analysis of variance of the effect of dormancy state, solution treatment, and its interaction treatment on the germination percentage of cabbage seeds.

Source of Variation	df	p value
Dormancy state	2	< 0.01
Solution	3	< 0.001
Dormancy state:Solution	6	> 0.05