Supplementary information

Growth and Nutritional Enhancement of Lisianthus (*Eustoma grandiflorum* (Raf.) Shinn.) via Dual AMF Inoculation under Phosphorus Regimes

Morteza Sheikh-Assadi [1], Mohammadreza Taheri [1], Azizollah Khandan-Mirkohi [1,*], Mesbah Babalar [1], Mahdi Khansefid [1], Mohammad Omidi [1] and Dionisios Gasparatos [2,*]

[1] Department of Horticultural Science, Faculty of Agricultural Science and Engineering, University of Tehran,Karaj 31587-77871, Iran; m.asadi66@ut.ac.ir (M.S.-A.); mrtaheri@ut.ac.ir (M.R.T.); mbabalar@ut.ac.ir (M.B.); m.omidighale@ut.ac.ir (M.O); mkhansefid@ut.ac.ir (M.K.)

[2] Laboratory of Soil Science and Agricultural Chemistry, Department of Natural Resources Management & Agricultural Engineering, School of Environment & Agricultural Engineering, Agricultural University of Athens, 11855 Athens, Greece

* Correspondence: khandan.mirkohi@ut.ac.ir (A.K.-M.); gasparatos@aua.gr (D.G)




Figure S1. The relation between shoot phosphorus concentration and days to flowering. *Outlier, not included in the regression.




Figure S2. Heat map of correlation among vegetative traits, ornamental features, and content of elements in Lisianthus plants. Shoot fresh weight (SFW), Shoot dry weight (SDW), Stem length (SL), Root dry weight (RDW), Days to flowering (DTF), Leaf area (LA), Root Fresh weight (RFW), Root length (RL), Number of stem per plant (NSP), Flower diameter (FD), Total flower fresh weight (TFFW), Flower fresh weight (FFW), Vase life (VL), Shoot P concentration (SPC), Shoot N concentration (SNC), Shoot Zn concentration (SZnC), Shoot Fe concentration (SFeC), Shoot K concentration (SKC), Shoot Ca concentration (SCaC), Root P concentration (RPC), Root K concentration (RKC), Root Ca concentration (RCaC), Root N concentration (RNC), Root Zn concentration (RZnC), Root Fe concentration (RFeC).