

Supporting Information

Detection of Acetaminophen and Its Glucuronide in Fingerprint by SALDI Mass Spectrometry Using Zeolite and Study of Time-Dependent Changes in Detected Ion Amount

Toshiki Horikoshi ¹, Chihiro Kitaoka ², Yosuke Fujii ³, Takashi Asano ⁴, Jiawei Xu ^{1,5} and Tatsuya Fujino ^{1,5*}

¹ Department of Applied Chemistry, Graduate School of Science and Engineering, Toyo University, 2100 Kujirai, Kawagoe 350-8585, Japan; s36d02100226@toyo.jp (T.H.); xu@toyo.jp (J.X.)

² Criminal Investigation Laboratory, Metropolitan Police Department, 2-1-1 Kasumigaseki, Chiyoda-ku, Tokyo 100-8929, Japan; kitaoka-chihiro@outlook.jp

³ Department of Chemistry, Graduate School of Science and Engineering, Tokyo Metropolitan University, 1-1 Minami-Osawa, Hachioji, Tokyo 192-0379, Japan; yosuke.fujii.221@gmail.com

⁴ Forensic Science Laboratory, Metropolitan Police Department, 2-1-1 Kasumigaseki, Chiyoda-ku, Tokyo 100-8929, Japan; asanofam@dc.mbn.or.jp

⁵ Bio-Nano Electronics Research Centre, Toyo University, 2100 Kujirai, Kawagoe 350-8585, Japan

* Correspondence: fujino048@toyo.jp




Figure S1. Time dependence of peak intensity at $m/z = 139$.