

**Table S1.** Distribution of minimum inhibitory concentration (MIC) of eight antimicrobial agents; MIC<sub>50</sub> and MIC<sub>90</sub> values for the studied *Streptococcus* spp. isolates from pigs (*n*=48).

Antimicrobial agent <sup>a</sup>	Number of Isolates with MIC (µg/mL) <sup>b</sup>												MIC <sub>50</sub>	MIC <sub>90</sub>
	0.003	0.004	0.006	0.008	0.012	<0.016	0.016	0.023	0.032	0.047	0.064	0.094		
PEN	2	16	13	4	3		6	3		1			0.006	0.016
AMC						8	18	13	1	5	3		0.016	0.047
CTX					1		8	8	13	6	5	7	0.032	0.094
	<b>0.25</b>	<b>0.38</b>	<b>0.5</b>	<b>0.75</b>	<b>1</b>	<b>1.5</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>8</b>	<b>12</b>		
CIP	3	9	18	16	1		1						0.5	0.75
GEN			3	7	10	10	9	4	5				1.5	3
	<b>0.064</b>	<b>0.094</b>	<b>0.125</b>	<b>0.19</b>	<b>0.25</b>	<b>0.38</b>	<b>0.5</b>	<b>0.75</b>	<b>1</b>	<b>1.5</b>	<b>2</b>	<b>&gt;256</b>		
ERY	1	6	11	6	4	5	2	9				4	0.19	0.75
	<b>0.125</b>	<b>0.19</b>	<b>0.25</b>	<b>0.38</b>	<b>0.5</b>	<b>0.75</b>	<b>1</b>	<b>1.5</b>	<b>3</b>	<b>8</b>	<b>12</b>	<b>&gt;32</b>		
SXT	6	10	3	7	4	6	2	4	2	1		3	0.38	3
	<b>0.094</b>	<b>0.125</b>	<b>0.19</b>	<b>0.25</b>	<b>0.38</b>	<b>0.5</b>	<b>4</b>	<b>12</b>	<b>16</b>	<b>24</b>	<b>32</b>	<b>48</b>		
DOX	2	3	5	10	2	2	1	3	11	5	2	2	0.5	24

<sup>a</sup> PEN – penicillin, AMC – amoxicillin/clavulanic acid, CTX – cefotaxime, CIP – ciprofloxacin, ERY – erythromycin, DOX – doxycycline, GEN – gentamicin, SXT – trimethoprim-sulfamethoxazole; <sup>b</sup> MIC range for tested antimicrobial agents.