

Supplementary materials.1. UPLC-ESI-MS/MS data of flavonoid ingredients characterized in the leaf extracts of *A. venetum* and *A. hendersonii*.

	Peak	RT (min)	UV λ_{max}	MS [m/Z]-	Formula	Tentative identification
<i>A. venetum</i>	1	10.113	202, 256, 355	464	C ₂₁ H ₂₀ O ₁₂	Hyperoside
	2	10.320	203, 256, 354	464	C ₂₁ H ₂₀ O ₁₂	Isoquercetin
	3	11.277	203, 256, 355	550	C ₂₄ H ₂₂ O ₁₅	Quercetin-3-O-(6-O-malonyl)-Galactoside
	4	11.420	203, 256, 355	550	C ₂₄ H ₂₂ O ₁₅	Quercetin-3-O-(6-O-malonyl)-glucoside
	5	12.693	203, 256, 355	505	C ₂₃ H ₂₂ O ₁₃	Quercetin-3-O-(6-O-acetyl)-galactoside
<i>A. hendersonii</i>	1	7.433	203, 256, 354	626	C ₂₇ H ₃₀ O ₁₇	Quercetin 3-sophoroside (Baimaside)
	2	10.450	203, 256, 354	464	C ₂₁ H ₂₀ O ₁₂	Isoquercetin
	3	11.383	203, 256, 355	549	C ₂₄ H ₂₂ O ₁₅	Quercetin-3-O-(6-O-malonyl)-Galactoside