

SUPPLEMENTARY INFORMATION

Bioavailability of Organosulfur Compounds after the Ingestion of Black Garlic by Healthy Humans

Alicia Moreno-Ortega^{1,2,*}, Gema Pereira-Caro^{1,2}, Iziar A. Ludwig^{3,†}, María-José Motilva^{3,‡} and José Manuel Moreno-Rojas^{1,2,*}

1 Department of Agroindustry and Food Quality, Andalusian Institute of Agricultural and Fisheries Research and Training (IFAPA), Alameda del Obispo, Avda. Menéndez-Pidal, 14004 Córdoba, Spain

2 Foods for Health Group, Maimonides Biomedical Research Institute of Cordoba (IMIBIC), 14004 Córdoba, Spain

3 Agrotecnio Center, XaRTA-TPV, Food Technology Department, Escola Tècnica Superior d'Enginyeria Agrària, University of Lleida, Avda. Alcalde Rovira Roure 191, 25198 Catalonia, Spain

* Correspondence: alicia.moreno.ortega@juntadeandalucia.es (A.M.-O.); josem.moreno.rojas@juntadeandalucia.es (J.M.M.-R.)

† Current address: Departamento de Ciencias de la Alimentación y Fisiología, Facultad de Farmacia y Nutrición, Universidad de Navarra, 31008 Pamplona, Spain.

‡ Current address: Instituto de Ciencias de la Vid y del Vino (ICVV) (Consejo Superior de Investigaciones Científicas-CSIC, Gobierno de La Rioja, UR), Finca “La Grajera”, Carretera de Burgos km 6, 26007 La Rioja, Spain.




Figure S1. HPLC-HRMS chromatograms of the identified *S*-Alk(en)yl-L-cysteine derivatives in urine after acute intake of Black Garlic.




Figure S2. HPLC-HRMS chromatograms of the identified γ -Glutamyl-S-alk(en)yl-L-cysteine derivatives in urine after acute intake of Black Garlic.