Cecal microbial diversity and metabolome reveal a reduction in growth due to oxidative stress caused by a low-energy diet in donkeys

Li L [1], Xiaoyu Guo [1], Yanli Zhao [1], Yongmei Guo [1], Binlin Shi [1], Yan Zhou [1], Yongwei Zhang [2] and Sumei Yan [1,*]

[1] Inner Mongolia Key Laboratory of Animal Nutrition and Feed Science, College of Animal Science, Inner Mongolia Agricultural University, Hohhot 010018, China; lily972021@163.com (L.L.); gxy_2594@163.com (X.G.); ylzhao2010@163.com (Y.Z.); ymguo2015@163.com (Y.G.); shibinlin@yeah.net (B.S.); 1454803209@qq.com (Y.Z.)

[2] Inner Mongolia Grassland Yulv Science and Technology Animal Husbandry Co., Ltd. Horinger County 011500, China. 1010142628@qq.com (Y.Z.)

* Correspondence: Correspondence: yansmimau@163.com



Figure S2. (A) PCA score plots of E2 vs. E1 in Pos. (B) PCA score plots of E2 vs. E1 in Neg. (C) OPLS-DA score plots of E2 vs. E1 in Pos. (D) OPLS-DA score plots of E2 vs. E1 in Neg. (E) Permutation test of OPLS-DA model for group E2 vs. E1 in Pos. (F) Permutation test of OPLS-DA model for group E2 vs. E1 in Neg. E1: low-energy diet; E2: high-energy diet.