antioxidants


MDPI

Supplementary Material


Isolation, Characterization, Antioxidant and Anticancer Activities of Compounds from *Erythrina caffra* Stem Bark Extract

Femi Olawale [1], Olusola Bodede [2], Mario Ariatti [1] and Moganavelli Singh [1,*]


[1] Nano-Gene and Drug Delivery Group, Discipline of Biochemistry, University of KwaZulu-Natal, Private Bag X54001, Durban 4000, South Africa; olawalefemi3@gmail.com (F.O.); ariattim@ukzn.ac.za (M.A.)
[2] Biodiscovery Centre, Department of Chemistry, University of Pretoria, Pretoria 0028, South Africa; olusolabodede@northeastern.edu
* Correspondence: singhm1@ukzn.ac.za; Tel.: +27-31-2607170




Supplementary Figure S1. Flow chart for the purification of anticancer compounds from *Erythrina caffra*.




Supplementary Figure S2a. ^{1}H NMR spectrum of compound 1.




Supplementary Figure S2b. ^{13}C NMR spectrum of compound 1.




Supplementary Figure S2c. HR-ESI-MS spectrum of compound 1.




Supplementary Figure S2d. ^{1}H NMR spectrum of compound 2




Supplementary Figure S2e. ^{13}C NMR spectrum of compound 2




Supplementary Figure S2f. HR-ESI-MS spectrum of compound 2




Supplementary Figure S2g. ^{1}H NMR spectrum of compound 3



Supplementary Figure S2h. ^{13}C NMR spectrum of compound 3



Supplementary Figure S2i. GC-MS spectrum of compound 3



Supplementary Figure S3. Quantitative analysis of multicaspase expression in HeLa and MCF-7 cells using flow cytometry.