

Effect of *Beauveria bassiana*-Seed Treatment on *Zea mays* L. Response Against *Spodoptera frugiperda*

Laiju Kuzhuppillymyal-Prabhakarankutty ¹, Fernando H. Ferrara-Rivero ², Patricia Tamez-Guerra ¹, Ricardo Gomez-Flores ¹, María Cristina Rodríguez-Padilla ¹ and María Julissa Ek-Ramos ^{1,*}

¹ Departamento de Microbiología e Inmunología-Laboratorio de Inmunología y Virología, Facultad de Ciencias Biológicas, Universidad Autónoma de Nuevo León, UANL, San Nicolás de los Garza, Nuevo León 66455, Mexico; laiju.kuzhuppillymyalpr@uanl.edu.mx (L.K.-P.); patricia.tamezgr@uanl.edu.mx (P.T.-G.); ricardo.gomezfl@uanl.edu.mx (R.G.-F.); cristina.rodriguezpd@uanl.edu.mx (M.C.R.-P.)

² Rancho La Verdad, Hacienda Ramírez, General Terán, Nuevo León 67405, México
fferrara@hexagonosmexicanos.com (F.H.F.-R.)

* Correspondence: maria.ekramos@uanl.edu.mx (M.J.E.-R.); Tel.: +52 (81) 8329 4115 ext. 6435, Fax: +52 (81) 8352 4212



Figure S1. 3rd instar *S. frugiperda* larva onto artificial diet.



Figure S2. *S. frugiperda* larva in *Z. mays* plant covered with a mesh cage.




Figure S3. Dimensions of Field trial 1.




Figure S4. Dimensions of Field trial 2.



Figure S5. Representative *S. frugiperda* larval instars found in the fields at the time of data collection.



Figure S6. *B. bassiana* treated plants showed more than two corn cobs per plant.