

Table S2. Primers used for qRT-PCR analysis.

Gene	Forward Primer	Reverse Primer
<i>Dazl</i>	ATGTCTGCCACAACCTCTGAG	CTGATTTCGGTTTCATCCATCCT
<i>Uchl1</i>	GATGCTGAACAAAGTGTTGGC	GGAGTTTCCGATGGTCTGCTT
<i>Utf1</i>	TGTCCCGGTGACTACGTCT	CCCAGAAGTAGCTCCGTCTCT
<i>Sycp3</i>	AGCCAGTAACCAGAAAATTGAGC	CCACTGCTGCAACACATTCATA
<i>Tex101</i>	ATCTTTCTTCTAATCGCCTCACG	GCTCAGCCTTTGAAGTCCAGT
<i>Pttg1</i>	TCTGATCCGCTGTACTCTCCT	AGGCGGCAATTCAACATCCA
<i>Acrv1</i>	TCAGCAACTTTCAAGCGAGTAT	CTCCTGAAGAGTGCTCACCTG
<i>Tssk1</i>	CTCAAGCGACGAGGCTACATC	ACCGCCACGTTGAACTTTAGG
<i>Spacal</i>	CCCGAGTCCGAAACCACAG	ACAACACACTTGGATTACCTC
<i>Tnp2</i>	TCACACCAGTAACCAGTGCAA	CCCTGAGCTACGCCTCTTAG
<i>Prm2</i>	ATGGTTTCGCTACCGAATGAGG	CTCCGCCTTCTGCATGACC
<i>Best1</i>	ACACAACACATTCTGGGTGC	CGCAAAGTACACACCTCATTCA
<i>Star</i>	ATGTTCTCTGCTACGTTCAAG	CCCAGTGCTCTCCAGTTGAG
<i>Hsd3b1</i>	TGGACAAAGTATTCCGACCAGA	GGCACACTTGCTTGAACACAG
<i>Cyp11a1</i>	AGGTCCTCAATGAGATCCCTT	TCCCTGTAAATGGGGCCATAC
<i>Cyp17a1</i>	GCCCAAGTCAAAGACACCTAAT	GTACCCAGGCGAAGAGAATAGA
<i>Hsd17b3</i>	AGGTTCTCGCAGCACCTTTTT	CATCGCCTGCTCCGGTAATC
<i>Nr5a1</i>	CCCAAGAGTTAGTGCTCCAGT	CTGGGCGTCCTTTACGAGG
<i>Insl3</i>	TCCTGGCTATGTCATTGC	TGTGGTCCTTGCTTACTG
<i>Il-1b</i>	GAAATGCCACCTTTTGACAGTG	TGGATGCTCTCATCAGGACAG
<i>Il-6</i>	AGTCCGGAGAGGAGACTTCA	ATTCCACGATTTCACAGAG
<i>Tnf</i>	CAGGCGGTGCCTATGTCTC	CGATCACCCGAAGTTCAGTAG
<i>Ccl2</i>	TTAAAAACCTGGATCGGAACCAA	GCATTAGCTTCAGATTTACGGGT
<i>Ccl7</i>	CCACATGCTGCTATGTCAAGA	ACACCGACTACTGGTGATCCT
<i>Nos2</i>	GTTCTCAGCCCAACAATACAAGA	GTGGACGGGTGCATGTCAC
<i>Il-4</i>	GGTCTCAACCCCCAGCTAGT	GCCGATGATCTCTCTCAAGTGAT
<i>Il-10</i>	CTTACTGACTGGCATGAGGATCA	GCAGCTCTAGGAGCATGTGG
<i>Ccl17</i>	TACCATGAGGTCACCTCAGATGC	GCACTCTCGGCCTACATTGG
<i>Gapdh</i>	AGGTCGGTGTGAACGGATTTG	TGTAGACCATGTAGTTGAGGTCA