

Table S1. Regression analysis of factors predicting clinical remission at T 12.

	Univariate Analysis			Multivariate Analysis		
	Odds Ratio	p-Value	95% IC	Odds Ratio	p-Value	95% IC
Male	1.44	0.57	0.41–5.07			
Age	0.99	0.48	0.95–1.02			
Body Mass Index	1.14	0.18	0.94–1.38			
Active smokers	0.50	0.48	0.07–3.38			
Crohn's disease	1.67	0.43	0.46–6.06			
5-aminosalicylic acid	2.14	0.24	0.59–7.77			
Topical corticosteroids	0.78	0.77	0.16–3.67			
Systemic corticosteroids	0.76	0.67	0.22–2.71			
Immunomodulators	0.80	0.83	0.10–6.32			
HBI T 0	0.94	0.63	0.75–1.20			
HBI T 6	0.65	0.07	0.40–1.04			
PMS T 0	0.59	0.09	0.32–1.09			
PMS T 6	0.47	0.02	0.25–0.89	0.59	0.11	0.31–1.12
C-RP T 0	0.47	0.25	0.13–1.69			
C-RP T 6	0.24	0.04	0.06–0.92	0.36	0.43	0.03–4.59
Hemoglobin T 0	2.14	0.04	1.04–4.43	1.78	0.50	0.33–9.49
Hemoglobin T 6	1.86	0.02	1.11–3.12	1.03	0.95	0.45–2.35
Fecal Calprotectin T 0	0.99	0.25	0.99–1.01			
Fecal Calprotectin T 6	0.99	0.08	0.99–1.00			
Vedolizumab concentrations T 6	1.02	0.03	1.002–1.03	1.01	0.36	0.99–1.04
Ulcerative colitis	1.01	0.80	0.99–1.02			