

Table S1. SS-OCT/SS-OCTA results stratified by duration of thalamic infarction.

	Group1(n=20)	Group 2(n=6)	Group 3(n=9)	<i>p</i> *	<i>p</i> 1	<i>p</i> 2	<i>p</i> 3
SVP, mm <sup>2</sup>	0.198 ± 0.022	0.203 ± 0.022	0.198 ± 0.035	0.502	<b>0.013</b>	<b>0.042</b>	0.576
ICP, mm <sup>2</sup>	0.163 ± 0.026	0.167 ± 0.016	0.175 ± 0.016	0.455	<b>0.004</b>	0.200	<b>0.023</b>
DCP, mm <sup>2</sup>	0.112 ± 0.023	0.118 ± 0.011	0.121 ± 0.024	0.498	0.128	0.980	0.153
RNFL, μm	16.94 ± 1.463	17.704 ± 1.165	17.173 ± 2.315	0.212	0.286	0.138	0.688
GCIPL, μm	61.667 ± 8.131	62.740 ± 5.620	62.161 ± 10.691	0.595	0.211	0.223	0.692

Group 1: < 1 month; Group 2: 1-6 months; Group 3: >6 months.

*p*\*value among the three groups;

*p*1: Group1 vs Group 2;

*p*2: Group1 vs Group 3;

*p*3: Group2 vs Group 3.

Values in bold indicate *p* < 0.05.