

# Longitudinal Coupling between Eating Disorder Psychopathology and Depression in Patients with Anorexia Nervosa and Bulimia Nervosa Treated with Enhanced Cognitive Behavior Therapy: A One-Year Follow-Up Study

Emanuele Cassioli, Eleonora Rossi, Michela Martelli, Francesca Arganini, Gabriele Giuranno, Serena Siviglia, Livio Tarchi, Marco Faldi, Giovanni Castellini \* and Valdo Ricca

Psychiatry Unit, Department of Health Sciences, University of Florence, Largo Brambilla 3, 50100 Florence, Italy

## Supplementary Materials

**Supplementary Table S1.** Characteristics of the sample at baseline and follow-up, divided by baseline diagnosis. Results of longitudinal analysis in patients are reported using age and BMI-adjusted unstandardized coefficients (time, group, and interaction fixed effects).

	Anorexia Nervosa		Bulimia Nervosa		Time Effect (b)	Group Effect (b)	Time*Group Interaction Effect (b)
	Baseline <i>T0</i> (n = 92)	Follow-up <i>T1</i> (n = 73)	Baseline <i>T0</i> (n = 65)	Follow-up <i>T1</i> (n = 48)			
Age (years)	24.41 ± 9.14	-	27.11 ± 10.86	-			
Education (years)	12.38 ± 3.09	-	13.52 ± 2.73	-			
Age of onset (years)	17.54 ± 4.03	-	17.97 ± 6.45	-			
EDE-Q Total Score	3.21 ± 1.64	2.29 ± 1.78	3.75 ± 1.44	2.73 ± 1.50	-1.10***	0.41	0.19
BDI	22.47 ± 11.59	14.00 ± 13.17	23.23 ± 11.24	13.62 ± 10.81	-9.10***	2.08	-0.36
SCL-90-R GSI	1.43 ± 0.78	1.07 ± 0.82	1.65 ± 0.68	1.13 ± 0.71	-0.42***	0.30	-0.06

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ ; BDI, Beck Depression Inventory; BMI, Body Mass Index; EDE-Q, Eating Disorders Examination Questionnaire; SCL-90-R GSI, Symptom Checklist-90-Revised Global Severity Index.