Supporting Information

Activated Carbon Ammonization: Effects of the Chemical Composition of the Starting Material and the Treatment Temperature

Silvia da C. Oliveira [1], Romulo C. Dutra [1], José J. L. León [1], Gesley A. V. Martins [1], Alysson M. A. Silva [2], Diana C. S. de Azevedo [3], Rafaelle G. Santiago [3], Daniel Ballesteros-Plata [4], Enrique Rodríguez-Castellón [4,*] and Marcos J. Prauchner [1,*]

[1] Institute of Chemistry, University of Brasilia, Campus Darcy Ribeiro, Brasilia, 70904-970, DF, Brazil; silviaoliveira.sco@gmail.com (S.d.C.O.); romulodutrac@gmail.com (R.C.D.); joselinares@unb.br (J.J.L.L.); gesley@unb.br (G.A.V.M.)
[2] Department of Mechanical Engineering, Faculty of Technology, University of Brasilia, Campus Darcy Ribeiro, Brasilia, 70904-970, DF, Brazil; alyssonmartins@unb.br
[3] Department of Chemical Engineering, University Federal of Ceará, Campus do Pici, Fortaleza, 60760-400, CE, Brazil; diana.azevedo@ufc.br (D.C.S.d.A.); rafaelle.santiago@gpsa.ufc.br (R.G.S.)
[4] Department of Inorganic Chemistry, Crystallography and Mineralogy, Inter-university Institute of Research in Biorefineries I3B, Faculty of Sciences, University of Málaga, 29071 Málaga, Spain; daniel.ballesteros@uma.es
* Correspondence: castellon@uma.es (E.R.-C.); marcosjp@unb.br (M.J.P.)




Figure S1. XPS survey spectra of the (a) unmodified AC WV and the samples resulting from its (b) complementary carbonization (WVC) and (c) nitrification (WVAc).




Figure S2. Binding Energies of XPS O 1*s* peaks for different functional oxygenated groups. (Based on reference 89 of the article).




Figure S3. XPS survey spectra of the ammonized samples.