

Risk Factors for Liver Decompensation and HCC in HCV-Cirrhotic Patients after DAAs: A Multicenter Prospective Study

Filomena Morisco, Alessandro Federico, Massimo Marignani, Mariarita Cannavò, Giuseppina Pontillo, Maria Guarino, Marcello Dallio, Paola Begini, Rosa G. Benigno, Flavia L. Lombardo and Tommaso Stroffolini

Table S1. Incidence rate (IR) per 100 person-years and relative rate (RR) of adverse events in SVR vs. no SVR cases (A), of Liver Decompensation and HCC in SVR according to Delta LSM (B) and Basal LSM value (C) in an Italian prospective study.

Characteristic						Any Event
Liver related death						
		IR (95% CI)		IR (95% CI)	RR (95% CI)	
A						
SVR	Yes		3.7 (2.9-4.8)	3.3 (1.3-8.2)		0.5 (0.2-1.0)
	SVR	No		9.0 (1.9-42.6)		12.3 (5.1-29.6)
			4.3 (1.1-17.2)			
Characteristic						
		Liver Decompensation		HCC		
		IR (95% CI)	RR (95% CI)	IR (95% CI)	RR (95% CI)	
B						
Delta LSM						
≥ 20%			0.5 (0.2-1.6)	0.3 (0.1-1.4)	2.3 (0.4-11.8)	
< 20%		1.2 (0.3-5.4)			0.8 (0.3-1.9)	
			0.6 (0.3-1.7)			
C						
Basal LSM value						
< 20 kPa			0.4 (0.1-1.1)	0.3 (0.1-0.9)	8.3 (2.3-30.2)	
≥ 20 kPa		5.5 (1.7-17.8)			2.5 (1.4-4.7)	
			2.2 (1.2-4.3)			