

A Simple Preoperative Blood Count to Stratify Prognosis in Isocitrate Dehydrogenase-Wildtype Glioblastoma Patients Treated with Radiotherapy Plus Concomitant and Adjuvant Temozolomide

Anne Clavreul, Jean-Michel Lemée, Gwénaëlle Soulard, Audrey Rousseau and Philippe Menei

Table S1. Relationship between hematological variables through Pearson's chi-squared test. Abbreviations: dNLR, derived neutrophil-to-lymphocyte ratio; LMR, lymphocyte-to-monocyte ratio; NLR, neutrophil-to-lymphocyte ratio; PLR, platelet-to-lymphocyte ratio; RBC, red blood cells; SII, systemic immune-inflammation index; SIRI, systemic inflammation response index; WBC, white blood cells.

	NLR		dNLR		Platelets		RBC	
	Pearson's r	P	Pearson's r	P	Pearson's r	P	Pearson's r	P
RBC	0.04	0.720	0.02	0.866	-0.22	0.046	1.00	/
WBC	0.26	0.017	0.36	< 0.001	0.20	0.062	-0.12	0.260
Neutrophils	0.38	< 0.001	0.39	< 0.001	0.23	0.033	-0.14	0.201
Lymphocytes	-0.20	0.067	-0.26	0.017	0.08	0.495	-0.15	0.169
Monocytes	-0.09	0.401	-0.13	0.225	0.16	0.146	-0.06	0.598
Platelets	0.05	0.654	0.08	0.489	1.00	/	-0.22	0.046
NLR	1.00	/	0.88	< 0.001	0.05	0.654	0.04	0.720
dNLR	0.88	< 0.001	1.00	/	0.08	0.489	0.02	0.866
LMR	-0.37	< 0.001	-0.38	< 0.001	-0.19	0.087	-0.03	0.792
PLR	0.30	0.005	0.36	< 0.001	0.22	0.046	0.07	0.507
SII	0.78	< 0.001	0.67	< 0.001	0.26	0.016	0.11	0.335
SIRI	0.52	< 0.001	0.49	< 0.001	0.10	0.385	-0.03	0.760