

Supplementary Files

Associations between CT-Based Preoperative Sarcopenia and Outcomes in Patients That Underwent Liver Resections

David Martin, Yaël Maeder, Kosuke Kobayashi, Michael Schneider, Joachim Koerfer, Emmanuel Melloul, Nermin Halkic, Martin Hübner, Nicolas Demartines, Fabio Becce and Emilie Uldry

Table S1. Clinical outcomes of patients that underwent major hepatectomies

Outcome	Overall (<i>n</i> = 190)	Non-sarcopenia (<i>n</i> = 73)	Sarcopenia (<i>n</i> = 117)	<i>p</i> -value
Length of stay (median, IQR)	12 (7–20)	11 (7–19)	13 (8–21)	0.196
30-day complications, (%)				
Any	118 (51)	51 (70)	67 (57)	0.092
Major	31 (16)	12 (16)	19 (16)	1.000
Bile leakage, (%)				0.122
A	9 (5)	7 (10)	2 (2)	
B	26 (14)	14 (19)	12 (10)	
C	3 (2)	-	3 (3)	0.532
Liver failure, (%)				
A	2 (1)	-	2 (2)	
B	4 (2)	1 (1)	3 (3)	
C	1 (1)	-	1 (1)	
30-day reoperation, (%)	24 (13)	6 (8)	16 (14)	0.352

IQR: interquartile range.