

Supplementary Table S1. Baseline Demographics of non-neoplastic versus neoplastic groups.

		Idiopathic (n=14)	Neoplastic causes (n=14)	Total
Sex	Female	12 (85.7%)	6 (42.9%)	18
	Male	2 (14.3%)	8 (57.1%)	10
Age at first symptoms (median and range in years)		9.3 (3.6–16)	11 (5–16.3)	9.7 (3.6–16.3)
Age at diagnosis of PST (median and range in years)		10.7 (3.8–16.2)	11.1 (5.2–16.5)	10.9 (3.8–16.5)
Presenting Symptoms	Polyuria/polydipsia	4	13	17
	Growth retardation	2	7	9
	Precocious puberty	7	1	8
	Delayed puberty	2	3	5
	Headache	2	3	5
	Visual problems	4	5	9
	Weight loss	1	2	3
	Lethargy	1	2	3
	Dizziness	0	1	1
	Poor appetite	2	1	3
	Decrease libido	0	1	1
Visual Problems	Visual field	1	1	2
	Visual acuity	0	0	0
	Papilledema	0	1	1
	Exotropia/diplopia	2	2	4
	Parinaud syndrome	0	1	1
	Raised intra-ocular pressure	0 1	1 0	1 1
	Amblyopia			
Biopsy performed	No	14 (100%)	3 (21.4%)	17
	Yes	0 (0%)	11 (78.6%)	11
Stalk thickness at initial MRI	Mild (<4mm)	11 (78.6%)	4 (28.6%)	15
	Moderate (4mm-<6.5mm)	3 (21.4%)	5 (35.7%)	8
	Severe (≥6.5mm)	0 (0%)	5 (35.7%)	5
Maximal stalk thickness during FU	<6.5mm	14 (100%)	2 (14.3 %)	16
	≥6.5mm	0 (0%)	12 (85.7 %)	12
Outcome	Alive	14	13	27
	Died	0	1	1