Supplemental Table S1: Survival of n = 142 patients with advanced pancreatic cancer receiving chemotherapy in dependence of an additional integrative therapeutic approach

	Median survival (months)	*1-year survival rate (%)*	*2-year survival rate (%)*	*5-year survival rate (%)*
Mistletoe and hyperthermia	18.9 [15.2; 24.5]	77.4 [62.9; 86.8]	37.3 [23.5; 51.0]	9.2 [2.6; 21.0]
Only mistletoe	11.2 [7.1; 14.2]	44.7 [29.9; 58.5]	12.5 [4.6; 24.4]	0 --
No mistletoe, no hyperthermia	8.6 [4.7; 15.4]	33.8 [15.4; 53.2]	9.7 [1.7; 26.3]	0 --

Survival times are given as months (median (range)) or percentage.

Supplemental Table S2: Survival (patients with advanced pancreatic cancer receiving chemotherapy) in dependence of the time of initiation of the integrative therapy (early vs late after diagnosis)

		Median survival (months)	*1-year survival rate (%)*	*2-year survival rate (%)*	*5-year survival rate (%)*
Mistletoe therapy	**early**	14.7 [7.1; 15.8]	56.5 [40.1; 70.0]	19.8 [8.8; 33.9]	0 --
	late	14.7 [12.8; 20.0]	65.7 [51.1; 76.8]	29.6 [17.7; 42.4	8.66 [2.5; 19.5]
Hyperthermia	**early**	16.0 [5.4; 22.6]	58.8 [32.5; 77.8]	26.1 [8.3; 48.6]	0 --
	late	23.5 [15.8; 26.4]	87.6 [70.2; 95.2]	43.4 [25.4; 60.1]	14.9 [4.2; 31.7]

Survival times are given as months (median (range)) or percentage.