

Symptom scales

← Lower is better

Fatigue

Other
Oslo/Viken



Difference (95% CI)

9.85 (8.59 , 11.12)
11.70 (9.56 , 13.83)

Nausea

Other
Oslo/Viken



2.75 (2.12 , 3.37)
3.05 (1.99 , 4.11)

Pain

Other
Oslo/Viken



2.87 (1.47 , 4.28)
4.50 (2.13 , 6.88)

Dyspnoea

Other
Oslo/Viken



1.28 (0.12 , 2.44)
2.39 (0.43 , 4.35)

Insomnia

Other
Oslo/Viken



4.25 (2.66 , 5.84)
6.37 (3.69 , 9.05)

Appetite loss

Other
Oslo/Viken



7.89 (6.77 , 9.01)
7.59 (5.69 , 9.48)

Constipation

Other
Oslo/Viken



3.93 (2.49 , 5.38)
4.75 (2.31 , 7.19)

Diarrhoea

Other
Oslo/Viken



-0.76 (-2.01 , 0.49)
-0.94 (-3.05 , 1.17)

Financial difficulties

Other
Oslo/Viken



2.83 (1.77 , 3.88)
2.88 (1.10 , 4.66)

0 10 20 30 40 50

Mean

Function scales

Higher is better →

Global quality of life

Other
Oslo/Viken



Difference (95% CI)

-9.29 (-10.40 , -8.17)
-10.31 (-12.19 , -8.43)

Physical functioning

Other
Oslo/Viken



-3.82 (-4.71 , -2.93)
-4.27 (-5.78 , -2.77)

Social functioning

Other
Oslo/Viken



-14.32 (-15.66 , -12.98)
-15.88 (-18.14 , -13.62)

Role functioning

Other
Oslo/Viken



-18.46 (-19.96 , -16.95)
-20.36 (-22.91 , -17.82)

Emotional functioning

Other
Oslo/Viken



-7.14 (-8.23 , -6.06)
-9.47 (-11.29 , -7.64)

Cognitive functioning

Other
Oslo/Viken



-3.99 (-5.07 , -2.91)
-3.77 (-5.59 , -1.95)

60 70 80 90

Mean

● Case ● Control