

Supplemental Table S2. Prevalence of co-infections of HPV5 with each indicated β -HPV genotype in the study population.

Genotype	Control	Obesity			AP			CRC		
	%	%	<i>p</i>	<i>OR</i> (95% CI)	%	<i>p</i>	<i>OR</i> (95% CI)	%	<i>p</i>	<i>OR</i> (95% CI)
HPV5-20	0	2.86	0.43	Inf (0.03-Inf)	0	1	0 (0-61.99)	0	0	0 (0-Inf)
HPV5-22	2.17	5.71	0.57	2.69 (0.13-164.24)	9.1	0.24	4.39 (0.22-270.74)	14	0.06	7.20 (0.87-336.78)
HPV5-96	0	5.71	0.18	Inf (0.25-Inf)	4.54	0.32	Inf (0.05-16.05)	8	0.12	Inf (0.62-Inf)
HPV5-100	0	2.86	0.43	Inf (0.03-Inf)	0	1	0 (0-Inf)	14	0.013	Inf (1.43-Inf)
HPV5-159	0	2.86	0.43	Inf (0.03-Inf)	0	1	0 (0-Inf)	14	0.013	Inf (1.43-Inf)

OR, odds ratio; CI, confidence interval; Inf., infinite.