


Supplementary Table S1. Comparison of 3-HPA bioconversion factors by *L. reuteri* strains

Strain	Glycerol (mM)	3-HPA (mM)	Yield (% mol/mol)	Reaction time (min)	Reference
<i>L. reuteri</i> ATCC 53608	400	235	58.8	45	[1]
<i>L. reuteri</i> ATCC 53608	200	170	85.0	120	[2]
<i>L. reuteri</i> 1063	250	140	56.0	120	[3]
<i>L. reuteri</i> NRRL B-14171	150	80	53.3	480	[4]
<i>L. reuteri</i> CG001	200	195	97.5	60	[5]
<i>L. reuteri</i> DSM 20016	250	180	72.0	60	[6]
<i>L. reuteri</i> BR201	500	378	75.6	60	This study



Supplementary Figure S1. Acrolein analysis HPLC chromatogram by DNPH derivatization. Acrolein-DNPH standard of 1 mg/L (a) and DNPH derivatized 3-HPA bioconversion sample (b).

## **References:**

1. Doleires, Y.; Beck, P.; Vollenweider, S.; Lacroix, C. Production of 3-Hydroxypropionaldehyde Using a Two-Step Process with *Lactobacillus reuteri*. *Appl. Microbiol. Biotechnol.* 2005, 68, 467–474.
2. Lüthi-Peng, Q.; Schärer, S.; Puhan, Z. Production and Stability of 3-Hydroxypropionaldehyde in *Lactobacillus reuteri*. *Appl. Microbiol. Biotechnol.* 2002, 60, 73–80.
3. Talarico, T.L.; Casas, I.A.; Chung, T.C.; Dobrogosz, W.J. Production and Isolation of Reuterin, a Growth Inhibitor Produced by *Lactobacillus reuteri*. *Antimicrob. Agents Chemother.* 1988, 32, 1854–1858.
4. Zamudio-Jaramillo, M.A.; Hernandez-Mendoza, A.; Robles, V.J.; Mendoza-Garcia, P.G.; Espinosa-De-Los-Monteros, J.J.; Garcia, H.S. Reuterin Production by *Lactobacillus reuteri* NRRL B-14171 Immobilized in Alginate. *J. Chem. Technol. Biot.* 2009, 84, 100–105.
5. Chen, G.; Yang, D.M.; Xiao, Y.Q.; Chen, H.W. Influence of Conditions on Reuterin Accumulation by the Resting Cell Biotransformation Process. *Chinese. J. Chem. Eng.* 2011, 19, 1023–1027.
6. Stevens, M.J.A.; Vollenweider, S.; Meile, L.; Lacroix, C. 1,3-Propanediol Dehydrogenases in *Lactobacillus reuteri*: Impact on Central Metabolism and 3-Hydroxypropionaldehyde Production. *Microb. Cell. Fact.* 2011, 10, 61.