


Supplementary Materials

L-Proline-Based Natural Deep Eutectic Solvents as Efficient Solvents and Catalysts for the Ultrasound-Assisted Synthesis of Aurones via Knoevenagel Condensation

Maria-Anna Karadendrou, Ioanna Kostopoulou, Vasiliki Kakokefalou, Andromachi Tzani and Anastasia Detsi *

Laboratory of Organic Chemistry, Department of Chemical Sciences, School of Chemical Engineering, National Technical University of Athens, Iroon Polytechniou 9, Zografou Campus, 15780 Athens, Greece;
mariannakaradendrou@mail.ntua.gr (M.-A.K.);
ioanna.th.kostopoulou@gmail.com (I.K.);
vickyfer@hotmail.gr (V.K.); atzani@mail.ntua.gr (A.T.)

* Correspondence: adetsi@chemeng.ntua.gr; Tel.: +30-2107724126






Figure S2. ¹³C NMR of NaDES Pro/Ox 1:1




Figure S3. ¹H NMR of NaDES Pro/Gly 1:2




Figure S4. ¹³C NMR of NaDES Pro/Gly 1:2






Figure S5. ¹H NMR of NaDES Pro/LA/W 1:2:2.5






Figure S8. ¹H NMR of recycled NaDES Pro/Gly 1:2 after run 7




Figure S9. ¹H NMR of NaDES Pro/Gly before use and after seven runs




Figure S10. ¹H NMR of compound 3a




Figure S11. ¹H NMR of compound 3b




Figure S12. ¹H NMR of compound 3c




Figure S13. ¹H NMR of compound 3d




Figure S14. ¹H NMR of compound 3e




Figure S15. ¹H NMR of compound 3f




Figure S16. ¹H NMR of compound 3g




Figure S17. ¹H NMR of compound 3h




Figure S18. ¹H NMR of compound **3i**