

MIL-88-Derived N and S Co-Doped Carbon Materials with Supplemental FeS_x to Enhance the Oxygen Reduction Reaction Performance

Yu Liu ^{1,2,†}, Yinghao Xu ^{2,3,†}, He Wang ^{2,3}, Jia Zhang ⁴, Haiyan Zhao ⁵, Li Chen ⁶, Ling Xu ^{1,*}, Yan Xie ^{2,*} and Jiahui Huang ²

¹ College of Chemistry and Materials Science, Inner Mongolia Minzu University, Inner Mongolia, Tongliao 028000, China; yuliu@dicp.ac.cn

² Dalian National Laboratory for Clean Energy, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Zhongshan Road 457, Dalian 116023, China; yinghao.xu@uzh.ch (Y.X.); hw122@ic.ac.uk (H.W.); jiahuihuang@dicp.ac.cn (J.H.)

³ Dalian Leicester Institute, Dalian University of Technology, Dalian 116024, China

⁴ State Key Laboratory of Catalysis, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Zhongshan Road 457, Dalian 116023, China; zhangjia610@dicp.ac.cn

⁵ Liaoning Key Laboratory of Plasma Technology, School of Physics and Materials Engineering, Dalian Minzu University, Liaoning, Dalian 116600, China; zhaohaiyan@dlmu.edu.cn

⁶ Shanghai Key Laboratory of Green Chemistry and Chemical Processes, School of Chemistry and Molecular Engineering, East China Normal University, Shanghai 200062, China; chenli@chem.ecnu.edu.cn

* Correspondence: tlxuling1979@163.com (L.X.); yanxie@dicp.ac.cn (Y.X.)

† These authors contributed equally to this work.




Figure S1. SEM images of (a) and (b) MIL-88-850.




Figure S2. EDX of FeNSC-850.




Figure S3. XRD patterns of (a) MIL-88-850, (b) FeNSC-750, (c) FeNSC-800 and (d) FeNSC-900.




Figure S4. Raman spectra of (a) FeNSC-850 and (b) MIL-88-850.




Figure S5. The survey spectra of (a) MIL-88-850 and (b) FeNSC-850.




Figure S6. N 1s XPS spectra of (a) MIL-88-850, (b) FeNSC-750, (c) FeNSC-800 and (d) FeNSC-900.