

Supplementary Material

Enhanced photoelectrochemical activity of TiO₂ nanotubes decorated with lanthanide ions for hydrogen production

Hyekyung Cho^{1,2}, Hyunku Joo¹, Hansung Kim², Ji-Eun Kim^{1,3},
Kyoung-Soo Kang¹, Hyeonmin Jung^{1,4} and Jaekyung Yoon^{1,*}

¹ Korea Institute of Energy Research, Hydrogen Research Department, 152 Gajeong-ro, Daejeon 34129, Republic of Korea; chk2946@gmail.com (H.C); hkjoo@kier.re.kr (H.Joo); cheeeun12@kier.re.kr (J.-E.K.); kskang@kier.re.kr (K.-S.K); nbc13@kier.re.kr (H.Jung)²

Department of Chemical and Biomolecular Engineering, Yonsei University, 50 Yonsei-ro, Seoul 03722, Republic of Korea; elchem@yonsei.ac.kr (H.K);

³ Department of Chemical and Biological Engineering, Korea University, 145 Anam-ro, Seoul 02841, Republic of Korea

⁴ Graduate School of Energy Science and Technology, Chungnam National University, 99 Daehak-ro, Daejeon 34134, Republic of Korea

* Correspondence: jyoona@kier.re.kr; Tel.: +82-42-860-3473

***Corresponding author.**

* Dr. Jaekyung Yoon: Tel.: +82-42-860-3473; E-mail: jyoona@kier.re.kr.




Figure S1. Schematic diagram of the photoelectrochemical cell.