Surface Alkali-Modified Nano-CeO_2 for Atmospherically Catalytic Polycondensation of CO_2 and Diol

Wenbin Zhong [1], Rongbin Xiao [1], Sheng Huang [2], Min Xiao [2], Shuanjin Wang [2], Dongmei Han [1,2]* and Yuezhong Meng [1,2,3,4]*

[1] School of Chemical Engineering and Technology, Sun Yat-sen University, Guangzhou 510275, China; zhongwb6@mail2.sysu.edu.cn (W.Z.); xiaorb158@163.com (R.X.)
[2] The Key Laboratory of Low-Carbon Chemistry &Energy Conservation of Guangdong Province/State Key Laboratory of Optoelectronic Materials and Technologies, School of Materials Science and Engineering, Sun Yat-sen University, Guangzhou 510275, China; huangsh47@mail.sysu.edu.cn (S.H.); stsxm@mail.sysu.edu.cn (M.X.); wangshj@mail.sysu.edu.cn (S.W.)
[3] Institute of Chemistry, Henan Academy of Sciences, Zhengzhou 45000, China
[4] College of Chemistry, Zhengzhou University, Zhengzhou 450001, China
* Correspondence: handongm@mail.sysu.edu.cn (D.H.); mengyzh@mail.sysu.edu.cn (Y.M.)




Figure S1. GPC curves of the crude product (black), purified product (green) after 24 h of reaction using CeO_2-11 as catalyst and HDO monomer (red).




Figure S2. ^{1}H NMR of purified oligomeric diols after catalyzed from CeO_2-11 for 24 h.




Figure S3. MALDI-TOF spectrum of the purified product after 24 h of reaction using CeO_2-11 as catalyst.