

Supplementary Materials

Transmembrane and tetratricopeptide repeat containing 4 is a novel diagnostic marker for prostate cancer with high specificity and sensitivity

Rania Makboul¹, Islam F. Abdelkawi², Dalia M. Badary¹, Mahmoud R.A. Hussein¹, John S. Rhim³, Eman A. Toraih⁴, Mourad Zerfaoui⁴, Zakaria Y. Abd Elmageed^{5#}

Departments of Pathology¹ and Urology², Faculty of Medicine, Assiut University, Assiut 71111, Egypt;

Department of Surgery³, Uniformed Services University of the Health Sciences, Bethesda, MD 20814;

Department of Surgery⁴, Tulane University School of Medicine, 1430 Tulane Avenue, New Orleans, LA 70112;

Department of Pharmacology⁵, Edward Via College of Osteopathic Medicine, University of Louisiana at Monroe, Monroe, LA 71203, USA

Figure S1. Expression of TMTC4 in a large panel of PCa cells. Protein cell lysates were collected from C4-2B, Du-145, PC-3, E006AA, RC77 T/E, E006A-hT, RC77 N/E, RWPE-1 and BPH1 cells and the expression of cellular TMTC4 was evaluated by Western blot analysis. Membranes were incubated with TMTC4 primary antibody and GAPDH was used as a loading protein.

