

The mechanoreception in *Drosophila melanogaster* oocyte under modeling micro- and hypergravity

Irina V. Ogneva^{1,2*}

¹ Cell Biophysics Laboratory, State Scientific Center of the Russian Federation Institute of Biomedical Problems of the Russian Academy of Sciences, 76a, Khoroshevskoye shosse, 123007 Moscow, Russia;

² Medical and biological physics department, I. M. Sechenov First Moscow State Medical University, 8-2 Trubetskaya Street, 119991, Moscow, Russia;

*Correspondence: iogneva@yandex.ru, iogneva@imbp.ru; Tel.: +7(499)195-63-98

Table S1. The stiffness (in pN/nm) of the *Drosophila melanogaster* oocytes under simulated micro- and hypergravity without or with addition agents.

Exposure time		0 min	30 min	90 min	210 min
Group and conditions of pre-incubation					
No pre-incubation	C	30.7 ± 1.0	30.8 ± 1.4	31.2 ± 1.2	29.6 ± 0.7
	s µg	30.4 ± 1.4	30.6 ± 1.3	25.5 ± 1.2*↓	24.8 ± 1.2*↓
	hg	31.4 ± 1.0	26.7 ± 1.8*↓	27.0 ± 2.4*↓	29.5 ± 1.4
Pre-incubation in 0.65% NaCl without any agents 3 hours, RT	C	31.0 ± 1.4	31.5 ± 0.9	31.3 ± 1.8	30.6 ± 1.6
	s µg	30.8 ± 1.5	31.1 ± 1.6	25.8 ± 1.8*↓	26.2 ± 1.4*↓
	hg	30.6 ± 1.1	26.7 ± 1.7*↓	26.3 ± 1.4*↓	31.2 ± 1.1
Pre-incubation in 0.65% NaCl + 2 µM cytochalasine B 3 hours, RT	C	30.4 ± 1.2	31.0 ± 1.6	30.5 ± 2.1	30.0 ± 1.1
	s µg	30.9 ± 1.4	31.5 ± 1.3	26.0 ± 1.2*↓	26.3 ± 1.8*↓
	hg	30.4 ± 1.2	26.7 ± 2.1*↓	26.3 ± 1.4*↓	30.8 ± 1.6
Pre-incubation in 0.65% NaCl + 10 µM colchicine 3 hours, RT	C	31.1 ± 2.1	31.5 ± 1.1	30.4 ± 1.5	30.4 ± 1.0
	s µg	31.1 ± 1.4	30.4 ± 1.2	25.8 ± 0.7*↓	26.0 ± 1.4*↓
	hg	31.5 ± 1.7	26.6 ± 2.0*↓	26.0 ± 1.2*↓	25.8 ± 0.9*↓
Pre-incubation in 0.65% NaCl + 40mM acrylamide 3 hours, RT	C	26.4 ± 0.8*↓	27.0 ± 1.4*↓	26.6 ± 1.4*↓	25.8 ± 1.2*↓
	s µg	26.2 ± 0.9*↓	27.1 ± 1.2*↓	25.9 ± 0.8*↓	26.7 ± 1.1*↓
	hg	26.7 ± 1.2*↓	25.9 ± 1.2*↓	26.2 ± 1.2*↓	25.3 ± 1.2*↓
Pre-incubation in 0.65% NaCl + 100nM calyculin A 3 hours, RT	C	36.1 ± 1.8*↑	35.7 ± 1.6*↑	35.6 ± 1.0*↑	35.3 ± 1.3*↑
	s µg	34.7 ± 1.6*↑	35.9 ± 1.5*↑	30.1 ± 1.2	30.2 ± 1.7
	hg	35.3 ± 1.4*↑	28.2 ± 2.0&↓	30.5 ± 1.1	30.4 ± 2.0

C – control, s µg – simulated microgravity, hg – hypergravity at the 2g level. RT – room temperature. * – p < 0.05 in comparison with the according control group without any agents, & – p < 0.1 in comparison with the according control group without any agents. ↑ – increase relative to the reference value. ↓ – decrease relative to the reference value.

Table S2. The cellular respiration (in pmol O₂·ml⁻¹·min⁻¹·oocyte⁻¹) of the *Drosophila melanogaster* oocytes under simulated micro- and hypergravity.

Group \ Exposure time		0 min	30 min	90 min	210 min
V0	C	11.7 ± 1.0	12.0 ± 0.4	11.2 ± 0.8	10.7 ± 1.6
	s µg	12.1 ± 0.4	12.2 ± 0.4	9.7 ± 0.4	10.3 ± 1.2
	hg	12.2 ± 1.6	12.2 ± 0.6	10.2 ± 1.0	9.1 ± 0.8
Vglu+mal	C	16.6 ± 1.2	16.6 ± 1.3	16.0 ± 1.8	16.6 ± 1.3
	s µg	17.7 ± 2.0	15.7 ± 0.5	14.4 ± 1.4	17.0 ± 1.6
	hg	15.8 ± 2.1	16.9 ± 2.0	14.5 ± 0.7	13.7 ± 1.3
Vmax	C	22.6 ± 1.8	21.3 ± 1.6	21.3 ± 0.9	21.6 ± 2.2
	s µg	20.8 ± 1.3	20.8 ± 1.2	20.9 ± 1.0	18.9 ± 1.9
	hg	22.6 ± 1.3	22.5 ± 1.0	19.0 ± 1.0	20.8 ± 1.3
V(II)	C	23.6 ± 1.3	24.2 ± 2.1	23.4 ± 2.0	22.7 ± 1.9
	s µg	22.9 ± 2.6	26.6 ± 2.6	22.9 ± 1.1	22.3 ± 2.2
	hg	22.8 ± 1.2	24.0 ± 1.2	22.7 ± 0.4	21.7 ± 1.1
V(IV)	C	265 ± 16	265 ± 22	277 ± 23	259 ± 20
	s µg	284 ± 21	278 ± 24	270 ± 12	262 ± 18
	hg	297 ± 24	287 ± 20	266 ± 21	264 ± 28

C – control, s µg – simulated microgravity, hg – hypergravity at the 2g level. V0 – basal oxygen absorption. Vglu+mal – oxygen absorption after supplement 10mM glutamate+5mM malate (substrates of the complex I respiratory chain). Vmax – maximal oxygen absorption after supplement 2mM ADP. V(II) – oxygen absorption after 0.5 µM rotenone implementation (inhibitor of the complex I respiratory chain) and 10 mM succinate addition (substrate of the complex II respiratory chain). V(IV) – oxygen absorption after 5 µM antimycin A implementation (inhibitor of the complex III respiratory chain) and 0.5 mM TMPD + 2 mM ascorbate addition (artificial substrates of the complex IV respiratory chain).