

Supplementary Materials

A New Synthetic Curcuminoid Displays Antitumor Activities in Metastasized Melanoma

Leonard Kaps ^{1,*†}, Adrian Klefenz ^{1,†}, Henry Traenckner ¹, Paul Schneider ¹, Ion Andronache ², Rainer Schobert ³, Bernhard Biersack ^{3,*} and Detlef Schuppan ^{1,4,*}

¹ Institute of Translational Immunology, University Medical Center, Johannes Gutenberg University Mainz, Germany; leonard.kaps@unimedizin-mainz.de, detlef.schuppan@unimedizin-mainz.de

² Research Center for Integrated Analysis and Territorial Management, University of Bucharest, Romania; ion.andronache@geo.unibuc.ro

³ Organic Chemistry 1, University Bayreuth, Germany; rainer.schobert@uni-bayreuth.de, bernhard.biersack@yahoo.com

⁴ Division of Gastroenterology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA

* Correspondence: leonard.kaps@unimedizin-mainz.de (L. K.), bernhard.biersack@yahoo.com (B. B.), detlef.schuppan@unimedizin-mainz.de (D. S.)

† These authors contributed equally to this work.




Figure S1. (A) Safety laboratory parameters of sera from (un)treated and healthy mice; (B) Biometric data of mice at sacrifice.