

Table S1. Distribution of psychopathological diagnoses between the first quarantine period, first reference period, second quarantine period, and the second reference period.

Diagnosis	Distribution among Admitted Patients			
	First Quarantine Period (March to June 2020)	First Reference Period (March to June 2019)	Second Quarantine Period (October 2020 to January 2021)	Second Reference Period (October 2019 to January 2020)
	# (%)	# (%)	# (%)	# (%)
Psychosis	14 (16.7)	21 (19.4)	20 (16.0)	21 (20.6)
Mood disorders	49 (58.3)	48 (44.4)	85 (68.0)	50 (49.0)
Post-traumatic stress disorder	10 (11.9)	16 (14.8)	13 (10.4)	12 (11.8)
Behavior disorders	23 (13.1)	11 (21.3)	7 (5.6)	19 (18.6)

Table S2. Distribution of self-injurious behaviors between the first quarantine period, first reference period, second quarantine period, and the second reference period.

Self-Injurious Behaviors	Distribution among Admitted Patients			
	First Quarantine Period (March to June 2020)	First Reference Period (March to June 2019)	Second Quarantine Period (October 2020 to January 2021)	Second Reference Period (October 2019 to January 2020)
	# (%)	# (%)	# (%)	# (%)
Non-suicidal self-injurious behaviors	45 (56.3)	37 (46.3)	71 (47.3)	40 (47.6)
Suicidal ideation	24 (30)	21 (26.3)	56 (37.3)	32 (38.1)
Suicide attempts	11 (13.8)	22 (27.5)	23 (15.3)	12 (14.3)